Antiderivatives and Riemann Sums

ClipSAT Printable Practice Suite (clipsat.org) • Topic 6.01

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 [Foundational] • Target: Basic antiderivatives

Find the general antiderivative of each function:

a $f(x) = x^4$


b $f(x) = \frac{1}{x}$

c $f(x) = e^x$

d $f(x) = \sec^2 x$

2 [Foundational] • Target: Basic antiderivatives

Find $\int (3x^2 - 4x + 5)\,dx$.



3 [Foundational] • Target: Basic antiderivatives

Find the particular antiderivative $F(x)$ of $f(x) = 6x^2 - 4$ satisfying $F(1) = 5$.

4 [Foundational] • Target: Riemann sums: left, right, and midpoint

Approximate $\int_0^4 x^2\,dx$ using a Left Riemann Sum with $n = 4$ equal subintervals.


5 [Foundational] • Target: Riemann sums: left, right, and midpoint

Approximate $\int_0^4 x^2\,dx$ using a Right Riemann Sum with $n = 4$ equal subintervals.

© ClipSAT — clipsat.org | All Rights Reserved

6 [Foundational] • Target: Riemann sums: left, right, and midpoint

Approximate $\int_0^4 x^2\,dx$ using a Midpoint Riemann Sum with $n = 4$ equal subintervals.



7 [Foundational] • Target: The Trapezoidal Rule


Use the Trapezoidal Rule with $n = 4$ to approximate $\int_0^4 x^2\,dx$.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

© ClipSAT — clipsat.org | All Rights Reserved

8 [Intermediate] • Target: The Trapezoidal Rule

A table gives $f(0) = 2$, $f(2) = 5$, $f(4) = 9$, $f(6) = 8$. Use the Trapezoidal Rule with these unequal-width data (widths of 2 each) to approximate $\int_0^6 f(x)\,dx$.




9 [Intermediate] • Target: Sums as limits; the definite integral

Explain how the definite integral $\int_a^b f(x)\,dx$ is defined as a limit of Riemann sums, and write the limit expression using sigma notation.

© ClipSAT — clipsat.org | All Rights Reserved

10 [Intermediate] • Target: Sums as limits: the definite integral

Determine whether the Left Riemann Sum overestimates or underestimates $\int_0^4 x^2\,dx$, and explain why using the shape of the graph.



11 [Intermediate] • Target: Riemann sums from a table of values

A table gives $f(0) = 3$, $f(1) = 5$, $f(2) = 8$, $f(3) = 10$, $f(4) = 9$. Use a Right Riemann Sum with 4 subintervals of width 1 to approximate $\int_0^4 f(x)\,dx$.

12 [Intermediate] • Target: Antiderivatives of trigonometric and exponential functions

Find $\int (3\sin x + 2e^x)\,dx$.


13 [Intermediate] • Target: Antiderivatives of trigonometric and exponential functions

Find the particular antiderivative $F(x)$ of $f(x) = \sec^2 x$ satisfying $F(0) = 2$.

© ClipSAT — clipsat.org | All Rights Reserved

14 **[Intermediate]** • **Target: Visualizing a Riemann sum approximation**

The graph shows $f(x) = x^2$ on $[0, 4]$. Use the graph to explain why the Right Riemann Sum $R_4 = 30$ (found earlier) overestimates the true area, and why the Left Riemann Sum $L_4 = 14$ underestimates it.



Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)


15 **[Challenging]** • **Target: An applied antiderivative problem** • **Cross-Lesson: 6.02 The Fundamental Theorem of Calculus**

A factory's production rate is $r(t) = 100 + 20t$ units/day, where t is days since a process change. Find the total production function $P(t)$ given $P(0) = 0$, then find the total production over the first 10 days.

© ClipSAT — clipsat.org | All Rights Reserved

16 [Challenging] • Target: More antiderivative practice • Cross-Lesson: 7.01 Differential Equations and Slope Fields

Find the general antiderivative of $f(x) = 4x^3 - \frac{2}{x^2} + \cos x$.



17 [Challenging] • Target: More antiderivative practice • Cross-Lesson: 8.01 Area Between Curves

Find the particular antiderivative $F(x)$ of $f(x) = x^2 - 3$ satisfying $F(0) = 4$.

18 [Challenging] • Target: Comparing Riemann sums to the exact integral • Cross-Lesson: 8.02 Volumes by the Disc and Washer Methods

For $\int_0^4 x^2\,dx$, the exact value is $\frac{64}{3} \approx 21.33$. Rank the four approximations found earlier — $L_4 = 14$, $R_4 = 30$, $M_4 = 21$, $T_4 = 22$ — from closest to farthest from the exact value, and explain which type of approximation (midpoint or trapezoidal) tends to be more accurate for a smooth curving function.

19 [Challenging] • Target: Riemann sums from a table of values • Cross-Lesson: 1.01 Limits Graphically and Numerically

Approximate $\int_0^7 5x^2\,dx$ using a Right Riemann Sum with $n = 7$ equal subintervals.

20 [Challenging] • Target: Antiderivatives of trigonometric and exponential functions • Cross-Lesson: 1.02 Continuity and the Intermediate Value Theorem

Approximate $\int_0^6 x^2\,dx$ using a Right Riemann Sum with $n = 6$ equal subintervals.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

© ClipSAT — clipsat.org | All Rights Reserved

21 [Foundational] • Target: Basic antiderivatives

Find C in the particular antiderivative $F(x) = x^3 - 7x + C$ of $f(x) = 3x^2 - 7$, satisfying $F(-1) = -8$.

a. -8

b. -12

c. -14

d. -16

22 [Foundational] • Target: Riemann sums: left, right, and midpoint

Find C in the particular antiderivative $F(x) = -2x^3 - 5x + C$ of $f(x) = -6x^2 - 5$, satisfying $F(2) = -9$.

a. 17


b. -9

c. 15

d. 19

23 [Foundational] • Target: The Trapezoidal Rule

Find C in the particular antiderivative $F(x) = 2x^3 + 9x + C$ of $f(x) = 6x^2 + 9$, satisfying $F(0) = 9$.



a. 7

b. 9

c. 10

d. 11

24 [Foundational] • Target: Sums as limits: the definite integral

Find C in the particular antiderivative $F(x) = 2x^3 + x + C$ of $f(x) = 6x^2 + 1$, satisfying $F(3) = -8$.

a. -65

b. -8

c. -67

d. -63

© ClipSAT — clipsat.org | All Rights Reserved

25 [Foundational] • Target: Riemann sums from a table of values

Find C in the particular antiderivative $F(x) = x^3 + 5x + C$ of $f(x) = 3x^2 + 5$, satisfying $F(3) = 4$.

a. 4

b. -36

c. -38

d. -40

26 [Foundational] • Target: Antiderivatives of trigonometric and exponential functions

Find C in the particular antiderivative $F(x) = -2x^3 + 6x + C$ of $f(x) = -6x^2 + 6$, satisfying $F(-3) = -8$.

a. -8

b. -46

c. -44

d. -42

27 [Foundational] • Target: Visualizing a Riemann sum approximation

Find C in the particular antiderivative $F(x) = -x^3 + C$ of $f(x) = -3x^2$, satisfying $F(2) = -1$.

a. 9

b. 5


c. -1

d. 7

© ClipSAT — clipsat.org | All Rights Reserved

28 [Foundational] • Target: An applied antiderivative problem

Approximate $\int_0^4 5x^2\,dx$ using a Right Riemann Sum with $n = 4$ equal subintervals.



a. 150

b. 145

c. 170

d. 70

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 [Intermediate] • Target: More antiderivative practice

Approximate $\int_0^6 2x^2\,dx$ using a Right Riemann Sum with $n = 6$ equal subintervals.

a. 194

b. 110

c. 180

d. 182

30 [Intermediate] • Target: Comparing Riemann sums to the exact integral

Approximate $\int_0^3 5x^2\,dx$ using a Right Riemann Sum with $n = 3$ equal subintervals.

a. 70

b. 85

c. 65

d. 25

© ClipSAT — clipsat.org | All Rights Reserved

31 [Intermediate] • Target: Basic Antiderivatives & Power Rule for Integration

Approximate $\int_0^4 3x^2\,dx$ using a Right Riemann Sum with $n = 4$ equal subintervals.

a. 42

b. 90

c. 102

d. 87

32 [Intermediate] • Target: Left and Right Riemann Sums

Approximate $\int_0^5 5x^2\,dx$ using a Right Riemann Sum with $n = 5$ equal subintervals.

a. 270


b. 275

c. 150

d. 300

33 [Intermediate] • Target: Midpoint & Trapezoidal Approximations

Approximate $\int_0^5 2x^2\,dx$ using a Right Riemann Sum with $n = 5$ equal subintervals.



a. 110

b. 120

c. 60

d. 108

© ClipSAT — clipsat.org | All Rights Reserved

34 [Intermediate] • Target: Limit of a Riemann Sum & Definite Integrals

Approximate $\int_0^6 f(x)\,dx$ using the Trapezoidal Rule (equal width 2), given $f(0) = 7$, $f(2) = 12$, $f(4) = 9$, $f(6) = 3$.

a. 54

b. 52

c. 62

d. 50

35 [Intermediate] • Target: Basic antiderivatives

Approximate $\int_0^3 f(x)\,dx$ using the Trapezoidal Rule (equal width 1), given $f(0) = 4$, $f(1) = 7$, $f(2) = 12$, $f(3) = 2$.

a. 22

b. 25

c. 23

d. 21

36 [Intermediate] • Target: Riemann sums: left, right, and midpoint

For $f(x) = 4x^2$ on $[0, 5]$, determine whether the Left and Right Riemann Sums overestimate or underestimate $\int_0^5 f(x)\,dx$.

a. Both Left and Right Riemann Sums underestimate.

b. Both Left and Right Riemann Sums overestimate.

c. Left Riemann Sum overestimates, and Right Riemann Sum underestimates.

d. Left Riemann Sum underestimates, and Right Riemann Sum overestimates.

37 [Intermediate] • Target: The Trapezoidal Rule

For $f(x) = 3x^2$ on $[0, 3]$, determine whether the Left and Right Riemann Sums overestimate or underestimate $\int_0^3 f(x)\,dx$.

a. Left Riemann Sum overestimates, and Right Riemann Sum underestimates.

b. Both Left and Right Riemann Sums overestimate.

c. Left Riemann Sum underestimates, and Right Riemann Sum overestimates.

d. Both Left and Right Riemann Sums underestimate.

© ClipSAT — clipsat.org | All Rights Reserved

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Sums as limits: the definite integral • Cross-Lesson: 8.02 Volumes by the Disc and Washer Methods

For $f(x) = -2x^2 + 8$ on $[0, 2]$, determine whether the Left and Right Riemann Sums overestimate or underestimate $\int_0^2 f(x)\,dx$.

a. Both Left and Right Riemann Sums underestimate.

b. Both Left and Right Riemann Sums overestimate.

c. Left Riemann Sum overestimates, and Right Riemann Sum underestimates.

d. Left Riemann Sum underestimates, and Right Riemann Sum overestimates.

39 [Challenging] • Target: Riemann sums from a table of values • Cross-Lesson: 1.01 Limits Graphically and Numerically

A factory's production rate is $r(t) = 187 + 10t$ units/day, where t is days since a process change. Find the total production $P(15)$ given $P(0) = 0$.

a. 2805

b. 3920

c. 3930

d. 3940

40 [Challenging] • Target: Antiderivatives of trigonometric and exponential functions • Cross-Lesson: 1.02 Continuity and the Intermediate Value Theorem

A factory's production rate is $r(t) = 86 + 18t$ units/day, where t is days since a process change. Find the total production $P(14)$ given $P(0) = 0$.

a. 2968

b. 1204

c. 2950

d. 2986

© ClipSAT — clipsat.org | All Rights Reserved

41 [Challenging] • Target: Basic Antiderivatives & Power Rule for Integration • Cross-Lesson: 2.01 The Definition of the Derivative

Find $\int (\frac{3}{x} + \frac{1}{x^2})dx$ for $x > 0$.

a. $3\ln x - \frac{1}{x} + C$

b. $3\ln x + \frac{1}{x} + C$

c. $-\frac{3}{x^2} - \frac{2}{x^3} + C$

d. $3\ln x - \frac{1}{2x} + C$

42 [Challenging] • Target: Left and Right Riemann Sums • Cross-Lesson: 2.02 Product, Quotient, and Trigonometric Derivatives

Approximate $\int_0^4 x^2 dx$ using a Left Riemann sum with $n = 4$ equal subintervals.

a. 14

b. 30

c. 21.33

d. 16

43 [Challenging] • Target: Left and Right Riemann Sums • Cross-Lesson: 3.01 The Chain Rule

If $f(x)$ is strictly increasing on $[a, b]$, what is the relationship between Left and Right Riemann sums and the true integral value I?

a. Left Sum $< I <$ Right Sum

b. Right Sum $< I <$ Left Sum

c. Left Sum $=$ Right Sum

d. Cannot be determined without concavity

© ClipSAT — clipsat.org | All Rights Reserved

44 [Challenging] • Target: Midpoint & Trapezoidal Approximations • Cross-Lesson: 3.02 Implicit Differentiation and Inverse Functions

If $f''(x) > 0$ (concave up) on $[a, b]$, how does the Trapezoidal Rule approximation compare to the exact integral?

a. Trapezoidal Rule overestimates the true value

b. Trapezoidal Rule underestimates the true value

c. Trapezoidal Rule is exact

d. Depends on whether f is increasing

45 [Challenging] • Target: Limit of a Riemann Sum & Definite Integrals • Cross-Lesson: 4.01 Rates of Change and Rectilinear Motion

Which definite integral is equal to $\lim_{n\to\infty} \sum_{i=1}^{n} (1 + \frac{2i}{n})^3 \frac{2}{n}$?

a. $\int_1^3 x^3 dx$

b. $\int_0^2 x^3 dx$

c. $\int_1^2 (1 + x)^3 dx$

d. $\int_0^1 (1 + 2x)^3 dx$

© ClipSAT — clipsat.org | All Rights Reserved