

z-Scores & Linear Transformations of Data

ANSWER KEY



Computing z-scores

- 1 A test has mean 70 and standard deviation 5. Find the z-score for a student who scored 82.

$$z = \frac{x - \mu}{\sigma} = \frac{82 - 70}{5} = \frac{12}{5} = 2.4.$$

- 2 Explain in words what a z-score of -1.5 means, in the context of a data value's position relative to its distribution.

A z-score of -1.5 means the data value lies 1.5 standard deviations below the mean of its distribution.

Comparing values across different distributions

- 3 On Test A (mean 75, standard deviation 6), a student scores 87. On Test B (mean 70, standard deviation 4), a different student scores 77. Compare the two students' relative performance using z-scores.

Test A: $z = \frac{87 - 75}{6} = \frac{12}{6} = 2$. Test B: $z = \frac{77 - 70}{4} = \frac{7}{4} = 1.75$. Since $2 > 1.75$, the student on Test A performed better relative to their own test's distribution, even though the raw score gap (12 vs. 7) might suggest otherwise.

Effect of adding a constant to every data value

- 4 A data set has mean 50 and standard deviation 8. If 10 is added to every value, find the new mean and new standard deviation.

Adding a constant shifts the mean by that constant but does not change the spread: new mean $= 50 + 10 = 60$; standard deviation stays 8.

- 5 Explain, without doing a full calculation, why adding a constant to every data value does not change the standard deviation.

Standard deviation measures how spread out the data is around the mean. Adding the same constant to every value shifts the entire data set (and its mean) by that constant, but every value's distance from the (also shifted) mean stays exactly the same — so the spread, and hence the standard deviation, is unaffected.

Effect of multiplying every data value by a constant

- 6 A data set has mean 20 and standard deviation 3. If every value is multiplied by 4, find the new mean and new standard deviation.

Multiplying by a constant scales both the mean and the standard deviation by that constant (standard deviation uses the absolute value of the constant): new mean = $20 \times 4 = 80$; new standard deviation = $3 \times 4 = 12$.

- 7 A distance measured in miles has mean 10 and standard deviation 2. Convert both to kilometers (multiply by 1.609), and find the new mean and standard deviation, to two decimal places.

New mean = $10 \times 1.609 = 16.09$ km. New standard deviation = $2 \times 1.609 = 3.22$ km.

Visualizing a standardized (z-score) distribution

- 8 The graph shows the standard normal distribution, which has mean 0 and standard deviation 1 — the distribution that results after standardizing any normal data set using z-scores.

a State the mean and standard deviation shown by this curve. Mean = 0 (the curve is centered at $x = 0$) and standard deviation = 1 (this is the standard normal curve).

b Explain why standardizing a data set using $z = \frac{x - \mu}{\sigma}$ always produces a distribution with mean 0 and standard deviation 1, regardless of the original data's mean and standard deviation. Subtracting μ shifts the mean to 0 (since subtracting a constant equal to the mean re-centers the data there), and dividing by σ rescales the standard deviation to $\frac{\sigma}{\sigma} = 1$ — combining a shift and a scaling that always produces mean 0, standard deviation 1.

Applying linear transformations in context

- 9 A factory's daily output (in units) has mean 500 and standard deviation 30. Management adds a fixed bonus batch of 50 units to every day's count, then reports output in dozens (divide by 12). Find the new mean and standard deviation, to two decimal places.

Adding 50: new mean = 550, standard deviation stays 30. Dividing by 12: new mean = $\frac{550}{12} \approx 45.83$ dozens, new standard deviation = $\frac{30}{12} = 2.5$ dozens.

Finding a raw value from a z-score

- 10 A distribution has mean $\mu = 64$ and standard deviation $\sigma = 5$. Find the data value corresponding to a z-score of $z = -1.8$.

$$x = \mu + z\sigma = 64 + (-1.8)(5) = 64 - 9 = 55.$$

- 11 A distribution has mean $\mu = 200$ and standard deviation $\sigma = 15$. If a data value has $z = 2.2$, find the data value.

$$x = \mu + z\sigma = 200 + 2.2(15) = 200 + 33 = 233.$$

Combined shift-and-scale transformations

- 12 A teacher curves test scores (mean 65, standard deviation 10) by first adding 5 points to every score, then multiplying the result by 1.1. Find the new mean and new standard deviation, to two decimal places.

After adding 5: mean = 70, standard deviation stays 10. After multiplying by 1.1: new mean = $70 \times 1.1 = 77$, new standard deviation = $10 \times 1.1 = 11$.

- 13 A weight distribution has mean $\mu = 68$ kg and standard deviation $\sigma = 9$ kg. Find the z-scores for weights of 77 kg and 50 kg.

$$z_{77} = \frac{77 - 68}{9} = \frac{9}{9} = 1. \quad z_{50} = \frac{50 - 68}{9} = \frac{-18}{9} = -2.$$

- 14 A national math exam has mean 500 and standard deviation 100, while a national verbal exam has mean 500 and standard deviation 110. A student scores 640 on the math exam and 625 on the verbal exam. Even though the verbal raw score is lower, use z-scores to determine which exam the student performed better on relative to other test-takers, and explain your reasoning.

Math z-score: $\frac{640 - 500}{100} = 1.4$. Verbal z-score: $\frac{625 - 500}{110} \approx 1.14$. Since $1.4 > 1.14$, the student performed better relative to other test-takers on the math exam, despite having a numerically higher raw score on the verbal exam — this shows why comparing raw scores directly across different distributions can be misleading, and z-scores put both exams on a common, comparable scale.

- 15 A shipping company records package weights with mean 18 kg and standard deviation 2.5 kg. The company then converts every measurement from kilograms to pounds by multiplying by 2.20462, and separately subtracts 1 kg from every package to account for packaging weight before converting. Describe, step by step, how each operation (the subtraction, then the multiplication) affects the mean and standard deviation.

Subtracting 1 kg from every package: new mean = $18 - 1 = 17$ kg; standard deviation is unchanged at 2.5 kg (subtracting a constant doesn't affect spread). Multiplying by 2.20462 to convert to pounds: new mean = $17 \times 2.20462 \approx 37.48$ lbs; new standard deviation = $2.5 \times 2.20462 \approx 5.51$ lbs (multiplying scales both the mean and the standard deviation by the same factor).

Multiple choice — easy 15

16 A distribution has mean $\mu = 51$ and standard deviation $\sigma = 25$. Find the z-score for a value of $x = 41$.

a. -0.40 ✓

b. 0.40

c. -0.90

d. 0.10

$$z = \frac{x - \mu}{\sigma} = \frac{41 - 51}{25} = -0.40.$$

17 A distribution has mean $\mu = 80$ and standard deviation $\sigma = 4$. Find the z-score for a value of $x = 126$.

a. -11.50

b. 11.00

c. 11.50 ✓

d. 12.00

$$z = \frac{x - \mu}{\sigma} = \frac{126 - 80}{4} = 11.50.$$

18 A distribution has mean $\mu = 30$ and standard deviation $\sigma = 2$. Find the z-score for a value of $x = 66$.

a. 18.50

b. -18.00

c. 18.00 ✓

d. 17.50

$$z = \frac{x - \mu}{\sigma} = \frac{66 - 30}{2} = 18.00.$$

19 A distribution has mean $\mu = 94$ and standard deviation $\sigma = 5$. Find the z-score for a value of $x = 71$.

a. 4.60

b. -5.10

c. -4.60 ✓

d. -4.10

$$z = \frac{x - \mu}{\sigma} = \frac{71 - 94}{5} = -4.60.$$

20 A distribution has mean $\mu = 50$ and standard deviation $\sigma = 2$. Find the z-score for a value of $x = -5$.

a. **-27.50 ✓**

b. -28.00

c. -27.00

d. 27.50

$$z = \frac{x - \mu}{\sigma} = \frac{-5 - 50}{2} = -27.50.$$

21 A distribution has mean $\mu = 85$ and standard deviation $\sigma = 2$. Find the z-score for a value of $x = 133$.

a. 24.50

b. **24.00 ✓**

c. -24.00

d. 23.50

$$z = \frac{x - \mu}{\sigma} = \frac{133 - 85}{2} = 24.00.$$

22 A distribution has mean $\mu = 77$ and standard deviation $\sigma = 25$. Find the z-score for a value of $x = 50$.

a. -0.58

b. 1.08

c. -1.58

d. **-1.08 ✓**

$$z = \frac{x - \mu}{\sigma} = \frac{50 - 77}{25} = -1.08.$$

23 A distribution has mean $\mu = 89$ and standard deviation $\sigma = 8$. Find the z-score for a value of $x = 40$.

a. **-6.12 ✓**

b. -6.62

c. 6.12

d. -5.62

$$z = \frac{x - \mu}{\sigma} = \frac{40 - 89}{8} = -6.12.$$

24 A distribution has mean $\mu = 98$ and standard deviation $\sigma = 5$. Find the z-score for a value of $x = 154$.

a. 10.70

b. 11.70

c. **11.20 ✓**

d. -11.20

$$z = \frac{x - \mu}{\sigma} = \frac{154 - 98}{5} = 11.20.$$

25 A distribution has mean $\mu = 78$ and standard deviation $\sigma = 8$. Find the z-score for a value of $x = 19$.

a. 7.38

b. **-7.38 ✓**

c. -7.88

d. -6.88

$$z = \frac{x - \mu}{\sigma} = \frac{19 - 78}{8} = -7.38.$$

26 A distribution has mean $\mu = 79$ and standard deviation $\sigma = 25$. Find the z-score for a value of $x = 30$.

a. -1.46

b. **-1.96 ✓**

c. -2.46

d. 1.96

$$z = \frac{x - \mu}{\sigma} = \frac{30 - 79}{25} = -1.96.$$

27 A distribution has mean $\mu = 80$ and standard deviation $\sigma = 20$. Find the z-score for a value of $x = 119$.

a. 1.95 ✓

b. 1.45

c. 2.45

d. -1.95

$$z = \frac{x - \mu}{\sigma} = \frac{119 - 80}{20} = 1.95.$$

28 A distribution has mean $\mu = 59$ and standard deviation $\sigma = 4$. Find the z-score for a value of $x = 63$.

a. 1.50

b. -1.00

c. 0.50

d. 1.00 ✓

$$z = \frac{x - \mu}{\sigma} = \frac{63 - 59}{4} = 1.00.$$

29 A distribution has mean $\mu = 69$ and standard deviation $\sigma = 2$. Find the z-score for a value of $x = 44$.

a. -12.00

b. 12.50

c. -12.50 ✓

d. -13.00

$$z = \frac{x - \mu}{\sigma} = \frac{44 - 69}{2} = -12.50.$$

30 A distribution has mean $\mu = 51$ and standard deviation $\sigma = 4$. Find the z-score for a value of $x = 50$.

- a. -0.75
- b. -0.55
- c. -0.25 ✓
- d. 0.25

$$z = \frac{x - \mu}{\sigma} = \frac{50 - 51}{4} = -0.25.$$

Multiple choice — medium 15

31 A distribution has mean $\mu = 85$ and standard deviation $\sigma = 25$. Find the data value corresponding to a z-score of $z = -1.4$.

- a. 49
- b. 51
- c. 120
- d. 50 ✓

$$x = \mu + z\sigma = 85 + (-1.4)(25) = 85 + (-35) = 50.$$

32 A distribution has mean $\mu = 168$ and standard deviation $\sigma = 10$. Find the data value corresponding to a z-score of $z = -2.8$.

- a. 139
- b. 196
- c. 141
- d. 140 ✓

$$x = \mu + z\sigma = 168 + (-2.8)(10) = 168 + (-28) = 140.$$

- 33 A distribution has mean $\mu = 49$ and standard deviation $\sigma = 5$. Find the data value corresponding to a z-score of $z = -1.2$.

- a. 42
- b. 55
- c. 43 ✓
- d. 44

$$x = \mu + z\sigma = 49 + (-1.2)(5) = 49 + (-6) = 43.$$

- 34 A distribution has mean $\mu = 76$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = 1.3$.

- a. 101
- b. 103
- c. 102 ✓
- d. 50

$$x = \mu + z\sigma = 76 + (1.3)(20) = 76 + (26) = 102.$$

- 35 A distribution has mean $\mu = 130$ and standard deviation $\sigma = 10$. Find the data value corresponding to a z-score of $z = -0.4$.

- a. 134
- b. 126 ✓
- c. 125
- d. 127

$$x = \mu + z\sigma = 130 + (-0.4)(10) = 130 + (-4) = 126.$$

- 36 A distribution has mean $\mu = 180$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = -2.2$.

- a. 224
- b. 137
- c. 135
- d. 136 ✓

$$x = \mu + z\sigma = 180 + (-2.2)(20) = 180 + (-44) = 136.$$

- 37 A distribution has mean $\mu = 101$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = -2.9$.

- a. 44
- b. 159
- c. 43 ✓
- d. 42

$$x = \mu + z\sigma = 101 + (-2.9)(20) = 101 + (-58) = 43.$$

- 38 A distribution has mean $\mu = 123$ and standard deviation $\sigma = 25$. Find the data value corresponding to a z-score of $z = 0.4$.

- a. 113
- b. 132
- c. 133 ✓
- d. 134

$$x = \mu + z\sigma = 123 + (0.4)(25) = 123 + (10) = 133.$$

- 39 A distribution has mean $\mu = 132$ and standard deviation $\sigma = 4$. Find the data value corresponding to a z-score of $z = -2.5$.

- a. 142
- b. 122 ✓
- c. 123
- d. 121

$$x = \mu + z\sigma = 132 + (-2.5)(4) = 132 + (-10) = 122.$$

- 40 A distribution has mean $\mu = 189$ and standard deviation $\sigma = 5$. Find the data value corresponding to a z-score of $z = 0.8$.

- a. 193 ✓
- b. 194
- c. 185
- d. 192

$$x = \mu + z\sigma = 189 + (0.8)(5) = 189 + (4) = 193.$$

- 41 A distribution has mean $\mu = 50$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = 2.4$.

- a. 98 ✓
- b. 97
- c. 99
- d. 2

$$x = \mu + z\sigma = 50 + (2.4)(20) = 50 + (48) = 98.$$

- 42 A distribution has mean $\mu = 144$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = 2.1$.

- a. 185
- b. 102
- c. 186 ✓
- d. 187

$$x = \mu + z\sigma = 144 + (2.1)(20) = 144 + (42) = 186.$$

- 43 A distribution has mean $\mu = 57$ and standard deviation $\sigma = 10$. Find the data value corresponding to a z-score of $z = -1.9$.

- a. 37
- b. 38 ✓
- c. 39
- d. 76

$$x = \mu + z\sigma = 57 + (-1.9)(10) = 57 + (-19) = 38.$$

- 44 A distribution has mean $\mu = 143$ and standard deviation $\sigma = 20$. Find the data value corresponding to a z-score of $z = 0.8$.

- a. 127
- b. 158
- c. 160
- d. 159 ✓

$$x = \mu + z\sigma = 143 + (0.8)(20) = 143 + (16) = 159.$$

- 45 A distribution has mean $\mu = 111$ and standard deviation $\sigma = 8$. Find the data value corresponding to a z-score of $z = 0.5$.

a. 115 ✓

b. 107

c. 114

d. 116

$$x = \mu + z\sigma = 111 + (0.5)(8) = 111 + (4) = 115.$$

Multiple choice — hard 15

- 46 On Test A (mean 79, standard deviation 4), a student scores 94. On Test B (mean 88, standard deviation 8), a different student scores 99. Using z-scores, which student performed better relative to their own test's distribution?

a. Test A ✓

b. Test B

c. Cannot be determined without more information

d. Both are exactly the same

Test A: $z = \frac{94 - 79}{4} \approx 3.75$. Test B: $z = \frac{99 - 88}{8} \approx 1.38$. Since $3.75 > 1.38$, Test A shows the stronger relative performance.

- 47 On Test A (mean 52, standard deviation 4), a student scores 53. On Test B (mean 54, standard deviation 5), a different student scores 61. Using z-scores, which student performed better relative to their own test's distribution?

a. Test A

b. Both are exactly the same

c. Cannot be determined without more information

d. Test B ✓

Test A: $z = \frac{53 - 52}{4} \approx 0.25$. Test B: $z = \frac{61 - 54}{5} \approx 1.40$. Since $1.40 > 0.25$, Test B shows the stronger relative performance.

- 48 On Test A (mean 67, standard deviation 6), a student scores 69. On Test B (mean 55, standard deviation 5), a different student scores 64. Using z-scores, which student performed better relative to their own test's distribution?

a. Test B ✓

b. Both are exactly the same

c. Cannot be determined without more information

d. Test A

Test A: $z = \frac{69 - 67}{6} \approx 0.33$. Test B: $z = \frac{64 - 55}{5} \approx 1.80$. Since $1.80 > 0.33$, Test B shows the stronger relative performance.

- 49 On Test A (mean 56, standard deviation 6), a student scores 76. On Test B (mean 67, standard deviation 4), a different student scores 72. Using z-scores, which student performed better relative to their own test's distribution?

a. Both are exactly the same

b. Cannot be determined without more information

c. Test B

d. Test A ✓

Test A: $z = \frac{76 - 56}{6} \approx 3.33$. Test B: $z = \frac{72 - 67}{4} \approx 1.25$. Since $3.33 > 1.25$, Test A shows the stronger relative performance.

- 50 On Test A (mean 90, standard deviation 10), a student scores 94. On Test B (mean 76, standard deviation 4), a different student scores 86. Using z-scores, which student performed better relative to their own test's distribution?

a. Both are exactly the same

b. Test A

c. Test B ✓

d. Cannot be determined without more information

Test A: $z = \frac{94 - 90}{10} \approx 0.40$. Test B: $z = \frac{86 - 76}{4} \approx 2.50$. Since $2.50 > 0.40$, Test B shows the stronger relative performance.

- 51 A data set has mean 75 and standard deviation 17. First, -4 is added to every value; then every value is multiplied by 4. Find the new mean and standard deviation as an ordered pair (mean, SD).

- a. (284, 17)
- b. (285, 68)
- c. (284, 68) ✓
- d. (300, 68)

After adding -4 : mean = 71, SD stays 17. After multiplying by 4: new mean = $4 \times 71 = 284$, new SD = $4 \times 17 = 68$. (284, 68).

- 52 A data set has mean 40 and standard deviation 18. First, 5 is added to every value; then every value is multiplied by 2. Find the new mean and standard deviation as an ordered pair (mean, SD).

- a. (90, 18)
- b. (80, 36)
- c. (91, 36)
- d. (90, 36) ✓

After adding 5: mean = 45, SD stays 18. After multiplying by 2: new mean = $2 \times 45 = 90$, new SD = $2 \times 18 = 36$. (90, 36).

- 53 A data set has mean 75 and standard deviation 18. First, 2 is added to every value; then every value is multiplied by 5. Find the new mean and standard deviation as an ordered pair (mean, SD).

- a. (375, 90)
- b. (386, 90)
- c. (385, 90) ✓
- d. (385, 18)

After adding 2: mean = 77, SD stays 18. After multiplying by 5: new mean = $5 \times 77 = 385$, new SD = $5 \times 18 = 90$. (385, 90).

- 54 A data set has mean 40 and standard deviation 3. First, -7 is added to every value; then every value is multiplied by 4. Find the new mean and standard deviation as an ordered pair (mean, SD).

a. (132, 12) ✓

b. (133, 12)

c. (160, 12)

d. (132, 3)

After adding -7 : mean = 33, SD stays 3. After multiplying by 4: new mean = $4 \times 33 = 132$, new SD = $4 \times 3 = 12$. (132, 12).

- 55 A data set has mean 61 and standard deviation 17. First, 3 is added to every value; then every value is multiplied by 5. Find the new mean and standard deviation as an ordered pair (mean, SD).

a. (305, 85)

b. (320, 85) ✓

c. (321, 85)

d. (320, 17)

After adding 3: mean = 64, SD stays 17. After multiplying by 5: new mean = $5 \times 64 = 320$, new SD = $5 \times 17 = 85$. (320, 85).

- 56 A distribution has mean $\mu = 97$ and standard deviation $\sigma = 4$. Find the z-scores for values of 113 and 84, as an ordered pair (z_{113}, z_{84}) , to two decimal places.

a. (4.30, -3.25)

b. (4.00, -3.25) ✓

c. (4.00, -3.55)

d. (-3.25 , 4.00)

$$z_{113} = \frac{113 - 97}{4} \approx 4.00. z_{84} = \frac{84 - 97}{4} \approx -3.25. (4.00, -3.25).$$

- 57 A distribution has mean $\mu = 41$ and standard deviation $\sigma = 8$. Find the z-scores for values of 52 and 31, as an ordered pair (z_{52}, z_{31}) , to two decimal places.

a. $(1.38, -1.25)$ ✓

b. $(1.68, -1.25)$

c. $(-1.25, 1.38)$

d. $(1.38, -1.55)$

$$z_{52} = \frac{52 - 41}{8} \approx 1.38. z_{31} = \frac{31 - 41}{8} \approx -1.25. (1.38, -1.25).$$

- 58 A distribution has mean $\mu = 82$ and standard deviation $\sigma = 8$. Find the z-scores for values of 104 and 70, as an ordered pair (z_{104}, z_{70}) , to two decimal places.

a. $(2.75, -1.50)$ ✓

b. $(-1.50, 2.75)$

c. $(2.75, -1.80)$

d. $(3.05, -1.50)$

$$z_{104} = \frac{104 - 82}{8} \approx 2.75. z_{70} = \frac{70 - 82}{8} \approx -1.50. (2.75, -1.50).$$

- 59 A distribution has mean $\mu = 53$ and standard deviation $\sigma = 5$. Find the z-scores for values of 74 and 47, as an ordered pair (z_{74}, z_{47}) , to two decimal places.

a. $(4.50, -1.20)$

b. $(4.20, -1.50)$

c. $(-1.20, 4.20)$

d. $(4.20, -1.20)$ ✓

$$z_{74} = \frac{74 - 53}{5} \approx 4.20. z_{47} = \frac{47 - 53}{5} \approx -1.20. (4.20, -1.20).$$

60 A distribution has mean $\mu = 50$ and standard deviation $\sigma = 3$. Find the z-scores for values of 57 and 43, as an ordered pair (z_{57}, z_{43}) , to two decimal places.

a. $(2.33, -2.63)$

b. $(2.33, -2.33)$ ✓

c. $(-2.33, 2.33)$

d. $(2.63, -2.33)$

$$z_{57} = \frac{57 - 50}{3} \approx 2.33. \quad z_{43} = \frac{43 - 50}{3} \approx -2.33. \quad (2.33, -2.33).$$