

Graphs of Inverse Trigonometric Functions & Their Derivatives (AA HL)

ANSWER KEY



Domain and range of arcsin

- 1 State the domain and range of $y = \arcsin x$.
Domain: $-1 \leq x \leq 1$. Range: $-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$.
-

Domain and range of arccos

- 2 State the domain and range of $y = \arccos x$.
Domain: $-1 \leq x \leq 1$. Range: $0 \leq y \leq \pi$.
-

Domain and range of arctan

- 3 State the domain and range of $y = \arctan x$.
Domain: all real numbers. Range: $-\frac{\pi}{2} < y < \frac{\pi}{2}$.
-

Evaluating inverse trigonometric functions exactly

- 4 Find the exact value of $\arcsin(\sqrt{3}/2)$, in radians.
 $\arcsin(\sqrt{3}/2) = \pi/3$, since $\sin(\pi/3) = \sqrt{3}/2$ and $\pi/3$ lies in the range $[-\pi/2, \pi/2]$.
- 5 Find the exact value of $\arccos(-1/2)$, in radians.
 $\arccos(-1/2) = 2\pi/3$, since $\cos(2\pi/3) = -1/2$ and $2\pi/3$ lies in the range $[0, \pi]$.
-

Differentiating arcsin using the chain rule

- 6 Differentiate $y = \arcsin(3x)$ with respect to x .
Using $\frac{d}{dx} \arcsin(u) = \frac{u'}{\sqrt{1-u^2}}$ with $u = 3x$, $u' = 3$: $\frac{dy}{dx} = \frac{3}{\sqrt{1-9x^2}}$.
-

Differentiating arctan using the chain rule

- 7 Differentiate $y = \arctan(x^2)$ with respect to x .
Using $\frac{d}{dx} \arctan(u) = \frac{u'}{1+u^2}$ with $u = x^2$, $u' = 2x$: $\frac{dy}{dx} = \frac{2x}{1+x^4}$.
-

Differentiating a product involving arccos

- 8 Differentiate $y = x \arccos(x)$ with respect to x .

By the product rule, $\frac{dy}{dx} = 1 \cdot \arccos(x) + x \cdot \frac{d}{dx} \arccos(x)$. Since $\frac{d}{dx} \arccos(x) = -1/\sqrt{1-x^2}$, the second term is $-x/\sqrt{1-x^2}$. So $\frac{dy}{dx} = \arccos(x) - \frac{x}{\sqrt{1-x^2}}$.

Finding a gradient at a point

- 9 Find the gradient of the curve $y = \arcsin(x/2)$ at $x = 1$.

Simplifying first: $\frac{dy}{dx} = \frac{1/2}{\sqrt{1-x^2/4}} = \frac{1}{\sqrt{4-x^2}}$ (multiplying top and bottom by 2). At $x = 1$, this is $\frac{1}{\sqrt{3}}$, i.e. $\sqrt{3}/3$.

Deriving the derivative of arctan from first principles

- 10 If $y = \arctan x$, then $x = \tan y$. Differentiate implicitly with respect to x to show that $\frac{dy}{dx} = \frac{1}{1+x^2}$.
Let $y' = \frac{dy}{dx}$. Differentiating $x = \tan y$ implicitly with respect to x gives $1 = \sec^2 y \cdot y'$. Since $\sec^2 y = 1 + \tan^2 y = 1 + x^2$ (using $\tan y = x$), this becomes $1 = (1+x^2)y'$, so $\frac{dy}{dx} = \frac{1}{1+x^2}$, as required.
-

Identifying the graph of arctan

- 11 Which of the following best describes the graph of $y = \arctan x$?

- a. An odd function with horizontal asymptotes $y = \pm \frac{\pi}{2}$ ✓
- b. An even function with a vertical asymptote at $x = 0$
- c. A periodic function with period 2π
- d. A function with domain restricted to $[-1, 1]$

$\arctan x$ is odd ($\arctan(-x) = -\arctan x$) and, as $x \rightarrow \pm \infty$, $\arctan x \rightarrow \pm \frac{\pi}{2}$, so its graph has horizontal (not vertical) asymptotes at $y = \pm \frac{\pi}{2}$ and its domain is all real numbers, not $[-1, 1]$.

Tangent line to an inverse trigonometric curve

- 12 Find the equation of the tangent line to $y = \arcsin x$ at $x = 0$.

At $x = 0$: $y = \arcsin(0) = 0$. The derivative is $\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$, which at $x = 0$ gives gradient 1. The tangent line is $y - 0 = 1(x - 0)$, i.e. $y = x$.