

Power Series and Taylor Series



Radius and interval of convergence

- 1 Find the radius of convergence of $\sum_{n=1}^{\infty} \frac{x^n}{n}$.
 - 2 Find the radius of convergence of $\sum_{n=0}^{\infty} \frac{x^n}{n!}$.
-

Taylor and Maclaurin series

- 3 Find the first three nonzero terms of the Maclaurin series for $f(x) = e^x$, using $f^{(n)}(0) = 1$ for all n .
 - 4 Find the third-degree Taylor polynomial for $f(x) = \cos x$ centered at $x = 0$.
 - 5 Use the Maclaurin series $\sin x = x - \frac{x^3}{6} + \frac{x^5}{120} - \dots$ to approximate $\sin(0.2)$ using the first two nonzero terms, to four decimal places.
-

Interval of convergence (including endpoints)

- 6 Find the interval of convergence of $\sum_{n=1}^{\infty} \frac{(x-2)^n}{n}$, checking both endpoints.
 - 7 Find the radius of convergence of $\sum_{n=1}^{\infty} n! x^n$.
-

Using known series

- 8 Using the known Maclaurin series $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$, write the Maclaurin series for e^{-x^2} (substitute $-x^2$ for x), giving the first three terms.
 - 9 Using $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$ for $|x| < 1$, write the power series for $\frac{1}{1+x^2}$ (substitute $-x^2$ for x).
-

Taylor series centered at a nonzero point

- 10 Find the first three terms of the Taylor series for $f(x) = \ln x$ centered at $x = 1$, using $f(1) = 0$, $f'(1) = 1$, $f''(1) = -1$.
- 11 Find the second-degree Taylor polynomial for $f(x) = \sqrt{x}$ centered at $x = 4$, given $f(4) = 2$, $f'(4) = \frac{1}{4}$, $f''(4) = -\frac{1}{32}$.

Error estimation with Taylor polynomials

- 12 Using the second-degree Maclaurin polynomial for $\cos x$, $T_2(x) = 1 - \frac{x^2}{2}$, estimate $\cos(0.3)$ to four decimal places, and compare to the true value $\cos(0.3) \approx 0.9553$.