

Volumes of Revolution



The disk and washer methods

- 1 Find the volume generated when the region under $y = \sqrt{x}$ on $[0, 4]$ is revolved about the x -axis.
 - 2 The region between $y = x^2$ and $y = 4$ is revolved about the x -axis. Using the washer method, set up and evaluate the volume integral.
-

The shell method

- 3 The region under $y = x^2$ on $[0, 2]$ is revolved about the y -axis. Use the shell method to find the volume.
 - 4 The region bounded by $y = x$, $y = 0$, and $x = 3$ is revolved about the y -axis. Find the volume using the shell method.
-

Comparing methods

- 5 The region under $y = x^2$ on $[0, 2]$ is revolved about the x -axis (not the y -axis). Find the volume using the disk method.
 - 6 The region bounded by $y = \sqrt{x}$ and $y = x$ (between their two intersection points) is revolved about the x -axis. Find the volume using the washer method.
-

Volumes about other axes

- 7 The region under $y = x^2$ on $[0, 2]$ is revolved about the line $y = -1$. Find the volume, using outer radius $x^2 + 1$.
 - 8 The region under $y = x^2$ on $[0, 2]$ is revolved about the line $x = 3$ (using the shell method with radius $3 - x$).
-

Volumes with trigonometric and exponential curves

- 9 The region under $y = \sin x$ on $[0, \pi]$ is revolved about the x -axis. Find the volume, using $\sin^2 x = \frac{1 - \cos 2x}{2}$.
- 10 The region under $y = e^{-x}$ on $[0, 1]$ is revolved about the x -axis. Find the volume, to three decimal places.

Setting up without evaluating

- 11 Set up (do not evaluate) the volume integral for the region under $y = \ln x$ on $[1, e]$, revolved about the x -axis.
- 12 Set up (do not evaluate) the volume integral for the region between $y = x^2$ and $y = \sqrt{x}$ (on $[0, 1]$) revolved about the y -axis, using the shell method.