

## Higher derivatives



- 1
- a  $\frac{dy}{dx} = -10x^{-6}$       b  $\frac{d^2y}{dx^2} = 60x^{-7}$       c  $\frac{d^3y}{dx^3} = -420x^{-8}$
- 2
- a  $\frac{dy}{dx} = 40x^7$       b  $\frac{d^2y}{dx^2} = 280x^6$       c  $\frac{d^3y}{dx^3} = 1680x^5$       d  $\frac{d^4y}{dx^4} = 8400x^4$
- e  $\frac{d^5y}{dx^5} = 33\,600x^3$
- 3
- a  $\frac{dy}{dx} = -8x^3 + 12x^2 - 14x + 8$       b  $\frac{dy}{dx} = -142$
- c  $\frac{d^2y}{dx^2} = -24x^2 + 24x - 14$       d  $\frac{d^2y}{dx^2} = -302$
- e  $\frac{d^3y}{dx^3} = -48x + 24$       f  $\frac{d^3y}{dx^3} = -264$
- 4
- a  $y'' = 12x^2 - 36x + 16$       b  $f''(x) = -\frac{6}{49}x^{-\frac{13}{7}}$
- c  $f''(x) = 108(3x + 2)^2$       d  $f''(x) = 24(x + 8)^{-4}$
- e  $y'' = -4(7 - 6x^2)^{-\frac{5}{3}}(2x^2 + 7)$       f  $y'' = -(6 - x)^{-\frac{3}{2}}$
- g  $f''(x) = 20x^3 + 6$       h  $f''(x) = -84x^5 + 36x^2 - 24x$

---

## Concavity

- 5
- a  $B, C, D$       b  $A$
- 6
- a  $f'(a) < 0, f''(a) > 0$       b  $f'(a) > 0, f''(a) < 0$
- c  $f'(a) > 0, f''(a) > 0$       d  $f'(a) < 0, f''(a) < 0$

7



a

i Positive

ii Minimum

iii  $f'(x) = 2x - 4$

iv  $f''(x) = 2$

v Concave up

b

i Negative

ii Maximum

iii  $f'(x) = -2x + 4$

iv  $f''(x) = -2$

v Concave down

8

a  $f'(x) = 8x + 3$

b  $f''(x) = 8$

c

|         |   |    |    |    |    |
|---------|---|----|----|----|----|
| $x$     | 0 | 1  | 2  | 3  | 4  |
| $f'(x)$ | 3 | 11 | 19 | 27 | 35 |

d Gradient is increasing at a constant rate.

e No, because  $f(x)$  is a quadratic function, and it is always concave up.

9

a  $f'(x) = -6x + 12$

b  $f''(x) = -6$

c Gradient is decreasing at a constant rate.

d No, because  $f(x)$  is a quadratic function, and it is always concave down.

10



a

- i No such  $x$ -values
- ii  $x > -1$
- iii The graph is always increasing for  $x \geq -1$ . There is no value of  $x$  in the function such that  $f''(x) = 0$  and concavity changes.

b

- i  $x > 2$
- ii No such  $x$ -values
- iii The graph is always decreasing for  $x \geq 2$ . There is no value of  $x$  in the function such that  $f''(x) = 0$  and concavity changes.

c

- i  $x > 0$
- ii  $x < 0$
- iii The graph is always decreasing for  $x < 0$  and for  $x > 0$ . There is no value of  $x$  in the function such that  $f''(x) = 0$  and concavity changes.

d

- i  $x < 0$  or  $x > 0$
- ii No such  $x$ -values
- iii The graph is always increasing for  $x < 0$  and always decreasing for  $x > 0$ . There is no value of  $x$  in the function such that  $f''(x) = 0$  and concavity changes.

11

a Horizontal translation of 6 units left

c  $(-6, 0)$ b  $(0, 0)$ 

d

|       |    |    |    |
|-------|----|----|----|
| $x$   | -7 | -6 | -5 |
| $y'$  | 3  | 0  | 3  |
| $y''$ | -6 | 0  | 6  |

e Yes, because  $f'(-6) = 0$ .f  $x > -6$ 

12

a Horizontal translation of 5 units left

c  $(-5, 0)$ b  $(0, 0)$ 

d

|       |    |    |    |
|-------|----|----|----|
| $x$   | -6 | -5 | -4 |
| $y'$  | 3  | 0  | 3  |
| $y''$ | -6 | 0  | 6  |

e Yes, because  $f'(-5) = 0$ .f  $x > -5$

13



a  $y'' = 12x^2 - 48x$

b  $(0, -9), (4, -265)$

c

|       |      |   |     |      |     |
|-------|------|---|-----|------|-----|
| $x$   | -2   | 0 | 2   | 4    | 6   |
| $y'$  | -128 | 0 | -96 | -128 | 0   |
| $y''$ | 144  | 0 | -48 | 0    | 144 |

d  $(0, -9)$  is a horizontal point of inflection and  $(4, -265)$  is an ordinary point of inflection.

e  $x < 0$  or  $x > 4$

14

a  $y'' = 24x - 32$

b  $\left(\frac{4}{3}, -\frac{206}{27}\right)$

c An ordinary point of inflection

d  $x < \frac{4}{3}$

15

a  $y'' = 18x + 12$

b  $\left(-\frac{2}{3}, -\frac{5}{9}\right)$

c Ordinary

d  $x < -\frac{2}{3}$

16

a  $y'' = 20x^3 - 6$

b  $\left(\sqrt[3]{\frac{3}{10}}, \left(\sqrt[3]{\frac{3}{10}}\right)^5 - 3\left(\sqrt[3]{\frac{3}{10}}\right)^2\right)$

c Ordinary

d  $x > \sqrt[3]{\frac{3}{10}}$

e  $x < \sqrt[3]{\frac{3}{10}}$

17

a  $x < -3$  or  $x > 0$

b  $-3 < x < 0$

c  $f''(x) = 6x + 9$

d  $x > -\frac{3}{2}$

e  $x < -\frac{3}{2}$

f  $x = -3$

g  $x = -\frac{3}{2}$

## Stationary points

18

a  $x = 5$

b  $f''(5) = 2$



19

- a  $(-4, 0), (2, -108)$
- b  $f''(x) = 6x + 6$
- c  $(-4, 0)$  is a maximum turning point and  $(2, -108)$  is a minimum turning point.
- d  $x < -1$

20

- a  $y' = 4x^3 + 12x^2$
- b  $y'' = 12x^2 + 24x$
- c  $(0, 2), (-3, -25)$
- d  $(0, 2)$  is a point of inflection, and  $(-3, -25)$  is a minimum point.
- e  $(0, 2), (-2, -14)$
- f  $(0, 2)$  is a horizontal point of inflection, while  $(-2, -14)$  is an ordinary point of inflection.
- g  $x < -2$  or  $x > 0$

21

- a  $f'(x) = (x - 5)(3x - 21)$
- b  $(5, 0), (7, -4)$
- c  $f''(x) = 6x - 36$
- d Local minimum at  $(7, -4)$ , local maximum at  $(5, 0)$
- e  $x < 6$

22

- a
  - i  $(3, -61), (-3, 47)$
  - ii  $(-3, 47)$  is a maximum turning point and  $(3, -61)$  is a minimum turning point.
- b
  - i  $(2, -18), (-2, 14)$
  - ii  $(-2, 14)$  is a maximum turning point and  $(2, -18)$  is a minimum turning point.
- c
  - i  $(-4, 112), (1, -13)$
  - ii  $(-4, 112)$  is a maximum turning point and  $(1, -13)$  is a minimum turning point
- d
  - i  $(-1, 2), (0, 3), (1, 2)$
  - ii  $(0, 3)$  is a maximum turning point,  $(-1, 2)$  and  $(1, 2)$  are minimum turning points

23



- a
- i  $x = -\frac{3}{7}, -\frac{1}{3}$
  - ii Maximum turning point at  $x = -\frac{3}{7}$  and minimum turning point at  $x = -\frac{1}{3}$
- b
- i  $x = 1, 7$
  - ii Maximum turning point at  $x = 1$  and minimum turning point at  $x = 7$
- c
- i  $x = -2$
  - ii Maximum turning point.
- d
- i  $x = 5$
  - ii Horizontal point of inflection
- e
- i  $x = 5$
  - ii Horizontal point of inflection.

24

- a  $x = -\frac{3}{5}, -\frac{1}{3}$
- b  $x = -\frac{7}{15}$
- c 2

25

- a  $y'' = 30(x^2 - 5)(x^2 - 1)$
- b  $(-\sqrt{5}, 0), (-1, -64), (1, -64), (\sqrt{5}, 0)$
- c  $(-\sqrt{5}, 0), (\sqrt{5}, 0)$

26

- a
- i (5, 4)
  - ii (5, 4)
  - iii Turning point. The graph changes from decreasing to increasing on either side of  $x = 5$ , but the concavity does not change at  $x = 5$ .
- b
- i (4, 5)
  - ii (4, 5)
  - iii The graph changes from decreasing to increasing at  $x = 4$ , but the concavity does not change at  $x = 4$ .
- c
- i (8, 6)
  - ii (8, 6)
  - iii The graph changes from decreasing to increasing around  $x = 8$ , but the concavity does not change around  $x = 8$ .

27

a  $a = b$

c They can be any real numbers.



b  $a = b$

28

a  $3a + 2b + 12 = 0$

b  $b = -3a$

c  $a = 4$

d  $b = -12$

29

a  $a = -2$

b  $c = -48$