

Points of intersection



1

a 2

b 1

c 0

d 1

e 0

f 2

g 1

h 1

i 0

j 0

2 $(3, 4), (-2, 9)$

3 Minimum value of $k = 1$

4

a $4x^2 - 5x + 14 = 0$

b -199

c No, because the discriminant is negative which means the quadratic has no solution.


5 5

6 $(-5, -67)$

7 $k = 7$

8

a
i




ii $(1, 6), (-3, -2)$

b
i



ii $(-1, 7), (4, -8)$




9

a $x = 0, \frac{5}{2}$

c $\left(\frac{5}{4}, \frac{25}{4}\right)$

b $x = \frac{5}{4}$

d



e $c = 6.25$

f $c < \frac{25}{4}$


10

a

x	-4	-3	-2
$y = x^2 + 6x$	-8	-9	-8
$y = 2x$	-8	-6	-4



b


c $(0, 0), (-4, -8)$

11

a

x	1	2	3
$y = 4x - x^2$	3	4	3
$y = -2x + 5$	3	1	-1

b



c $(1, 3), (5, -5)$

12

a No

b $(-2, 7)$


13 Example answers:



14

a


i



ii $(-1, 1), (1, 1)$

b

i



ii $(-1, 2), (1, 2)$



Quadratic systems of equations

15

a

i $x = \pm 2$

ii $(-2, 0), (2, 0)$

b

i $x = \pm 2$

ii $(-2, 4), (2, 4)$

c

i $x = 0$

ii $(0, 0)$

d

i $x = 3, -5$

ii $(3, 25), (-5, 9)$

e

i $x = 7$

ii $(7, 8)$

f

i $x = 5, 4$

ii $(5, 54), (4, 38)$

g

i $x = \frac{1}{4}$

ii $\left(\frac{1}{4}, -\frac{109}{16}\right)$

h

i $x = \frac{21}{2}$

ii $\left(\frac{21}{2}, \frac{5}{4}\right)$

i

i $x = 2$

ii $(2, 6)$

j

i $x = \pm\sqrt{10}$

ii $(-\sqrt{10}, -8), (\sqrt{10}, -8)$

16

a

$(0, 0), (-7, 49)$

b

$(3, 18), (-7, 28)$

c

$(3, -7)$

d

$(0, 0), (1, 16)$

e

$(7, -7), (1, -7)$

f

$(-4, 42)$

g

$(2, 2), (-3, -3)$

h

$(3, 0), \left(\frac{7}{2}, \frac{1}{4}\right)$

i

$(2, 1), (-2, 1)$

17

a

$x = \frac{1 + \sqrt{17}}{2}, \frac{1 - \sqrt{17}}{2}$

b

$\left(\frac{1 + \sqrt{17}}{2}, -4\right), \left(\frac{1 - \sqrt{17}}{2}, -4\right)$

18 $(0, 16), (-1, 15), (-2, 12), (-3, 9), (-4, 4)$



19

a $(-2, 3), (2, -9)$

c $(2, -4), (0, 0)$

b $(-4, -12), (-3, -12)$

d i $x = 4, -3$ ii $(4, 0), (-3, 21)$

e i $x = -4, -5$ ii $(-4, 2), (-5, 5)$


20 $y = -x - \frac{21}{4}$

21 a $x = -4 \pm 2\sqrt{3}$

b i Yes ii Yes iii Yes

22 a $c = -8$

b $a = -2$






Quadratic equations from systems

23 $3x^2 - 2x + 9 = 0$

24

a



b 2


c $(0, 4), (-1, 5)$

25 $y = 2x + 6$

26

a $-3x + 1$

b




c $x = -1, 2$

27



28

a



b $(-3, 9), (3, 9)$

c $x = \pm 3$

Applications

29 $y = 1, \frac{8}{3}$

30

a $x = 0, x = 6$

b $y = 144$

c Example answer:

The equilibrium quantity occurs when 6 thousand units are being produced, which has a corresponding price of 144 dollars per unit.

