# Trigonometry

ClipSAT Printable Practice Suite (clipsat.org) • Topic 6.01

**ANSWER KEY — COMPLETE WORKED SOLUTIONS**

## Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

**1** **[Foundational]** • **Target: Right triangle trigonometric ratios**

In a right triangle, angle $\theta$ has an opposite side of 8 and hypotenuse 17. Find $\sin\theta$, $\cos\theta$, and $\tan\theta$.

The adjacent side: $\sqrt{17^2 - 8^2} = \sqrt{289 - 64} = \sqrt{225} = 15$. $\sin\theta = \frac{8}{17}$, $\cos\theta = \frac{15}{17}$, $\tan\theta = \frac{8}{15}$.

**2** **[Foundational]** • **Target: Right triangle trigonometric ratios**

A ladder 15 feet long leans against a wall, making a $70^\circ$ angle with the ground. Find the height reached on the wall, to the nearest tenth.

Height $= 15\sin(70^\circ) \approx 15(0.9397) \approx 14.1$ feet.

**3** **[Foundational]** • **Target: The unit circle and radian measure**

Convert $150^\circ$ to radians, in terms of $\pi$.

$150^\circ \times \frac{\pi}{180^\circ} = \frac{5\pi}{6}$.

**4** **[Foundational]** • **Target: The unit circle and radian measure**

Convert $\frac{3\pi}{4}$ radians to degrees.

$\frac{3\pi}{4} \times \frac{180^\circ}{\pi} = 135^\circ$.

**5** **[Foundational]** • **Target: The unit circle and radian measure**

Find the exact value of $\sin\left(\frac{\pi}{6}\right)$ and $\cos\left(\frac{\pi}{4}\right)$, using the unit circle.

$\sin\left(\frac{\pi}{6}\right) = \frac{1}{2}$. $\cos\left(\frac{\pi}{4}\right) = \frac{\sqrt{2}}{2}$.

**6** **[Foundational]** • **Target: Basic trigonometric identities**

Use the Pythagorean identity to find $\cos\theta$, given that $\sin\theta = \frac{3}{5}$ and $\theta$ is in the first quadrant.

$\sin^2\theta + \cos^2\theta = 1$, so $\cos^2\theta = 1 - \frac{9}{25} = \frac{16}{25}$. Since $\theta$ is in the first quadrant, $\cos\theta$ is positive: $\cos\theta = \frac{4}{5}$.

**7** **[Foundational]** • **Target: Graphs of sine and cosine functions**

State the amplitude and period of $y = 3\sin(2x)$.

Amplitude: $|3| = 3$. Period: $\frac{2\pi}{2} = \pi$.

## Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

© ClipSAT — clipsat.org | All Rights Reserved

**8** [Intermediate] • Target: Graphs of sine and cosine functions

State the amplitude, period, and vertical shift of $y = 4\cos(x) + 2$.

Amplitude: $4$. Period: $2\pi$ (unchanged, since there is no coefficient on $x$ inside the cosine). Vertical shift: up 2 units.

**9** [Intermediate] • Target: Law of Sines

In triangle $ABC$, $A = 40^\circ$, $B = 65^\circ$, and $a = 10$. Find side $b$, to the nearest tenth, using the Law of Sines.

Law of Sines: $\frac{a}{\sin A} = \frac{b}{\sin B}$. $\frac{10}{\sin 40^\circ} = \frac{b}{\sin 65^\circ}$. $b = \frac{10\sin 65^\circ}{\sin 40^\circ} \approx \frac{10(0.9063)}{0.6428} \approx 14.1$.

**10** [Intermediate] • Target: Law of Cosines

In triangle $ABC$, $a = 7$, $b = 9$, and $C = 60^\circ$. Find side $c$, using the Law of Cosines.

$c^2 = a^2 + b^2 - 2ab\cos C = 49 + 81 - 2(7)(9)\cos 60^\circ = 130 - 126(0.5) = 130 - 63 = 67$. $c = \sqrt{67} \approx 8.2$.

**11** [Intermediate] • Target: Applications of trigonometry to real-world problems

This question has two parts. A surveyor stands 200 feet from the base of a tower and measures the angle of elevation to the top as $38^\circ$. (a) Find the height of the tower, to the nearest foot. (b) If the surveyor moves 50 feet closer to the tower, find the new angle of elevation, to the nearest degree, using the height found in part (a).

(a) Height $= 200\tan(38^\circ) \approx 200(0.7813) \approx 156$ feet. (b) New distance: $200 - 50 = 150$ feet. New angle: $\theta = \arctan\left(\frac{156}{150}\right) = \arctan(1.04) \approx 46^\circ$.

**12** [Intermediate] • Target: Reciprocal trigonometric functions

If $\cos\theta = \frac{3}{5}$, find $\sec\theta$ and $\tan\theta$ (assuming $\theta$ is in the first quadrant).

$\sec\theta = \frac{1}{\cos\theta} = \frac{5}{3}$. Since $\sin^2\theta = 1 - \frac{9}{25} = \frac{16}{25}$, $\sin\theta = \frac{4}{5}$ (positive in the first quadrant).
$\tan\theta = \frac{\sin\theta}{\cos\theta} = \frac{4/5}{3/5} = \frac{4}{3}$.

**13** [Intermediate] • Target: Angle of elevation and depression

From the top of a 100-foot cliff, the angle of depression to a boat is $25^\circ$. Find the horizontal distance from the base of the cliff to the boat, to the nearest foot.

The angle of depression from the top equals the angle of elevation from the boat. Using the tangent ratio: $\tan(25^\circ) = \frac{100}{d}$, so $d = \frac{100}{\tan(25^\circ)} \approx \frac{100}{0.4663} \approx 214$ feet.

© ClipSAT — clipsat.org | All Rights Reserved

**14** **[Intermediate]** • **Target: A multi-part triangle problem combining Law of Sines and Law of Cosines**

This question has two parts. Triangle $ABC$ has $a = 8$, $b = 11$, and $C = 50^\circ$. (a) Use the Law of Cosines to find side $c$, to the nearest tenth. (b) Use the Law of Sines to find angle $A$, to the nearest degree.

(a) $c^2 = 8^2 + 11^2 - 2(8)(11)\cos 50^\circ = 64 + 121 - 176(0.6428) \approx 185 - 113.1 \approx 71.9$. $c \approx \sqrt{71.9} \approx 8.5$. (b) $\frac{a}{\sin A} = \frac{c}{\sin C}$: $\frac{8}{\sin A} = \frac{8.5}{\sin 50^\circ}$. $\sin A = \frac{8\sin 50^\circ}{8.5} \approx \frac{8(0.766)}{8.5} \approx 0.721$, so $A \approx 46^\circ$.

---

## Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

**15** **[Challenging]** • **Target: Cofunction and negative angle identities** • **Cross-Lesson: 7.01 Statistics & Probability**

Use the cofunction identity to find $\cos(30^\circ)$ given that $\sin(60^\circ) = \frac{\sqrt{3}}{2}$.

The cofunction identity states $\cos\theta = \sin(90^\circ - \theta)$, so $\cos(30^\circ) = \sin(90^\circ - 30^\circ) = \sin(60^\circ) = \frac{\sqrt{3}}{2}$.

**16** **[Challenging]** • **Target: Cofunction and negative angle identities** • **Cross-Lesson: 8.01 Matrices, Vectors & Complex Numbers**

Use the fact that sine is an odd function to find $\sin(-30^\circ)$, given $\sin(30^\circ) = \frac{1}{2}$.

Since sine is odd, $\sin(-\theta) = -\sin(\theta)$: $\sin(-30^\circ) = -\sin(30^\circ) = -\frac{1}{2}$.

**17** **[Challenging]** • **Target: Graphs of sine and cosine functions** • **Cross-Lesson: 1.01 Numbers, Percents & Ratios**

In right triangle $ABC$ with $C = 90^\circ$, $a = 5$, and $c = 13$. What is the value of $\sin A + \cos A$?

By the Pythagorean theorem, the adjacent leg is $b = \sqrt{13^2 - 5^2} = \sqrt{169 - 25} = 12$. Then $\sin A = \frac{a}{c} = \frac{5}{13}$ and $\cos A = \frac{b}{c} = \frac{12}{13}$. Adding them together: $\sin A + \cos A = \frac{5+12}{13} = \mathbf{\frac{17}{13}}$.

**18** **[Challenging]** • **Target: Law of Sines** • **Cross-Lesson: 2.01 Elementary & Intermediate Algebra**

Two sides of a non-degenerate triangle measure 7 inches and 11 inches. What is the range of possible perimeters for this triangle?

Let $x$ be the third side. By the Triangle Inequality Theorem, $11 - 7 < x < 11 + 7 \Rightarrow 4 < x < 18$. The perimeter is $P = 7 + 11 + x = 18 + x$. Substituting the strict bounds for $x$:
$18 + 4 < P < 18 + 18 \Rightarrow$ **$22 < P < 36$** inches

**19** **[Challenging]** • **Target: Law of Cosines** • **Cross-Lesson: 3.01 Functions & Graphs**

What is the period and amplitude of the trigonometric function $f(x) = -6\cos(4x - \pi) + 3$?

For any function in standard form $f(x) = A\cos(Bx - C) + D$: Amplitude: $|A| = |-6| =$ **$6$**.Period: $T = \frac{2\pi}{B} = \frac{2\pi}{4} =$ **$\frac{\pi}{2}$**.Midline: $y = D = 3$.

© ClipSAT — clipsat.org | All Rights Reserved

**20** [Challenging] • Target: Trigonometric equations • Cross-Lesson: 4.01 Coordinate Geometry

A circle has a radius of 12 cm. What is the area of a sector formed by a central angle of $75^\circ$?

The formula for the area of a circle sector is: $A = \frac{\theta}{360^\circ}\pi r^2 = \frac{75}{360}\pi(12^2) = \frac{5}{24} \cdot 144\pi = 5 \cdot 6\pi = \mathbf{30\pi}$ cm$^2$

## Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

**21** [Foundational] • Target: Right triangle trigonometric ratios

In a right triangle, the angle $\theta$ has an opposite side of length 20, adjacent side 21, and hypotenuse 29. Find $\tan\theta$.

**a.** $\frac{29}{21}$

**b.** $\frac{20}{21}$ ✓

**c.** $\frac{29}{20}$

**d.** $\frac{21}{20}$

$\tan\theta = opposite/adjacent = 20/21 = \frac{20}{21}$.

**22** [Foundational] • Target: The unit circle and radian measure

In triangle $ABC$, $a = 14$, $b = 7$, and $C = 105^\circ$. Find side $c$, to the nearest tenth, using the Law of Cosines.

**a.** 18.7

**b.** 17.2 ✓

**c.** 20.6

**d.** 15.7

$c^2 = a^2 + b^2 - 2ab\cos C = 196 + 49 - 2(14)(7)\cos 105^\circ = 245 - 196(-0.2588) \approx 295.7$. $c \approx \sqrt{295.7} \approx 17.2$.

**23** [Foundational] • Target: Basic trigonometric identities

In $\Delta ABC$, side $AB$ has length 6, side $AC$ has length 10, and the measure of $\angle BAC$ is $120^\circ$. The angle bisector of $\angle BAC$ intersects side $BC$ at point $D$. What is the length of segment $AD$?

**a.** $\frac{15}{4}$ ✓

**b.** $\frac{15\sqrt{3}}{4}$

**c.** $\frac{15}{2}$

**d.** $\frac{30}{7}$

We can find the length of the angle bisector $AD$ by decomposing the area of $\Delta ABC$ into the sum of the areas of $\Delta ABD$ and $\Delta ACD$. Since $AD$ bisects $\angle BAC = 120^\circ$, we have: $\angle BAD = \angle CAD = 60^\circ$ Using the sine formula for the area of a triangle, Area $= \frac{1}{2}ab\sin\theta$:

Area$(\Delta ABC) = \frac{1}{2} \cdot AB \cdot AC \cdot \sin(120^\circ) = \frac{1}{2} \cdot 6 \cdot 10 \cdot \frac{\sqrt{3}}{2} = 15\sqrt{3}$ The sum of the areas of the tw.

© ClipSAT — clipsat.org | All Rights Reserved

**24** [Foundational] • Target: Graphs of sine and cosine functions

In $\Delta ABC$, the length of side $AC$ is 5, the length of side $BC$ is 8, and the measure of $\angle C$ is $60^\circ$. What is the value of $\cos A$?

**a.** $\frac{1}{7}$ ✓

**b.** $\frac{2}{7}$

**c.** $\frac{5}{8}$

**d.** $\frac{11}{14}$

To find $\cos A$, we first determine the length of the third side, $c = AB$, using the Law of Cosines with the known angle $C = 60^\circ$: $c^2 = a^2 + b^2 - 2ab\cos C$ Substitute $a = BC = 8$, $b = AC = 5$, and $\cos(60^\circ) = \frac{1}{2}$: $c^2 = 8^2 + 5^2 - 2(8)(5)\left(\frac{1}{2}\right) = 64 + 25 - 40 = 49$ Taking the positive square root gives $c = AB = 7$. Next, apply the Law of Cosines rearranged for $\cos A$: $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$ Substitute $a = 8$, $b = 5$, and $c = 7$.

**25** [Foundational] • Target: Law of Sines

In a right triangle, the side opposite $\theta$ is 8, the side adjacent is 6, and the hypotenuse is 10. Find $\tan\theta$.

**a.** 0

**b.** $\frac{3}{4}$

**c.** $\frac{4}{3}$ ✓

**d.** $\frac{2}{25}$

$\tan\theta = \frac{\text{opp}}{\text{adj}} = \frac{8}{6} = \frac{4}{3}$.

**26** [Foundational] • Target: Law of Cosines

In a triangle, sides $a = 15$ and $b = 14$ meet at an included angle of $123^\circ$. Using the Law of Cosines, which expression gives side $c$?

**a.** $\sqrt{15^2 + 14^2 + 2(15)(14)\cos 123^\circ}$

**b.** $\sqrt{15^2 - 14^2 - 2(15)(14)\cos 123^\circ}$

**c.** $\sqrt{15^2 + 14^2 - 2(15)(14)\cos 123^\circ}$ ✓

**d.** $15^2 + 14^2 - 2(15)(14)\cos 123^\circ$

Law of Cosines: $c^2 = a^2 + b^2 - 2ab\cos C \Rightarrow c = \sqrt{15^2 + 14^2 - 2(15)(14)\cos 123^\circ}$.

© ClipSAT — clipsat.org | All Rights Reserved

**27** [Foundational] • Target: Trigonometric equations

In a right triangle, the side opposite angle $\theta$ is 18, the adjacent side is 24, and the hypotenuse is 30. What is the sine of $\theta$?

**a.** $\frac{3}{4}$

**b.** $\frac{5}{3}$

**c.** $\frac{4}{5}$

**d.** $\frac{3}{5}$ ✓

$\sin(\theta) = \frac{\text{opposite}}{\text{hypotenuse}} = \frac{18}{30} = \frac{3}{5}$

**28** [Foundational] • Target: Applications of trigonometry to real-world problems

In the right-angled triangle ABC shown below, $\angle ABC = 90^\circ$, $AB = 3$, and $BC = 4$. What is the length of the hypotenuse AC?

**a.** **5** ✓

**b.** 7

**c.** 12

**d.** 25

This question involves a direct application of the Pythagorean theorem in a right-angled triangle. The Pythagorean theorem states that in a right-angled triangle, the square of the hypotenuse (the side opposite the right angle) is equal to the sum of the squares of the other two sides. In triangle ABC, with $\angle ABC = 90^\circ$, AB and BC are the two shorter sides (legs), and AC is the hypotenuse. According to the Pythagorean theorem: $AC^2 = AB^2 + BC^2$ Given $AB = 3$ and $BC = 4$: $AC^2 = 3^2 + 4^2$.

## Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

**29** [Intermediate] • Target: Reciprocal trigonometric functions

In the right-angled triangle shown, what is the length of side $AC$ if $AB = 10$ and $\angle B = 30^\circ$?

**a.** **5** ✓

**b.** 10

**c.** $5\sqrt{3}$

**d.** 20

In the right-angled triangle $ABC$, we are given $AB$ as the hypotenuse and $\angle B = 30^\circ$. We need to find the length of the side opposite to $\angle B$, which is $AC$. Using the sine function, we have: $\sin(\angle B) = \frac{\text{Opposite}}{\text{Hypotenuse}} = \frac{AC}{AB}$ Substituting the given values: $\sin(30^\circ) = \frac{AC}{10}$ We know that $\sin(30^\circ) = \frac{1}{2}$. So, $\frac{1}{2} = \frac{AC}{10}$. Multiplying both sides by 10 to solve for $AC$: $AC = 10 \times \frac{1}{2}$.

© ClipSAT — clipsat.org | All Rights Reserved

**30** [Intermediate] • Target: Angle of elevation and depression

A circle with radius 8 has a sector with area $16\pi$. What is the measure of the central angle of this sector in radians?

**a.** $\frac{\pi}{4}$

**b.** $\frac{\pi}{2}$ ✓

**c.** $\frac{2\pi}{3}$

**d.** $\pi$

Area $= \frac{1}{2}r^2\theta \Rightarrow 16\pi = \frac{1}{2}(64)\theta = 32\theta \Rightarrow \theta = \frac{16\pi}{32} = \frac{\pi}{2}$.

**31** [Intermediate] • Target: Solving trigonometric equations over a restricted domain

What is the reference angle of $140^\circ$?

**a.** $64^\circ$

**b.** **$40^\circ$** ✓

**c.** $66^\circ$

**d.** $31^\circ$

In Quadrant II, reference angle $= 180^\circ - 140^\circ = 40^\circ$.

**32** [Intermediate] • Target: A multi-part triangle problem combining Law of Sines and Law of Cosines

In a right triangle, the hypotenuse is 10 and one angle is $30^\circ$. What is the length of the side opposite the $30^\circ$ angle?

**a.** 6

**b.** **5** ✓

**c.** 10

**d.** 4

$\sin(30^\circ) = \frac{1}{2} = \frac{opp}{10} \Rightarrow opp = 5$

**33** [Intermediate] • Target: Cofunction and negative angle identities

A right triangle has legs 5 and 12 with hypotenuse 13. Find $\tan\theta$, where $\theta$ is the angle opposite the side of length 5.

**a.** $\frac{12}{13}$

**b.** $\frac{13}{5}$

**c.** $\frac{12}{5}$

**d.** $\frac{5}{12}$ ✓

$\tan\theta = \text{opposite/adjacent} = \frac{5}{12}$.

© ClipSAT — clipsat.org | All Rights Reserved

**34** [Intermediate] • Target: Right triangle trigonometric ratios

If $\theta$ is an angle in Quadrant III, what is the sign of $\tan\theta$?

**a.** **positive ✓**

**b.** undefined

**c.** negative

**d.** zero

In Quadrant III, $\tan\theta$ is positive.

**35** [Intermediate] • Target: The unit circle and radian measure

If $\theta$ is an angle in Quadrant II, what is the sign of $\tan\theta$?

**a.** undefined

**b.** **negative ✓**

**c.** zero

**d.** positive

In Quadrant II, $\tan\theta$ is negative.

**36** [Intermediate] • Target: Basic trigonometric identities

If $\theta$ is an angle in Quadrant IV, what is the sign of $\sin\theta$?

**a.** **negative ✓**

**b.** zero

**c.** undefined

**d.** positive

In Quadrant IV, $\sin\theta$ is negative.

**37** [Intermediate] • Target: Graphs of sine and cosine functions

If $\theta$ is an angle in Quadrant I, what is the sign of $\tan\theta$?

**a.** **positive ✓**

**b.** undefined

**c.** negative

**d.** zero

In Quadrant I, $\tan\theta$ is positive.

## Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

© ClipSAT — clipsat.org | All Rights Reserved

**38** [Challenging] • Target: Law of Sines • Cross-Lesson: 2.01 Elementary & Intermediate Algebra

In a right triangle, angle $A$ and angle $B$ are acute angles such that $\sin(A) = \frac{7}{11}$. What is the value of $\cos(B)$?

**a.** $\frac{7}{11}$ ✓

**b.** $\frac{4}{11}$

**c.** $\frac{8}{11}$

**d.** $\frac{11}{7}$

In any right triangle, the two acute angles $A$ and $B$ are complementary, meaning $A + B = 90^\circ$ (or $B = 90^\circ - A$). By the cofunction identity: $\cos(B) = \cos(90^\circ - A) = \sin(A)$ Since $\sin(A) = \frac{7}{11}$, we have $\cos(B) = \frac{7}{11}$.

**39** [Challenging] • Target: Law of Cosines • Cross-Lesson: 3.01 Functions & Graphs

What is the domain of the real-valued function $f(x) = \sqrt{2x - 14}$?

**a.** $x \geq -7$

**b.** $x \leq 7$

**c.** $x > 7$

**d.** $x \geq 7$ ✓

For the function $f(x) = \sqrt{2x - 14}$ to produce real values, the radicand must be non-negative: $2x - 14 \geq 0$ Add $14$ to both sides: $2x \geq 14$ Divide by $2$: $x \geq \frac{14}{2} = 7$ Thus, the domain is all real numbers such that $x \geq 7$.

**40** [Challenging] • Target: Trigonometric equations • Cross-Lesson: 4.01 Coordinate Geometry

What is the radian measure of an angle of $330^\circ$?

**a.** $\frac{11\pi}{6}$ ✓

**b.** $\frac{\pi}{33}$

**c.** $\frac{330\pi}{360}$

**d.** $\frac{330}{180\pi}$

To convert an angle from degrees to radians, multiply by $\frac{\pi}{180^\circ}$: $\text{Radians} = 330^\circ \times \frac{\pi}{180^\circ} = \frac{330\pi}{180} = \frac{11\pi}{6}$ .

**41** [Challenging] • Target: Applications of trigonometry to real-world problems • Cross-Lesson: 5.01 Plane Geometry

In a right triangle $ABC$ where angle $C = 90^\circ$, if $\cos(A) = \frac{12}{13}$, what is the value of $\tan(A)$?

**a.** $\frac{5}{12}$ ✓

**b.** $\frac{12}{5}$

**c.** $\frac{5}{13}$

**d.** $\frac{13}{12}$

Using $\sin^2(A) + \cos^2(A) = 1$, $\sin(A) = \sqrt{1 - (12/13)^2} = \frac{5}{13}$. Then $\tan(A) = \frac{\sin(A)}{\cos(A)} = \frac{5/13}{12/13} = \frac{5}{12}$.

© ClipSAT — clipsat.org | All Rights Reserved

**42** [Challenging] • Target: Reciprocal trigonometric functions • Cross-Lesson: 7.01 Statistics & Probability

A point $P(x, y)$ on the unit circle corresponds to an angle of $\theta$. If $\cos(\theta) = -\frac{3}{5}$ and $\theta$ is in Quadrant II, what is $\sin(\theta)$?

**a.** $-\frac{4}{5}$

**b.** $\frac{4}{5}$ ✓

**c.** $-\frac{3}{4}$

**d.** $\frac{3}{4}$

In Quadrant II, sine is positive: $\sin(\theta) = \sqrt{1 - \cos^2(\theta)} = \sqrt{1 - \left(-\frac{3}{5}\right)^2} = \sqrt{\frac{16}{25}} = \frac{4}{5}$.

**43** [Challenging] • Target: Angle of elevation and depression • Cross-Lesson: 8.01 Matrices, Vectors & Complex Numbers

An inscribed angle in a circle intercepts an arc with a measure of $110^\circ$. What is the measure of the inscribed angle?

**a.** 45°

**b.** **55°** ✓

**c.** 70°

**d.** 110°

By the Inscribed Angle Theorem, the measure of an inscribed angle is half the measure of its intercepted arc: $\frac{110^\circ}{2} = 55^\circ$.

**44** [Challenging] • Target: Solving trigonometric equations over a restricted domain • Cross-Lesson: 1.01 Numbers, Percents & Ratios

A right triangle has legs of length $5\sqrt{3}$ and 5. What is the measure of the smallest angle in degrees?

**a.** $30^\circ$ ✓

**b.** $45^\circ$

**c.** $60^\circ$

**d.** $15^\circ$

Tangent of smallest angle = $5/(5\sqrt{3}) = 1/\sqrt{3} \Rightarrow 30^\circ$.

**45** [Challenging] • Target: A multi-part triangle problem combining Law of Sines and Law of Cosines • Cross-Lesson: 2.01 Elementary & Intermediate Algebra

What is the measure of each interior angle of an equilateral triangle?

**a.** 45°

**b.** **60°** ✓

**c.** 75°

**d.** 90°

All angles in an equilateral triangle are equal: $180^\circ/3 = 60^\circ$.

© ClipSAT — clipsat.org | All Rights Reserved