
Polynomial End Behavior & Zeros

ClipSAT Printable Practice Suite (clipsat.org) • Topic 1.01

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

A polynomial $f(x)$ of degree 4 with real coefficients has leading coefficient 2 and zeros at $x = -1$ (multiplicity 2) and $x = 2 + i$. Write $f(x)$ in fully expanded standard form.



2 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

When the polynomial $P(x) = 2x^3 + kx^2 - 7x + 6$ is divided by $(x - 2)$, the remainder is 12. Find the value of k, and determine whether $(x + 3)$ is a factor of $P(x)$.


3 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

Determine the end behavior of $P(x) = -4x^5 + 7x^3 - 2x + 9$ as $x \to -\infty$ and as $x \to +\infty$ using limit notation, and state the maximum number of local extrema.

© ClipSAT — clipsat.org | All Rights Reserved


4 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

The graph of the exponential function $f(x) = a \cdot \log_3(x - h) + k$ is shown below. The function has a vertical asymptote at $x = 3$ and passes through the points $(4, 5)$ and $(12, 7)$. Determine the value of the parameter a.



5 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)


The graph of the quadratic function $f(x) = x^2 - bx + c$ is shown below. The vertex of the parabola is at $(2, -3)$, and it passes through the point $(0, 1)$. Determine the value of the parameter b.



© ClipSAT — clipsat.org | All Rights Reserved


6 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

For the rational function $r(x) = \frac{2x^2 - 2x - 12}{x^2 - 9}$, find: (a) the domain, (b) the coordinates of any removable discontinuity (hole), (c) the vertical and horizontal asymptotes, and (d) the x- and y-intercepts.



7 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

Consider the quadratic function $f(x) = x^2 - 4x + 3$. What are the x-intercepts of the graph of this function?




Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

© ClipSAT — clipsat.org | All Rights Reserved


8 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Consider the polynomial function $f(x) = x^2 - 5x + 6$. What are the roots of the function $f(x)$?



9 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)


The graph of the exponential function $f(x) = 2^x$ is given. A point P on the graph has a y-coordinate of 8. What is the x-coordinate of point P?



© ClipSAT — clipsat.org | All Rights Reserved

10 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Find the average rate of change of the polynomial function $f(x) = 2x^3 - 5x + 4$ over the closed interval $[-1, 2]$.



11 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Solve the rational inequality $\frac{x+4}{x-2} \leq 3$ and express the solution set in interval notation.

12 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Find the equation of the slant (oblique) asymptote of the rational function $g(x) = \frac{3x^2 - 5x + 7}{x - 2}$, and compute $\lim_{x \to 2^+} g(x)$.

13 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

The graph of $y = f(x)$ is transformed into $y = -2f(3x - 6) + 5$. Describe the exact sequence of geometric transformations applied to $f(x)$, and find the image of the point $(3, -1)$ on $y = f(x)$.

14 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Let $z_1 = 2\left(\cos\frac{\pi}{4} + i\sin\frac{\pi}{4}\right)$ and $z_2 = 3\left(\cos\frac{5\pi}{12} + i\sin\frac{5\pi}{12}\right)$. Find the product $z_1 z_2$ in rectangular form $a + bi$, and compute z_1^6 using De Moivre's Theorem.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 3.02 Polar Coordinates & Polar Graphs

Solve the polynomial inequality $x^3 - 4x^2 + x + 6 \leq 0$.

© ClipSAT — clipsat.org | All Rights Reserved

16 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 4.01 Inverse & Composite Functions

A geometric sequence has third term $g_3 = 18$ and sixth term $g_6 = -\frac{16}{3}$. Find the common ratio r, the first term g_1, and the infinite sum S_∞.

17 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 4.02 Piecewise-Defined & Implicit Functions

A log-log regression plot of $(\ln x, \ln y)$ produces the linear fit $\ln(\hat{y}) = \ln(5) - 2.5\ln(x)$ for $x > 0$. Write $\hat{y}$ as an explicit power function of x, and find the factor by which $\hat{y}$ changes when x is multiplied by 4.

18 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 4.03 Parametric Equations & Modeling


Use the Binomial Theorem to find the coefficient of x^3y^2 in the expansion of $(2x - 3y)^5$.

19 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 1.02 Rational Functions & Asymptotes

Find the 12th term and the sum of the first 25 terms of the arithmetic sequence with $a_1 = -7$ and common difference $d = 4$.

20 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 1.03 Rates of Change & Concavity

A point P is on the unit circle such that its terminal ray makes an angle of θ with the positive x-axis. The coordinates of P are $\left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$. If a point Q on the unit circle has an angle ϕ such that $\phi = \theta + \frac{\pi}{3}$, what is the y-coordinate of point Q?



Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

© ClipSAT — clipsat.org | All Rights Reserved

21 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

Determine the end behavior of the polynomial function $f(x) = -3x^5 + 7x^3 - 2x + 9$.

a. As $x \to \infty$, $f(x) \to -\infty$; as $x \to -\infty$, $f(x) \to \infty$

b. As $x \to \infty$, $f(x) \to \infty$; as $x \to -\infty$, $f(x) \to -\infty$

c. As $x \to \infty$, $f(x) \to -\infty$; as $x \to -\infty$, $f(x) \to -\infty$

d. As $x \to \infty$, $f(x) \to \infty$; as $x \to -\infty$, $f(x) \to \infty$

22 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

If a polynomial $p(x)$ satisfies $\lim_{x \to \infty} p(x) = -\infty$ and $\lim_{x \to -\infty} p(x) = -\infty$, which of the following could be the leading term of $p(x)$?

a. $-4x^6$

b. $2x^6$

c. $-5x^7$

d. $3x^5$

23 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

Consider $g(x) = (2 - x)(x + 3)^2(2x - 1)$. What is the leading coefficient of $g(x)$?

a. -2

b. 2

c. -1

d. 4

24 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

Let $f(x) = (x - 4)^3(x + 2)^2(x - 1)$. At which zero does the graph of f touch the x-axis and turn around without crossing?

a. $x = -2$

b. $x = 4$

c. $x = 1$

d. $x = 2$

© ClipSAT — clipsat.org | All Rights Reserved

25 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

A polynomial $p(x)$ of degree 4 has a zero at $x = 3$ of multiplicity 2, and no other real zeros. If $p(0) = 18$, which equation represents $p(x)$ if its remaining zeros are non-real complex conjugates $\pm i$?

a. $p(x) = 2(x - 3)^2(x^2 + 1)$

b. $p(x) = (x - 3)^2(x^2 + 1)$

c. $p(x) = 2(x - 3)^2(x^2 - 1)$

d. $p(x) = \frac{1}{2}(x - 3)^2(x^2 + 1)$

26 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

At $x = c$, $f(x) = (x - c)^3 g(x)$ where $g(c) \neq 0$. What is the behavior of the graph of f at $(c, 0)$?

a. The graph crosses the x-axis with an inflection/flattening point (horizontal tangent)

b. The graph touches the x-axis and turns around symmetrically

c. The graph has a vertical asymptote at $x = c$

d. The graph crosses the x-axis linearly without flattening

27 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

When $f(x) = 2x^3 - 5x^2 + kx - 8$ is divided by $(x - 3)$, the remainder is 7. Find the value of k.

a. -4

b. 4

c. -2

d. 5

28 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

Which of the following is a factor of $P(x) = x^3 - 6x^2 + 11x - 6$?

a. $x - 2$

b. $x + 2$

c. $x - 4$

d. $x + 1$

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

© ClipSAT — clipsat.org | All Rights Reserved

29 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A polynomial $q(x)$ satisfies $q(-4) = 0$, $q(1) = 0$, and $q(5) = 12$. Which statement must be true?

a. $(x + 4)$ and $(x - 1)$ are factors of $q(x)$

b. $(x - 4)$ and $(x + 1)$ are factors of $q(x)$

c. The remainder when $q(x)$ is divided by $(x + 5)$ is 12

d. $q(x)$ must be a quadratic polynomial

30 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

According to the Rational Root Theorem, which of the following is NOT a possible rational zero of $f(x) = 3x^3 - 2x^2 + 5x - 4$?

a. $\frac{3}{2}$

b. $\frac{4}{3}$

c. ± 1

d. ± 2

31 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A polynomial with real coefficients has degree 3. Two of its zeros are 4 and $2 - 3i$. What is the third zero?

a. $2 + 3i$

b. $-2 + 3i$

c. -4

d. $-2 - 3i$

32 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

What is the total number of complex zeros (counting multiplicity) of $P(x) = (x^2 + 4)^2(x - 5)^3$?

a. 7

b. 5

c. 4

d. 3

33 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

Consider a cubic polynomial $P(x) = ax^3 + bx^2 + cx + d$. The graph of $P(x)$ has a y-intercept at $(0, 12)$. It is known that $P(x)$ has a local maximum at $x = -1$ and a local minimum at $x = 3$. If the value of the local maximum is 20, determine the value of the constant term d of the polynomial.

a. 12

b. 10

c. 8

d. 6

© ClipSAT — clipsat.org | All Rights Reserved

34 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)**

For the polynomial $f(x) = -1x^3 + 3x - 1$, describe the end behavior of f.

a. down on the left, up on the right (as $x \to -\infty$, $f(x) \to -\infty$; as $x \to +\infty$, $f(x) \to +\infty$)

b. down on both ends (as $x \to -\infty$, $f(x) \to -\infty$; as $x \to +\infty$, $f(x) \to -\infty$)

c. up on the left, down on the right (as $x \to -\infty$, $f(x) \to +\infty$; as $x \to +\infty$, $f(x) \to -\infty$)

d. up on both ends (as $x \to -\infty$, $f(x) \to +\infty$; as $x \to +\infty$, $f(x) \to +\infty$)

35 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)**

The polynomial $p(x) = x^4 - 5x^2 + 4$. Which of the following are all its real zeros?

a. $x = \pm 1, \pm 2$

b. $x = \pm 2$

c. $x = 1, 2$

d. $x = \pm 1$

36 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)**

How many real zeros can a degree-3 polynomial have?

a. Exactly 3

b. 1 or 3


c. 0 or 2

d. 2 or 4

© ClipSAT — clipsat.org | All Rights Reserved

37 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)**

Consider the function $f(x) = x^3 - 6x^2 + 9x + 1$. A line tangent to the graph of $f(x)$ at $x = 1$ intersects the graph at another point P. Determine the x-coordinate of point P.



a. 4

b. 5

c. 3


d. 2

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

© ClipSAT — clipsat.org | All Rights Reserved

38 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 4.03 Parametric Equations & Modeling

Consider the quadratic function $f(x) = x^2 - 6x + 5$. What are the x-intercepts of the graph of $f(x)$?



a. x = 1 and x = 5


b. x = -1 and x = -5

c. x = 1 and x = -5

d. x = -1 and x = 5

39 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 1.02 Rational Functions & Asymptotes

The graph of the quadratic function $f(x) = x^2 - 4x + 3$ is shown below. The graph intersects the x-axis at two points. What is the y-coordinate of the vertex of this parabola?



a. -1

b. 1


c. -2

d. 0

© ClipSAT — clipsat.org | All Rights Reserved

40 [Challenging] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14) • Cross-Lesson: 1.03 Rates of Change & Concavity

The graph of a quadratic function $f(x) = ax^2 + bx + c$ is shown below. The vertex of the parabola is at $(2, -4)$, and the function passes through the point $(4, 0)$. Determine the value of the coefficient a.



a. 1

b. -1

c. 2

d. -2

41 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 2.01 Exponential Functions & Growth Models

The graph of the cubic polynomial $P(x) = x^3 + ax^2 + bx + c$ has a local maximum at $(-1, 8)$ and a local minimum at $(3, -24)$. Find the value of a.

a. -3

b. -1

c. 1

d. 3

42 [Challenging] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28) • Cross-Lesson: 2.02 Logarithmic Functions & Properties

Consider the polynomial function $f(x) = x^3 - 5x^2 + 2x + 8$. What is the value of $f(2)$?

a. -4

b. 0

c. 4

d. 8

© ClipSAT — clipsat.org | All Rights Reserved

43 [Challenging] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37) • Cross-Lesson: 3.01 Trigonometric Functions & the Unit Circle

By Rational Root Theorem, possible rational roots of $2x^3 - 5x + 3$ include:

a. ± 6

b. $\pm \frac{3}{2}$

c. ± 4

d. ± 7

44 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 3.02 Polar Coordinates & Polar Graphs

Complex roots of polynomials with real coefficients come in:

a. Pairs of same sign

b. Conjugate pairs

c. Groups of three

d. No pattern

45 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 4.01 Inverse & Composite Functions

Vertical asymptote of $\frac{x+1}{x-4}$:

a. $x = 1$

b. $x = 4$

c. $x = -1$

d. $x = -4$

© ClipSAT — clipsat.org | All Rights Reserved