Sampling Distributions & Central Limit Theorem

ClipSAT Printable Practice Suite (clipsat.org) • Topic 3.04

ANSWER KEY — COMPLETE WORKED SOLUTIONS

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 **[Foundational] • Target: What is a sampling distribution?**

Explain the difference between a population distribution and the sampling distribution of a sample mean $\bar{x}$.

The population distribution describes the values of individual members of the entire population. The sampling distribution of $\bar{x}$ describes the values that the sample mean would take if you repeatedly took many samples of the same size from that population and computed the mean of each sample — it describes variability between different samples' means, not between individuals.

2 **[Foundational] • Target: What is a sampling distribution?**

Explain why the sampling distribution of $\bar{x}$ has less variability than the original population distribution.

A sample mean averages together multiple individual values, and averaging tends to smooth out extreme individual values — some will be higher and some lower than the true mean, partially canceling out. This averaging effect makes sample means cluster more tightly around the population mean than individual values do.

3 **[Foundational] • Target: Mean and standard deviation of the sampling distribution**

A population has mean $\mu = 60$ and standard deviation $\sigma = 12$. For samples of size $n = 36$, find the mean and standard deviation of the sampling distribution of $\bar{x}$.

Mean of $\bar{x}$: $\mu_{\bar{x}} = \mu = 60$ (unbiased — the sampling distribution's mean equals the population mean). Standard deviation of $\bar{x}$ (standard error): $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{12}{\sqrt{36}} = \frac{12}{6} = 2$.

4 **[Foundational] • Target: Mean and standard deviation of the sampling distribution**

Using the same population, find the standard error for a larger sample of size $n = 144$, and explain what happens to the standard error as sample size increases.

$\sigma_{\bar{x}} = \frac{12}{\sqrt{144}} = \frac{12}{12} = 1$. As sample size increases, the standard error decreases (specifically, it decreases proportionally to $\frac{1}{\sqrt{n}}$) — larger samples give more precise (less variable) estimates of the population mean.

© ClipSAT — clipsat.org | All Rights Reserved

5 [Foundational] • Target: The Central Limit Theorem

State the Central Limit Theorem in your own words, including what happens to the shape of the sampling distribution of $\bar{x}$ as sample size n increases.

The Central Limit Theorem states that as sample size n increases, the sampling distribution of the sample mean $\bar{x}$ becomes approximately normal, regardless of the shape of the original population distribution — even if the population itself is skewed or unusual in shape. A commonly used guideline is that $n \geq 30$ is generally large enough for this approximation to be reasonable.

6 [Foundational] • Target: The Central Limit Theorem

Explain why the Central Limit Theorem is considered one of the most important results in statistics, referencing its practical usefulness.

The Central Limit Theorem is powerful because it allows statisticians to use normal-distribution-based methods (like z-scores and confidence intervals) for the sample mean even when the underlying population distribution is unknown or clearly not normal — as long as the sample size is reasonably large, this makes a huge range of inference procedures broadly applicable.

7 [Foundational] • Target: Applying the Central Limit Theorem

A population of package weights has mean $\mu = 50$ lbs and standard deviation $\sigma = 8$ lbs (population shape unknown). For a random sample of $n = 64$ packages, find the probability the sample mean weight exceeds 52 lbs, given $P(Z > 2) \approx 0.0228$.

Since $n = 64 \geq 30$, the CLT justifies treating $\bar{x}$ as approximately normal. $\sigma_{\bar{x}} = \frac{8}{\sqrt{64}} = \frac{8}{8} = 1$. $z = \frac{52 - 50}{1} = 2$. $P(\bar{x} > 52) = P(Z > 2) \approx 0.0228$, or about 2.28%.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

8 [Intermediate] • Target: Sampling distribution of a sample proportion

A population has a true proportion $p = 0.4$ favoring a proposal. For samples of size $n = 100$, find the mean and standard deviation of the sampling distribution of the sample proportion $\hat{p}$, using $\sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$.

Mean: $\mu_{\hat{p}} = p = 0.4$. Standard deviation: $\sigma_{\hat{p}} = \sqrt{\frac{0.4(0.6)}{100}} = \sqrt{\frac{0.24}{100}} = \sqrt{0.0024} \approx 0.049$.

9 [Intermediate] • Target: Sampling distribution of a sample proportion

State the two conditions typically required to treat the sampling distribution of $\hat{p}$ as approximately normal.

Both $np \geq 10$ and $n(1 - p) \geq 10$ must hold (sometimes stated as needing at least 10 expected successes and 10 expected failures) — this ensures the sampling distribution of $\hat{p}$ is not too skewed to approximate with a normal curve.

© ClipSAT — clipsat.org | All Rights Reserved

10 **[Intermediate]** • **Target: Comparing sampling variability across sample sizes**

Explain, in the context of opinion polling, why a poll of 1000 people gives a more reliable estimate of a population proportion than a poll of 50 people, connecting your answer to the formula for $\sigma_{\hat{p}}$.

Since $\sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$ decreases as n increases, a larger sample size produces a smaller standard error for the sample proportion — meaning the sample proportion from a poll of 1000 people tends to land closer to the true population proportion, on average, than a poll of only 50 people, which is subject to much more sample-to-sample variability.

11 **[Intermediate]** • **Target: Applying the sampling distribution of the sample mean**

A population of commute times has mean $\mu = 32$ minutes and standard deviation $\sigma = 10$ minutes. For a random sample of $n = 25$ commuters, find the probability the sample mean commute time exceeds 35 minutes, given $P(Z > 1.5) \approx 0.0668$.

$\sigma_{\bar{x}} = \frac{10}{\sqrt{25}} = \frac{10}{5} = 2$. $z = \frac{35-32}{2} = 1.5$. $P(\bar{x} > 35) = P(Z > 1.5) \approx 0.0668$, or about 6.68%.

12 **[Intermediate]** • **Target: Sample size and the CLT threshold**

A population of insurance claim amounts is strongly right-skewed. Explain why a sample of size $n = 15$ might not be large enough to safely use CLT-based normal methods, but a sample of size $n = 200$ likely would be.

The Central Limit Theorem's normal approximation improves as sample size increases, and it takes a larger sample size to overcome a strongly skewed population shape. With only $n = 15$, the sampling distribution of $\bar{x}$ may still noticeably reflect the skewness of the original population, making normal-based methods unreliable. With $n = 200$ (well beyond the common $n \geq 30$ guideline), the sampling distribution of $\bar{x}$ is much more likely to be close to normal, even though the original population itself is.

13 **[Intermediate]** • **Target: Standard error vs. standard deviation**

Explain the difference between the standard deviation of an individual population value and the standard error of a sample mean $\bar{x}$.

The standard deviation (σ) measures the spread of individual values within the population. The standard error ($\sigma_{\bar{x}} = \sigma/\sqrt{n}$) measures the spread of sample means across many hypothetical samples of the same size — it is always smaller than σ (for $n > 1$), since averaging reduces variability.

© ClipSAT — clipsat.org | All Rights Reserved

14 **[Intermediate]** • **Target: Standard error vs. standard deviation**

A quality control engineer at a bottling plant knows that individual bottle fill volumes are approximately normal with mean 500 mL and standard deviation 8 mL. Every hour, the engineer measures a sample of 16 bottles and computes the sample mean fill volume. Describe the shape, mean, and standard deviation of the sampling distribution of this hourly sample mean, and explain whether the Central Limit Theorem's usual $n \geq 30$ guideline is even needed to justify treating this sampling distribution as normal.

Shape: approximately normal. Mean: $\mu_{\bar{x}} = 500$ mL (same as the population mean). Standard deviation: $\sigma_{\bar{x}} = \frac{8}{\sqrt{16}} = \frac{8}{4} = 2$ mL. The $n \geq 30$ CLT guideline is not needed here, because the underlying population of individual fill volumes is already stated to be approximately normal — when the population itself is normal, the sampling distribution of $\bar{x}$ is exactly normal for any sample size, including small ones like $n = 16$.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 **[Challenging]** • **Target: Standard error vs. standard deviation** • **Cross-Lesson: 1.03 Describing Distributions & Linear Regression**

A political pollster samples 900 voters and calculates a sample proportion $\hat{p} = 0.53$ supporting a ballot measure, from a true population proportion of $p = 0.50$. Verify that both conditions ($np \geq 10$ and $n(1-p) \geq 10$) are satisfied for treating the sampling distribution of $\hat{p}$ as approximately normal, then find the standard deviation of this sampling distribution, to four decimal places.

$np = 900(0.50) = 450 \geq 10$ ✓. $n(1-p) = 900(0.50) = 450 \geq 10$ ✓. Both conditions are satisfied. Standard deviation: $\sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}} = \sqrt{\frac{(0.50)(0.50)}{900}} = \sqrt{\frac{0.25}{900}} \approx 0.0167$.

16 **[Challenging]** • **Target: Standard error vs. standard deviation** • **Cross-Lesson: 2.01 Sampling Methods & Bias**

A large factory produces light bulbs whose individual lifespans are strongly right-skewed (most bulbs fail early, but a few last an unusually long time), with population mean $\mu = 800$ hours and standard deviation $\sigma = 250$ hours. An engineer takes a random sample of $n = 64$ bulbs and computes the sample mean lifespan. Explain, step by step, why it is reasonable to treat this sampling distribution of $\bar{x}$ as approximately normal despite the population itself being skewed, then compute the standard error of $\bar{x}$.

Even though the individual bulb lifespans are strongly skewed, the Central Limit Theorem states that the sampling distribution of the sample mean becomes approximately normal as sample size increases, regardless of the population's original shape. Since $n = 64$ comfortably exceeds the common guideline of $n \geq 30$, it is reasonable to treat $\bar{x}$'s sampling distribution as approximately normal here. Standard error: $\sigma_{\bar{x}} = \frac{250}{\sqrt{64}} = \frac{250}{8} = 31.25$ hours.

© ClipSAT — clipsat.org | All Rights Reserved

17 **[Challenging] • Target: Standard error vs. standard deviation • Cross-Lesson: 2.02 Experimental Design & Scope of Inference**

An online retailer knows that 22% of all website visitors make a purchase. On a particular day, the retailer records the browsing sessions of a random sample of 500 visitors and calculates the sample proportion who made a purchase. Verify the two conditions needed to treat this sampling distribution as approximately normal, then compute the standard deviation of the sampling distribution of the sample proportion, to four decimal places, and describe what this standard deviation represents in the context of comparing many different days' worth of visitor samples.

$np = 500(0.22) = 110 \geq 10$ ✓. $n(1-p) = 500(0.78) = 390 \geq 10$ ✓. Both conditions are satisfied. $\sigma_{\hat{p}} = \sqrt{\frac{(0.22)(0.78)}{500}} = \sqrt{\frac{0.1716}{500}} = \sqrt{0.0003432} \approx 0.0185$. This standard deviation describes how much the sample proportion of purchasers would typically vary from one day's random sample of 500 visitors to another, if the true purchase rate stayed fixed at 22% — most daily sample proportions would fall within roughly ± 0.0185 of 0.22.

18 **[Challenging] • Target: Standard error vs. standard deviation • Cross-Lesson: 3.01 Probability Rules & Conditional Probability**

Level 2 · MediumExplain the difference between a single-blind experiment and a double-blind experiment, and describe why a placebo is used in medical trials.

Single-Blind: Either the subjects do not know which treatment they are receiving, OR the evaluators who measure the response do not know which treatment was administered. Double-Blind: NEITHER the subjects nor the researchers/evaluators interacting with them and measuring response variables know which treatment was administered. Placebo Purpose: A placebo provides an inactive comparison baseline that isolates the physiological effect of the active medication from psychological improvements caused simply by receivin.

19 **[Challenging] • Target: What is a sampling distribution? • Cross-Lesson: 3.02 Discrete Random Variables & Binomial Distributions**

Level 3 · AdvancedTen athletes participate in a vertical jump training program. Their jump heights (in inches) before and after the program are recorded. To test whether the program increased jump height, explain why a paired t-test is appropriate rather than a two-sample t-test, state the null and alternative hypotheses, and specify the degrees of freedom.

Why Paired: The observations are paired on the same subjects (before and after measurements on each athlete). The two samples are dependent, not independent. Hypotheses: Letting $\mu_d = \mu_{\text{after}} - \mu_{\text{before}}$ be the true mean difference in jump height: $H_0 : \mu_d = 0 \quad \text{vs.} \quad H_a : \mu_d > 0$. Degrees of Freedom: Since this is a one-sample test on differences: $df = n - 1 = 10 - 1 = \mathbf{9}$.

© ClipSAT — clipsat.org | All Rights Reserved

20 [Challenging] • Target: Mean and standard deviation of the sampling distribution • Cross-Lesson: 3.03 The Normal Distribution

Level 2 · MediumFind the critical value z^* required to construct a 98% confidence interval for a population proportion, rounded to three decimal places.

For a 98% confidence interval, the central area is 0.98, leaving 0.02 split equally between both tails (0.01 in each tail). The area to the left of the upper critical value is: $1 - 0.01 = 0.99$. Using the inverse normal function: $z^* = \text{invNorm}(\text{area} = 0.99, \mu = 0, \sigma = 1) \approx \mathbf{2.326}$.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

21 [Foundational] • Target: What is a sampling distribution?

A population has mean $\mu = 100$ and standard deviation $\sigma = 60$. For samples of size $n = 100$, find the standard deviation of the sampling distribution of $\bar{x}$ (the standard error), using $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$.

a. 6 ✓

b. 5

c. 7

d. 60

$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \frac{60}{\sqrt{100}} = \frac{60}{10} = 6.$

22 [Foundational] • Target: Mean and standard deviation of the sampling distribution

A population has a true proportion $p = 0.40$. For samples of size $n = 200$, find the standard deviation of the sampling distribution of the sample proportion $\hat{p}$, to four decimal places, using $\sigma_{\hat{p}} = \sqrt{p(1-p)/n}$.

a. 0.0346 ✓

b. 0.0246

c. 0.2400

d. 0.0446

$\frac{0.40(0.60)}{200} = 0.0012$. $\sigma_{\hat{p}} = \sqrt{0.0012} \approx 0.0346.$

© ClipSAT — clipsat.org | All Rights Reserved

23 [Foundational] • Target: The Central Limit Theorem

A population has mean $\mu = 143$ and standard deviation $\sigma = 90$ (population shape unknown). For a random sample of $n = 36$, find the probability the sample mean is above 158, using the Central Limit Theorem.

a. 11.87%

b. 15.87% ✓

c. 84.13%

d. 19.87%

Since $n = 36 \geq 30$, the CLT justifies treating $\bar{x}$ as approximately normal. $\sigma_{\bar{x}} = 90/\sqrt{36} = 15$. $z = (158 - 143)/15 = 1$. $P(\bar{x} > 158) = P(Z > 1) \approx 15.87\%$.

24 [Foundational] • Target: Applying the Central Limit Theorem

A population has mean $\mu = 48$ and standard deviation $\sigma = 132$ (population shape unknown). For a random sample of $n = 144$, find the probability the sample mean is below 81, using the Central Limit Theorem.

a. 96.87%

b. 95.87%

c. 0.13%

d. 99.87% ✓

Since $n = 144 \geq 30$, the CLT justifies treating $\bar{x}$ as approximately normal. $\sigma_{\bar{x}} = 132/\sqrt{144} = 11$. $z = (81 - 48)/11 = 3$. $P(\bar{x} < 81) = P(Z < 3) \approx 99.87\%$.

25 [Foundational] • Target: Sampling distribution of a sample proportion

A pollster samples $n = 500$ voters, from a population with true proportion $p = 0.20$ supporting a measure. Verify that both conditions ($np \geq 10$ and $n(1-p) \geq 10$) are satisfied, then find the standard deviation of the sampling distribution of $\hat{p}$, to four decimal places.

a. 0.0259

b. 0.2000

c. 0.0179 ✓

d. 0.0099

$np = 500(0.20) = 100.00 \geq 10$. $n(1-p) = 500(0.80) = 400.00 \geq 10$. Both conditions hold. $p(1-p)/n = 0.20(0.80)/500 = 0.0003$. $\sigma_{\hat{p}} = \sqrt{0.0003} \approx 0.0179$.

© ClipSAT — clipsat.org | All Rights Reserved

26 [Foundational] • Target: Comparing sampling variability across sample sizes

A population has standard deviation $\sigma = 160$. Find the sample size n needed so that the standard error $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$ equals 20.

a. 72

b. 64 ✓

c. 56

d. 8

$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$: $20 = \frac{160}{\sqrt{n}}$, so $\sqrt{n} = \frac{160}{20} = 8$, giving $n = 8^2 = 64$.

27 [Foundational] • Target: Applying the sampling distribution of the sample mean

A researcher is studying the average height of adult males in a city. They collected a sample of 100 adult males and found the sample mean height to be 175 cm with a sample standard deviation of 8 cm. If the margin of error for a 95% confidence interval is calculated to be approximately 1.57 cm, what is the minimum sample size required to achieve a margin of error of at most 1 cm, assuming the sample standard deviation remains the same?

a. 154

b. 246

c. **197** ✓

d. 97

The formula for the margin of error (ME) for a confidence interval for the mean is given by $ME = z^* \frac{\sigma}{\sqrt{n}}$, where z^* is the critical value for the desired confidence level, σ is the population standard deviation (or sample standard deviation as an estimate), and n is the sample size. For a 95% confidence interval, the critical value z^* is approximately 1.96. From the given information, we have $ME = 1.57$ cm, $\sigma \approx 8$ cm, and $n = 100$. We can verify the given margin of.

28 [Foundational] • Target: Sample size and the CLT threshold

A sample of size $n = 64$ has a sample mean of 80 and standard deviation 16. What is the standard error of the mean?

a. 0.25

b. 1.0

c. **2.0** ✓

d. 4.0

Standard error of the mean $\text{SE} = \frac{s}{\sqrt{n}} = \frac{16}{\sqrt{64}} = \frac{16}{8} = 2.0$.

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

© ClipSAT — clipsat.org | All Rights Reserved

29 [Intermediate] • Target: Standard error vs. standard deviation

A researcher wants a 95% confidence interval for a proportion with margin of error at most 0.043, using a planning value $p^* = 0.4$. What sample size n is required ($z^* = 1.96$)?

a. 529

b. **499 ✓**

c. 484

d. 469

$n = \left(\frac{z^*}{ME}\right)^2 p^*(1-p^*) = \left(\frac{1.96}{0.043}\right)^2 (0.4)(0.6) = 498.64$, round up to $n = 499$.

30 [Intermediate] • Target: What is a sampling distribution?

The length, X mm, of components manufactured by an industrial process is normally distributed with mean μ mm and standard deviation σ mm. A quality assurance engineer tests the null hypothesis $H_0 : \mu = 20.0$ against the alternative hypothesis $H_1 : \mu > 20.0$ at the 5% significance level using an initial random sample of size $n = 25$. The critical region for the sample mean $\bar{X}$ is found to be $\bar{X} \geq 20.658$. Using the standard normal critical value $z_{0.05} = 1.645$: 1. Find the value of σ. 2. Hence, determine the minimum sample size, m, required so that in a subsequent test of $H_0 : \mu = 20.0$ against $H_1 : \mu > 20.0$ at the same 5% significance level, the probability of committing a Type II error when the true mean is $\mu = 21.0$ is at most 0.05.

a. **44 ✓**

b. 43

c. 11

d. 22

We solve the problem in two stages: **Step 1: Determine σ** Under H_0, the distribution of the sample mean is $\bar{X} \sim N\left(20.0, \frac{\sigma^2}{25}\right)$. The critical value for a upper one-tailed test at the 5% level corresponds to $z = 1.645$: $\frac{20.658 - 20.0}{\frac{\sigma}{\sqrt{25}}} = 1.645$ $\frac{0.658}{\frac{\sigma}{5}} = 1.645 \Rightarrow \frac{3.29}{\sigma} = 1.645 \Rightarrow \sigma = \frac{3.29}{1.645} = 2.0$ mm **Step 2: Determine the minimum sample size m** For.

© ClipSAT — clipsat.org | All Rights Reserved

31 [Intermediate] • Target: Mean and standard deviation of the sampling distribution

A quality control process monitors the weight, W grams, of chocolate bars produced by a machine. A random sample of 50 chocolate bars was taken, and their weights were recorded. The sample mean weight was 102.5 grams, and the sample standard deviation was 3.5 grams. A) Calculate the standard error of the mean. B) Construct a 95% confidence interval for the mean weight of the chocolate bars. C) The machine is calibrated to produce chocolate bars with a mean weight of $\mu_0 = 100$ grams. Based on your confidence interval, what conclusion can be drawn about the calibration of the machine?

a. The machine is likely calibrated correctly, as μ_0 is within the interval.

b. **The machine is likely not calibrated correctly, as μ_0 is outside the interval.** ✓

c. The machine is likely calibrated correctly, as the sample mean is close to μ_0.

d. The machine is likely not calibrated correctly, as the sample standard deviation is too large.

A) The standard error of the mean (SE) is calculated using the formula: $SE = \frac{s}{\sqrt{n}}$, where s is the sample standard deviation and n is the sample size. Given $s = 3.5$ grams and $n = 50$. $SE = \frac{3.5}{\sqrt{50}} \approx \frac{3.5}{7.071} \approx 0.495$ grams. B) To construct a 95% confidence interval for the mean weight, we use the formula: $CI = \bar{x} \pm z^* \times SE$, where $\bar{x}$ is the sample mean and z^* is the critical z-value for a 95% confidence level. For a 95% confidence lev.

32 [Intermediate] • Target: The Central Limit Theorem

A sample of 5 numbers has a mean of 12 and a standard deviation of 3. If every number in the sample is multiplied by 2, what are the new mean and standard deviation?

a. Mean = 24, SD = 3

b. **Mean = 24, SD = 6** ✓

c. Mean = 12, SD = 6

d. Mean = 24, SD = 9

Multiplying every value by a constant $c = 2$ scales both the mean and the standard deviation by 2: new mean = $12 \times 2 = 24$, new SD = $3 \times 2 = 6$.

33 [Intermediate] • Target: Applying the Central Limit Theorem

For a sample of size $n = 30$ (so $df = 29$), find the critical value t^* for a 99% confidence interval for the mean.

a. 3.256

b. **2.756** ✓

c. 3.056

d. 1.956

With $df = 29$ and 99% confidence, $t^* = 2.756$ (from the t-distribution).

© ClipSAT — clipsat.org | All Rights Reserved

34 [Intermediate] • Target: Sampling distribution of a sample proportion

Increasing sample size n primarily:

a. Increases bias

b. **Reduces variability ✓**

c. Increases variability

d. Has no effect on confidence intervals

Larger n → smaller standard error → less variability.

35 [Intermediate] • Target: Comparing sampling variability across sample sizes

A randomised block design blocks subjects by gender into 3 blocks of 57 subjects each, with random assignment to treatments within each block. What is the total sample size?

a. **171 ✓**

b. 156

c. 196

d. 186

Total $= 3 \times 57 = 171$ subjects.

36 [Intermediate] • Target: Applying the sampling distribution of the sample mean

A randomised block design blocks subjects by fitness level into 3 blocks of 20 subjects each, with random assignment to treatments within each block. What is the total sample size?

a. 45

b. 85

c. 50

d. **60 ✓**

Total $= 3 \times 20 = 60$ subjects.

37 [Intermediate] • Target: Sample size and the CLT threshold

A randomised block design blocks subjects by region into 2 blocks of 42 subjects each, with random assignment to treatments within each block. What is the total sample size?

a. 59

b. **84 ✓**

c. 74

d. 109

Total $= 2 \times 42 = 84$ subjects.

© ClipSAT — clipsat.org | All Rights Reserved

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 **[Challenging]** • **Target: Standard error vs. standard deviation** • **Cross-Lesson: 3.01 Probability Rules & Conditional Probability**

A randomised block design blocks subjects by prior experience into 3 blocks of 29 subjects each, with random assignment to treatments within each block. What is the total sample size?

a. 77

b. 97

c. **87** ✓

d. 72

Total $= 3 \times 29 = 87$ subjects.

© ClipSAT — clipsat.org | All Rights Reserved

39 **[Challenging] • Target: What is a sampling distribution? • Cross-Lesson: 3.02 Discrete Random Variables & Binomial Distributions**

A political poll surveyed 1200 likely voters. The results indicated that 55% of respondents plan to vote for Candidate A, with a margin of error of ± 3 percentage points. If the poll is adjusted to include only likely voters who are registered as "strong supporters" (who constitute 60% of the overall surveyed population), and assuming the proportion of "strong supporters" who favor Candidate A remains the same as the overall proportion, what is the most accurate statement regarding the new confidence interval for the proportion of strong supporters who favor Candidate A? Let P be the proportion of all likely voters who plan to vote for Candidate A, and let n be the sample size. The margin of error for a proportion is typically calculated as $z\sqrt{\frac{P(1-P)}{n}}$, where z is the critical value for the desired confidence level. For the initial poll: $P = 0.55$ $n = 1200$ Margin of Error = 0.03 This implies a confidence interval for P of $[0.52, 0.58]$. Now, consider the subset of "strong supporters," which represents 60% of the population. Let P_{ss} be the proportion of strong supporters who favor Candidate A. We are given that the proportion of "strong supporters" who favor Candidate A is assumed to be the same as the overall proportion, so $P_{ss} \approx P = 0.55$. The sample size for this subset, assuming it's representative, would be $n_{ss} = 0.60 \times n = 0.60 \times 1200 = 720$. The new margin of error for this subgroup would be $ME_{ss} = z\sqrt{\frac{P_{ss}(1-P_{ss})}{n_{ss}}}$. Since $n_{ss} < n$, the margin of error will increase (making the interval wider) if the proportion P_{ss} were significantly different, or it would decrease if n_{ss} were larger. However, if the proportion is the same, a smaller sample size will lead to a larger margin of error.

a. The new confidence interval for strong supporters will be wider than the original interval. ✓

b. The new confidence interval for strong supporters will be narrower than the original interval.

c. The new confidence interval for strong supporters will be exactly the same as the original interval.

d. It is impossible to determine if the new confidence interval is wider or narrower without knowing the z-score.

The margin of error for a proportion is given by $ME = z\sqrt{\frac{P(1-P)}{n}}$. In the original poll, $n = 1200$. In the adjusted poll focusing on strong supporters, the effective sample size is $n_{ss} = 0.60 \times 1200 = 720$. Since the proportion P is assumed to be the same for both groups, $P_{ss} \approx P$. The new margin of error is $ME_{ss} = z\sqrt{\frac{P(1-P)}{n_{ss}}}$. Because $n_{ss} = 720\sqrt{\frac{P(1-P)}{n}}$. This means that $ME_{ss} > ME$. A larger margin of error results in a wider conf.

© ClipSAT — clipsat.org | All Rights Reserved

40 [Challenging] • Target: Mean and standard deviation of the sampling distribution • Cross-Lesson: 3.03 The Normal Distribution

A data set of 100 observations has a mean of 50 and standard deviation 10. According to Chebyshev's inequality, at least what percentage of the data must lie between 30 and 70?

a. 50%

b. 68%

c. **75%** ✓

d. 95%

The interval 30 to 70 is $50 \pm 2(10)$, which is $k = 2$ standard deviations from the mean. Chebyshev's inequality guarantees that at least $1 - 1/k^2 = 1 - 1/4 = 3/4 = 75\%$ of data lies within 2 standard deviations for ANY distribution.

41 [Challenging] • Target: The Central Limit Theorem • Cross-Lesson: 4.01 Confidence Intervals & Hypothesis Testing

A researcher wants to estimate a population proportion with a margin of error of at most $\pm 2\%$ at a 95% confidence level. An earlier study with $n = 400$ had a margin of error of $\pm 5\%$. Approximately what sample size n is required?

a. 1,000

b. 1,600

c. **2,500** ✓

d. 4,000

$\text{MOE} \propto 1/\sqrt{n} \Rightarrow \frac{\text{MOE}_1}{\text{MOE}_2} = \sqrt{\frac{n_2}{n_1}} \Rightarrow \frac{5}{2} = \sqrt{\frac{n_2}{400}} \Rightarrow 2.5^2 = \frac{n_2}{400} \Rightarrow 6.25 \times 400 = 2,500.$

42 [Challenging] • Target: Applying the Central Limit Theorem • Cross-Lesson: 4.02 Chi-Square Tests & Regression Inference

A data set X has mean μ and standard deviation σ. A new data set Y is formed by the transformation $y_i = 3x_i - 5$ for each value in X. What are the mean and standard deviation of Y?

a. Mean = $3\mu - 5$, SD = $3\sigma - 5$

b. **Mean =** $3\mu - 5$**, SD =** 3σ ✓

c. Mean = 3μ, SD = 3σ

d. Mean = $\mu - 5$, SD = σ

Linear transformation properties: $E(3X - 5) = 3\mu - 5$. The constant -5 does not affect spread, while multiplication by 3 scales standard deviation: $\text{SD}(3X - 5) = |3|\sigma = 3\sigma$.

© ClipSAT — clipsat.org | All Rights Reserved

43 [Challenging] • Target: Sampling distribution of a sample proportion • Cross-Lesson: 1.01 Measures of Center & Spread

A student takes 4 tests and has a mean score of 85 with standard deviation 0. What were the student's individual test scores?

a. 80, 85, 85, 90

b. **All four scores were 85 ✓**

c. 75, 85, 85, 95

d. 82, 84, 86, 88

A standard deviation of 0 means there is zero variability among the values; every single score must equal the mean of 85.

44 [Challenging] • Target: Comparing sampling variability across sample sizes • Cross-Lesson: 1.02 Z-Scores & Linear Transformations

In estimating a population proportion, what is the effect on the margin of error of increasing the sample size (with the same confidence level), assuming all else is held constant?

a. The margin of error tends to increase

b. **The margin of error tends to decrease ✓**

c. The margin of error stays exactly the same

d. The margin of error becomes zero

Increasing the sample size (with the same confidence level) generally causes: the margin of error tends to decrease.

45 [Challenging] • Target: Applying the sampling distribution of the sample mean • Cross-Lesson: 1.03 Describing Distributions & Linear Regression

In estimating a population proportion, what is the effect on the margin of error of decreasing the sample size (with the same confidence level), assuming all else is held constant?

a. **The margin of error tends to increase ✓**

b. The margin of error tends to decrease

c. The margin of error stays exactly the same

d. The margin of error becomes zero

Decreasing the sample size (with the same confidence level) generally causes: the margin of error tends to increase.

© ClipSAT — clipsat.org | All Rights Reserved