Triangles & Congruence

ClipSAT Printable Practice Suite (clipsat.org) • Topic 3.01

ANSWER KEY — COMPLETE WORKED SOLUTIONS

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 **[Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)**

A data set recorded for review: equations of lines consists of the four values $\{15, 19, 23, 29\}$. What is the arithmetic mean? Show your complete algebraic work.

Final Answer: 21.5. The sum is $15 + 19 + 23 + 29 = 86$, and dividing by 4 gives 21.5.

2 **[Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)**

For the model $f(x) = 6x + 3$ in review: equations of lines, evaluate $f(3)$.

Final Answer: 21. Substitute $x = 3$: $f(3) = 6(3) + 3 = 21$.

3 **[Foundational] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)**

In ΔABC studying equations of parallel and perpendicular lines, $m\angle A = 45^\circ$ and $m\angle B = 45^\circ$. What is $m\angle C$? Show your complete algebraic work.

Final Answer: 90°. Using the triangle angle-sum theorem, $m\angle C = 180^\circ - (45^\circ + 45^\circ) = 90^\circ$.

4 **[Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)**

[Triangles & Congruence · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

5 **[Foundational] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)**

In a $45°$–$45°$–$90°$ right triangle, the hypotenuse has length $10\sqrt{2}$ cm. What is the length of each leg, and what is the area of the triangle?

Step 1: Find leg length s: In a $45°$–$45°$–$90°$ special right triangle, the hypotenuse is $s\sqrt{2}$: $s\sqrt{2} = 10\sqrt{2} \Rightarrow s = 10$ cm. Step 2: Calculate the area: Since both legs form the base and height: Area $= \frac{1}{2}s^2 = \frac{1}{2}(10^2) = \frac{1}{2}(100) = 50$ cm^2.

6 **[Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)**

[Right Triangles & Trigonometry · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

© ClipSAT — clipsat.org | All Rights Reserved

7 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

Each interior angle of a regular polygon measures 144°. How many sides does the polygon have?

Solution: The fastest approach is to find the exterior angle first. Since interior and exterior angles form a linear pair: $\text{Exterior angle} = 180^\circ - 144^\circ = 36^\circ$. Since the exterior angles of any regular n-gon sum to 360°: $n = \frac{360^\circ}{36^\circ} = 10$. The polygon is a regular decagon with 10 sides.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

8 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

In right ΔABC with right angle at C, altitude CD is drawn to hypotenuse AB. If $AD = 4$ and $DB = 9$, find the length of leg BC.

Step 1: Compute total hypotenuse AB: $AB = AD + DB = 4 + 9 = 13$ Step 2: Apply the Geometric Mean Leg Rule: The leg BC is the geometric mean of the hypotenuse AB and the adjacent segment DB: $BC^2 = AB \cdot DB = 13 \cdot 9 = 117$ Step 3: Simplify the radical: $BC = \sqrt{117} = \sqrt{9 \times 13} = 3\sqrt{13} \approx 10.82$

9 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

Points A, B, C lie on a circle with center O. Inscribed angle $\angle ABC$ intercepts minor arc AC and has measure 38°. What is the measure of the central angle $\angle AOC$ intercepting the same arc?

Solution: The measure of a central angle is twice the measure of an inscribed angle that intercepts the same arc: $m\angle AOC = 2 \times m\angle ABC = 2 \times 38^\circ = 76^\circ$. The central angle $\angle AOC$ has measure 76°, and the measure of minor arc AC is also 76°.

10 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

[Similarity & Proportional Reasoning · Practice 4] If the governing relation for this system is $g(t) = 6t + 6$, evaluate $g(10)$ and find the inverse input t when $g(t) = 72$.

Substitute $t = 10$: $g(10) = 6(10) + 6 = \mathbf{66}$. Setting $6t + 6 = 72$ gives $6t = 66 \Rightarrow t = \mathbf{11}$.

11 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

[Circles · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

12 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

[Coordinate Geometry · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

© ClipSAT — clipsat.org | All Rights Reserved

13 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)**

[Foundations & Reasoning · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

14 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)**

[Polygons & Quadrilaterals · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 **[Challenging]** • **Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)** • **Cross-Lesson: 8.01 Area, Surface Area & Volume**

[Area, Surface Area & Volume · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

16 **[Challenging]** • **Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)** • **Cross-Lesson: 9.01 Transformations & Symmetry**

[Transformations & Symmetry · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

17 **[Challenging]** • **Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)** • **Cross-Lesson: 1.01 Foundations & Reasoning**

In a course grading system, Homework is weighted 20%, Midterm Exam is weighted 35%, and Final Exam is weighted 45%. A student scored 92 on Homework and 76 on the Midterm Exam. What is the minimum score the student must achieve on the Final Exam to earn an overall course average of at least 84?

Step 1: Set up the weighted average inequality with final exam score F:
$0.20(92) + 0.35(76) + 0.45(F) \geq 84$. Step 2: Compute the accumulated points:
$0.20(92) = 18.4, \quad 0.35(76) = 26.6 \Rightarrow 18.4 + 26.6 = 45.0$. Step 3: Isolate F:
$45.0 + 0.45F \geq 84 \Rightarrow 0.45F \geq 84 - 45 = 39 \Rightarrow F \geq \frac{39}{0.45} = \frac{3900}{45} = \frac{260}{3} \approx 86.67$. The student must score at least $\mathbf{86.7}$ (or $\mathbf{\frac{260}{3}}$) on the Final Exam.

© ClipSAT — clipsat.org | All Rights Reserved

18 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 10.01 Coordinate Geometry

A bivariate regression model for student preparation hours x and exam grade y yields the line of best fit $\hat{y} = 4.8x + 52$. A student who studied for 8 hours scored 94 on the exam. (a) What is the model's predicted score for this student? (b) Compute and interpret the student's regression residual.

(a) Predicted Score $\hat{y}$: Substitute $x = 8$ into the regression equation:
$\hat{y} = 4.8(8) + 52 = 38.4 + 52 = \mathbf{90.4}$. (b) Residual Calculation & Interpretation:
$\text{Residual} = y - \hat{y} = 94 - 90.4 = \mathbf{+3.6}$. Because the residual is positive (+3.6), the student's actual score was 3.6 points higher than the score predicted by the regression line (the model underestimated the student's performance).

19 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 11.01 Probability

Section I of EST I Math contains 20 questions in 25 minutes. If a student answers the first 12 questions in an average of 50 seconds each, how many seconds per question are available for the remaining 8 questions?

Time for first 12 questions $= 12 \times 50 = 600$ seconds (10 minutes). Total time $= 25 \times 60 = 1500$ seconds. Remaining time $= 1500 - 600 = 900$ seconds. For the remaining 8 questions:
$\frac{900}{8} = 112.5$ seconds per question (1 minute 52.5 seconds per question).

20 [Challenging] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14) • Cross-Lesson: 2.01 Lines & Angles

A student has scores of 78, 85, 92, and 81 on four exams. What score must the student earn on the fifth exam to achieve an overall arithmetic mean of 85 across all five exams?

Step 1: Determine the required sum of all 5 exam scores: $\sum x = 5 \times 85 = 425$. Step 2: Sum the first four exam scores: $78 + 85 + 92 + 81 = 336$. Step 3: Subtract the accumulated total from the required total to find the 5th score x_5: $x_5 = 425 - 336 = \mathbf{89}$. The student must score 89 on the fifth exam.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

21 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

In ΔABC studying parallel lines and transversals, $m\angle A = 42^\circ$ and $m\angle B = 45^\circ$. What is $m\angle C$?

a. 93° ✓

b. 103°

c. 85°

d. 138°

Using the triangle angle-sum theorem, $m\angle C = 180^\circ - (42^\circ + 45^\circ) = 93^\circ$.

© ClipSAT — clipsat.org | All Rights Reserved

22 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

In ΔABC studying proving lines parallel, $m\angle A = 43^\circ$ and $m\angle B = 45^\circ$. What is $m\angle C$?

a. 92° ✓

b. 102°

c. 84°

d. 137°

Using the triangle angle-sum theorem, $m\angle C = 180^\circ - (43^\circ + 45^\circ) = 92^\circ$.

23 [Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

For the model $f(x) = 3x + 5$ in extension: perpendiculars and distance, evaluate $f(3)$.

a. 14 ✓

b. 17

c. 9

d. 18

Substitute $x = 3$: $f(3) = 3(3) + 5 = 14$.

24 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

In a triangle, two angles measure 14° and 101°. What is the measure of the third angle?

a. 101°

b. **65°** ✓

c. 115°

d. 14°

Triangle angles sum to 180°. $180 - 14 - 101 = 65^\circ$.

25 [Foundational] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

An isosceles triangle has a vertex angle of 12°. What is the measure of each base angle?

a. **84°** ✓

b. 6°

c. 12°

d. 168°

Base angles are equal: $(180 - 12)/2 = 84^\circ$.

© ClipSAT — clipsat.org | All Rights Reserved

26 [Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

In ΔABC and ΔXYZ, we are given that $AB = XY$, $BC = YZ$, and $\angle B = \angle Y$. Which of the following additional conditions would NOT be sufficient to prove that $\Delta ABC \cong \Delta XYZ$?

a. $\angle A = \angle X$

b. $AC = XZ$

c. $\angle C = \angle Z$

d. $AB = XZ$ ✓

We are given that two sides and the included angle of ΔABC are equal to the corresponding sides and included angle of ΔXYZ ($AB = XY$, $BC = YZ$, and $\angle B = \angle Y$). This satisfies the SAS (Side-Angle-Side) congruence criterion, which is sufficient to prove that the triangles are congruent. Therefore, any additional condition that, when combined with the given information, results in a valid congruence criterion is not the correct answer to the question asking for a condition that is NOT s.

27 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

In ΔABC, medians $\overline{AD}$ and $\overline{BE}$ intersect at the centroid G. The lengths of the medians are $AD = 12$ and $BE = 9$. Which of the following represents the complete range of all possible lengths for side $\overline{BC}$?

a. $4 < BC < 20$ ✓

b. $2 < BC < 14$

c. $3 < BC < 21$

d. $2 < BC < 10$

1. **Apply the Centroid Median Division Property:** The centroid G divides each median into segments in a $2:1$ ratio, with the longer segment adjacent to the vertex. - For median $\overline{AD}$: $GD = \frac{1}{3}AD = \frac{1}{3}(12) = 4$. - For median $\overline{BE}$: $BG = \frac{2}{3}BE = \frac{2}{3}(9) = 6$. 2. **Apply the Triangle Inequality Theorem to ΔBGD:** In ΔBGD, the three side lengths are $BG = 6$, $GD = 4$, and BD. For a non-degenerate triangle to exist, the third side must sati.

28 [Foundational] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

In triangles ABC and DEF, it is given that $AB \cong DE$ and $\angle B \cong \angle E$. Which of the following additional conditions is NOT sufficient by itself to guarantee that $\Delta ABC \cong \Delta DEF$?

a. $AC \cong DF$ ✓

b. $BC \cong EF$

c. $\angle A \cong \angle D$

d. $\angle C \cong \angle F$

We are given one pair of congruent sides ($AB \cong DE$) and one pair of congruent angles ($\angle B \cong \angle E$). We evaluate each choice: 1. If $AC \cong DF$: The given angle $\angle B$ is not between the two known sides AB and AC. This setup corresponds to Side-Side-Angle (SSA), which is not a valid congruence criterion because two non-congruent triangles can share these measurements (the ambiguous case). Thus, this condition is NOT sufficient. 2. If $BC \cong EF$: The angle $\angle B$ is the included an.

© ClipSAT — clipsat.org | All Rights Reserved

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)**

Can segments of length 10, 12, and 12 form a triangle?

a. Only if the triangle is isosceles

b. No, they cannot form a triangle

c. **Yes, they can form a triangle ✓**

d. Only if the triangle is right

Triangle Inequality Theorem: the sum of any two sides must exceed the third. $10 + 12 = 22 > 12$, $10 + 12 = 22 > 12$, $12 + 12 = 24 > 10$, so yes.

30 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)**

Two triangles have two pairs of corresponding angles and a non-included side are congruent. Which congruence postulate proves the triangles congruent? (variant 5)

a. ASA

b. SAS

c. SSS

d. **AAS ✓**

When two pairs of corresponding angles and a non-included side are congruent, the triangles are congruent by AAS.

31 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)**

Two triangles have two pairs of corresponding angles and the included side are congruent. Which congruence postulate proves the triangles congruent? (variant 14)

a. SSS

b. **ASA ✓**

c. SAS

d. AAS

When two pairs of corresponding angles and the included side are congruent, the triangles are congruent by ASA.

© ClipSAT — clipsat.org | All Rights Reserved

32 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

Two triangles have two pairs of corresponding sides and the included angle are congruent. Which congruence postulate proves the triangles congruent? (variant 2)

a. AAS

b. ASA

c. SSS

d. **SAS ✓**

When two pairs of corresponding sides and the included angle are congruent, the triangles are congruent by SAS.

33 [Intermediate] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

Two triangles have all three pairs of corresponding sides are congruent. Which congruence postulate proves the triangles congruent? (variant 2)

a. AAS

b. SAS

c. **SSS ✓**

d. ASA

When all three pairs of corresponding sides are congruent, the triangles are congruent by SSS.

34 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

Triangles $ABC \cong DEF$ by SAS. If $\angle A = (3x + 7)^\circ$ and the corresponding $\angle D = (1x + 13)^\circ$, find $\angle A$.

a. **16° ✓**

b. 24°

c. 3°

d. 164°

By CPCTC, $\angle A = \angle D$, so $3x + 7 = 1x + 13$. Solving gives $x = 3$, so $\angle A = 16^\circ$.

35 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

Triangle with $\angle A = 30^\circ$, $\angle B = 90^\circ$, side $a = 5$ (opposite A). Find side b (Law of Sines).

a. **10 ✓**

b. 5

c. 12

d. $\frac{5}{2}$

$\frac{a}{\sin A} = \frac{b}{\sin B} \Rightarrow b = \frac{5 \sin 90^\circ}{\sin 30^\circ} = 10.$

© ClipSAT — clipsat.org | All Rights Reserved

36 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

A 30-60-90 right triangle has a short leg of length 21. What is the length of the longer leg?

a. $21\sqrt{3}$ ✓

b. 21

c. 42

d. $21\sqrt{2}$

In a 30-60-90 triangle, longer leg $=$ short leg $\times \sqrt{3} = 21\sqrt{3}$.

37 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

From a point 30 m from the base of a tower, the angle of elevation to the top is 60°. Find the height of the tower, rounded to two decimal places.

a. 54.96

b. 30

c. **51.96** ✓

d. 25.98

height $=$ distance $\times \tan(60^\circ) = 30 \times 1.7321 = 51.96$ m.

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 10.01 Coordinate Geometry

A 30-60-90 right triangle has a short leg of length 18. What is the length of the hypotenuse?

a. 18

b. $18\sqrt{2}$

c. $18\sqrt{3}$

d. **36** ✓

In a 30-60-90 triangle, hypotenuse $= 2 \times$ short leg $= 2 \times 18 = 36$.

39 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 11.01 Probability

A 45-45-90 right triangle has legs of length 40. What is the length of the hypotenuse?

a. $40\sqrt{3}$

b. $40\sqrt{2}$ ✓

c. 40

d. 80

In a 45-45-90 triangle, hypotenuse $=$ leg $\times \sqrt{2} = 40\sqrt{2}$.

© ClipSAT — clipsat.org | All Rights Reserved

40 [Challenging] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14) • Cross-Lesson: 2.01 Lines & Angles

In a right triangle, one acute angle measures 45° and the side adjacent to it is 59. Find the length of the side opposite the 45° angle, rounded to two decimal places.

a. 60.5

b. 57.0

c. **59.0** ✓

d. 70.8

opposite = adjacent × tan(45°) = 59 × 1.0 = 59.0.

41 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 4.01 Similarity & Proportional Reasoning

In a right triangle, one acute angle measures 30° and the hypotenuse is 49. Find the length of the side adjacent to the 30° angle, rounded to two decimal places.

a. 40.43

b. 43.93

c. **42.43** ✓

d. 50.92

adjacent = hyp × cos(30°) = 49 × 0.866 = 42.43.

42 [Challenging] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28) • Cross-Lesson: 5.01 Right Triangles & Trigonometry

A right triangle has legs of length 22 and 29. What is the length of the hypotenuse? (Simplify if possible.)

a. $5\sqrt{53}$ ✓

b. $\sqrt{1321}$

c. $\sqrt{1329}$

d. 51

By the Pythagorean Theorem, $c = \sqrt{22^2 + 29^2} = \sqrt{1325} = 5\sqrt{53}$.

43 [Challenging] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37) • Cross-Lesson: 6.01 Polygons & Quadrilaterals

In a right triangle, one acute angle measures 30° and the hypotenuse is 40. Find the length of the side opposite the 30° angle, rounded to two decimal places.

a. 24.0

b. 18.0

c. 21.5

d. **20.0** ✓

opposite = hyp × sin(30°) = 40 × 0.5 = 20.0.

© ClipSAT — clipsat.org | All Rights Reserved

44 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 7.01 Circles

A right triangle has a hypotenuse of length 388 and one leg of length 260. What is the length of the other leg?

a. 388

b. **288** ✓

c. 260

d. 128

By the Pythagorean Theorem, $b = \sqrt{388^2 - 260^2} = \sqrt{82944} = 288$.

45 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 8.01 Area, Surface Area & Volume

Two coastal observation stations, A and B, are situated 12 km apart along a straight shoreline running due east-west, with station B located due east of station A. Both stations detect a distressed vessel at point C offshore (north of the shoreline). From station A, the vessel is on a bearing of 030°. From station B, the vessel is on a bearing of 315°. What is the area of ΔABC, in square kilometers? *(Note: Bearings are measured clockwise from true north.)*

a. $108 - 36\sqrt{3}$ ✓

b. $108 + 36\sqrt{3}$

c. $36\sqrt{3} - 36$

d. $216 - 72\sqrt{3}$

1. **Determine the interior angles of ΔABC**: - The baseline segment AB runs due east from A to B, corresponding to a bearing of 090°. - The bearing from A to C is 030°. The interior angle at A is: $\angle CAB = 90^\circ - 30^\circ = 60^\circ$ - From station B, due west toward A is along a bearing of 270°. The vessel C is on a bearing of 315°. Thus, the interior angle at B is: $\angle CBA = 315^\circ - 270^\circ = 45^\circ$ - The interior angle at C is.

© ClipSAT — clipsat.org | All Rights Reserved