Statistics & Probability

ClipSAT Printable Practice Suite (clipsat.org) • Topic 4.01

ANSWER KEY — COMPLETE WORKED SOLUTIONS

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

For the model $f(x) = 5x + 4$ in exponential equations, evaluate $f(3)$.

Final Answer: 19. Substitute $x = 3$: $f(3) = 5(3) + 4 = 19$.

2 [Foundational] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

[Statistics & Probability · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

3 [Foundational] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

[AI SL — Specific Topics · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

4 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

[Overview · Practice 2] If the governing relation for this system is $g(t) = 4t + 4$, evaluate $g(6)$ and find the inverse input t when $g(t) = 32$.

Substitute $t = 6$: $g(6) = 4(6) + 4 = \mathbf{28}$. Setting $4t + 4 = 32$ gives $4t = 28 \Rightarrow t = \mathbf{7}$.

5 [Foundational] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

[Number & Algebra · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

6 [Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

[Functions · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

© ClipSAT — clipsat.org | All Rights Reserved

7 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

[Calculus · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

8 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

A bag contains 5 red and 4 green marbles. Three marbles are drawn at random without replacement. Find the probability that: (a) all three are red, (b) at least two are red.

Total marbles = 9. Number of ways to choose 3 marbles = $\binom{9}{3} = \frac{9 \times 8 \times 7}{3 \times 2 \times 1} = 84$. (a) All 3 are red: $\binom{5}{3} = 10$. Probability: $P(\text{all red}) = \frac{10}{84} = \frac{5}{42} \approx 0.119$ (b) 'At least two red' means exactly 2 red and 1 green, or exactly 3 red: Ways for 2 red, 1 green: $\binom{5}{2} \times \binom{4}{1} = 10 \times 4 = 40$. Ways for 3 red: $\binom{5}{3} = 10$. Total favorable outcomes: $40 + 10 = 50$. Probability.

9 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

Events A and B are such that $P(A) = 0.6$, $P(B) = 0.5$, and $P(A \cup B) = 0.8$. Find $P(A \cap B)$, calculate $P(A|B)$, and determine whether A and B are independent.

1. Use the addition rule: $P(A \cup B) = P(A) + P(B) - P(A \cap B) \Rightarrow 0.8 = 0.6 + 0.5 - P(A \cap B) \Rightarrow P(A \cap B) = 1.1 - 0.8 = 0.3$. 2. Conditional probability: $P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.3}{0.5} = 0.6$ 3. Check for independence: $P(A \cap B) = 0.3$ and $P(A) \times P(B) = 0.6 \times 0.5 = 0.30$. Since $P(A \cap B) = P(A)P(B)$ (or equivalently $P(A|B) = P(A) = 0.6$), events A and B are independent.

10 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

The table shows the time, t minutes, taken by 50 students to complete a homework task. Intervals: $0 < t \leq 10$ (freq: 8), $10 < t \leq 20$ (freq: 14), $20 < t \leq 30$ (freq: 18), $30 < t \leq 40$ (freq: 10). Calculate an estimate of the mean time taken.

Find class midpoints (x) and products (fx):Midpoint 5: $8 \times 5 = 40$Midpoint 15: $14 \times 15 = 210$Midpoint 25: $18 \times 25 = 450$Midpoint 35: $10 \times 35 = 350$ $\sum fx = 40 + 210 + 450 + 350 = 1050$ Estimated Mean $= \frac{\sum fx}{\sum f} = \frac{1050}{50} = 21$ minutes

11 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A continuous random variable X has probability density function $f(x) = 2e^{-2x}$ for $x \geq 0$ and 0 elsewhere. Find the cumulative distribution function $F(x)$ and determine the exact median of X.

1. Find CDF for $x \geq 0$: $F(x) = \int_0^x 2e^{-2t}\,dt = [-e^{-2t}]_0^x = 1 - e^{-2x}$. 2. Set $F(m) = 0.5$ to find the median m: $1 - e^{-2m} = 0.5 \Rightarrow e^{-2m} = 0.5 \Rightarrow -2m = \ln(0.5) = -\ln 2 \Rightarrow m = \frac{\ln 2}{2}$.

© ClipSAT — clipsat.org | All Rights Reserved

12 **[Intermediate]** • **Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)**

Using the same bag (5 red, 3 blue), find the probability of drawing exactly one red ball.

$P(\text{RB}) + P(\text{BR}) = \frac{5}{8} \cdot \frac{3}{7} + \frac{3}{8} \cdot \frac{5}{7} = \frac{15}{56} + \frac{15}{56} = \frac{30}{56} = \frac{15}{28}$.

13 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)**

[Probability & Statistics · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Substitute $t = 8$: $g(8) = 5(8) + 5 = \mathbf{45}$. Setting $5t + 5 = 50$ gives $5t = 45 \Rightarrow t = \mathbf{9}$.

14 **[Intermediate]** • **Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)**

[Statistics & Probability (HL) · Practice 2] If the governing relation for this system is $g(t) = 4t + 4$, evaluate $g(6)$ and find the inverse input t when $g(t) = 32$.

Substitute $t = 6$: $g(6) = 4(6) + 4 = \mathbf{28}$. Setting $4t + 4 = 32$ gives $4t = 28 \Rightarrow t = \mathbf{7}$.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 **[Challenging]** • **Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)** • **Cross-Lesson: 3.01 Geometry & Trigonometry**

A continuous random variable X has probability density function $f(x) = kx(2 - x)$ for $0 \le x \le 2$ and $f(x) = 0$ elsewhere. Determine: (a) the constant k; (b) the median of X; (c) the variance $\text{Var}(X)$.

1. Normalization condition: $\int_0^2 k(2x - x^2)\,dx = 1 \Rightarrow k\left[x^2 - \frac{x^3}{3}\right]_0^2 = k\left(4 - \frac{8}{3}\right) = k\left(\frac{4}{3}\right) = 1 \Rightarrow k = \frac{3}{4}$. 2. Median m: By visual symmetry of the parabola $f(x) = \frac{3}{4}x(2 - x)$ about $x = 1$, $m = 1$. We verify:
$F(1) = \int_0^1 \frac{3}{4}(2x - x^2)\,dx = \frac{3}{4}\left[x^2 - \frac{x^3}{3}\right]_0^1 = \frac{3}{4}\left(\frac{2}{3}\right) = \frac{1}{2}$. ✓ 3. Variance: By symmetry, $E[X] = 1$..

16 **[Challenging]** • **Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)** • **Cross-Lesson: 5.01 Calculus**

Rotate the Cartesian axes to eliminate the cross-product term in $5x^2 - 4xy + 8y^2 = 36$. Find the angle of rotation θ, standard form in the rotated system (x', y'), and the lengths of the semi-major and semi-minor axes.

Step 1: Determine the rotation angle θ: Here $A = 5$, $B = -4$, and $C = 8$.
$\cot(2\theta) = \frac{A - C}{B} = \frac{5 - 8}{-4} = \frac{-3}{-4} = \frac{3}{4} > 0$. Since $\cot(2\theta) = 3/4$, by right triangle trigonometry $\cos(2\theta) = \frac{3}{5}$.
Using half-angle formulas for $\theta \in (0, \pi/4)$: $\cos\theta = \sqrt{\frac{1 + \cos(2\theta)}{2}} = \sqrt{\frac{1 + 3/5}{2}} = \sqrt{\frac{4}{5}} = \frac{2}{\sqrt{5}}$, .

© ClipSAT — clipsat.org | All Rights Reserved

17 **[Challenging]** • **Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)** • **Cross-Lesson: 6.01 AI SL — Specific Topics**

(IB AA/AI HL) Given the initial value problem $\frac{dy}{dx} = x + y$ with $y(0) = 1$, use Euler's numerical method with step size $h = 0.1$ to approximate $y(0.2)$. Compare your numerical result with the exact analytical solution.

1. Step formula: $x_{n+1} = x_n + h$, $y_{n+1} = y_n + h(x_n + y_n)$ with $h = 0.1$, $x_0 = 0$, $y_0 = 1$. Step 1 (to $x_1 = 0.1$): $y_1 = y_0 + 0.1(x_0 + y_0) = 1 + 0.1(0 + 1) = 1 + 0.1 = 1.1$. Step 2 (to $x_2 = 0.2$): $y_2 = y_1 + 0.1(x_1 + y_1) = 1.1 + 0.1(0.1 + 1.1) = 1.1 + 0.1(1.2) = 1.1 + 0.12 = 1.22$. Euler's approximation is $y(0.2) \approx 1.22$. 2. Exact analytical solution: Rewrite as $\frac{dy}{dx} - y = x$. Integrating factor $\mu = e^{-x}$.

18 **[Challenging]** • **Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)** • **Cross-Lesson: 1.01 Number & Algebra**

(IB AI HL) A market transition matrix $T = (0.7, 0.2; 0.3, 0.8)$ represents weekly customer movement between Brand A and Brand B. Find: (a) the probability that a customer initially with Brand A is with Brand B after 2 weeks; (b) the long-term steady-state market distribution.

(a) State distribution after 2 weeks is $\mathbf{p}_2 = T^2\mathbf{p}_0$. With $\mathbf{p}_0 = (1; 0)$: $T^2 = (0.7, 0.2; 0.3, 0.8)(0.7, 0.2; 0.3, 0.8) = (0.49 + 0.06, 0.14 + 0.16; 0.21 + 0.24, 0.06 + 0.64) = (0.55, 0.30; 0.45, 0.70)$. Thus $\mathbf{p}_2 = (0.55; 0.45)$, so the probability of being with Brand B after 2 weeks is 0.45 (45%). (b) Steady-state vector $\mathbf{s} = (s_1; s_2)$ satisfies $(T - I)\mathbf{s} = \mathbf{0}$.

19 **[Challenging]** • **Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)** • **Cross-Lesson: 2.01 Functions**

Let $\mathbf{a} = (2; -1; 3)$ and $\mathbf{b} = (1; 4; -2)$. Find the cross product $\mathbf{a} \times \mathbf{b}$, and determine the exact area of the triangle with adjacent sides $\mathbf{a}$ and $\mathbf{b}$.

1. Compute the cross product via determinant: $\mathbf{a} \times \mathbf{b} = |\mathbf{i}, \mathbf{j}, \mathbf{k}; 2, -1, 3; 1, 4, -2| = \mathbf{i}[(-1)(-2) - 3(4)] - \mathbf{j}[2(-2) - 3(1)] + \mathbf{k}[2(4) - (-1)(1)]$ $= \mathbf{i}(2 - 12) - \mathbf{j}(-4 - 3) + \mathbf{k}(8 + 1) = -10\mathbf{i} + 7\mathbf{j} + 9\mathbf{k} = (-10; 7; 9)$. 2. The area of the triangle is half the magnitude of the cross product.

20 **[Challenging]** • **Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)** • **Cross-Lesson: 3.01 Geometry & Trigonometry**

Two independent random variables are normally distributed as $X \sim N(50, 16)$ and $Y \sim N(45, 9)$. Find the probability that $X > Y$.

1. Define the difference variable $W = X - Y$. The condition $X > Y$ is equivalent to $W > 0$. 2. Since X and Y are independent normals, W is normally distributed: Mean: $\mu_W = E[X] - E[Y] = 50 - 45 = 5$. Variance: $\sigma_W^2 = \text{Var}(X) + \text{Var}(Y) = 16 + 9 = 25 \Rightarrow \sigma_W = 5$. Thus $W \sim N(5, 25)$. 3. Compute $P(W > 0)$: $z = \frac{0 - 5}{5} = -1.0$. $P(W > 0) = P(Z > -1.0) = P(Z < 1.0) = \Phi(1.0) \approx 0.8413 \approx 0.841$.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

© ClipSAT — clipsat.org | All Rights Reserved

21 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

For the model $f(x) = 5x + 3$ in logic, evaluate $f(2)$.

a. 13 ✓

b. 18

c. 10

d. 17

Substitute $x = 2$: $f(2) = 5(2) + 3 = 13$.

22 [Foundational] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

For the model $f(x) = 6x + 4$ in proof by deduction, evaluate $f(2)$.

a. 16 ✓

b. 22

c. 12

d. 20

Substitute $x = 2$: $f(2) = 6(2) + 4 = 16$.

23 [Foundational] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A data set recorded for logic consists of the four values $\{11, 15, 19, 25\}$. What is the arithmetic mean?

a. 15.5

b. 16.5

c. 17.5 ✓

d. 19.5

The sum is $11 + 15 + 19 + 25 = 70$, and dividing by 4 gives 17.5.

24 [Foundational] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A data set recorded for proof by deduction consists of the four values $\{12, 16, 20, 26\}$. What is the arithmetic mean?

a. 16.5

b. 17.5

c. 18.5 ✓

d. 20.5

The sum is $12 + 16 + 20 + 26 = 74$, and dividing by 4 gives 18.5.

© ClipSAT — clipsat.org | All Rights Reserved

25 [Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

A data set recorded for exponential equations consists of the four values $\{15, 19, 23, 29\}$. What is the arithmetic mean?

a. 19.5

b. 20.5

c. 21.5 ✓

d. 23.5

The sum is $15 + 19 + 23 + 29 = 86$, and dividing by 4 gives 21.5.

26 [Foundational] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

The box plot below shows the distribution of scores for a class of 30 students on a recent test. The interquartile range (IQR) is 22 marks. The median score is 75 marks. The minimum score is 30 marks, and the maximum score is 98 marks. Let the first quartile be denoted by Q_1 and the third quartile be denoted by Q_3. If the upper whisker extends to 98 marks, and the third quartile Q_3 is 86 marks, find the value of Q_1 .

a. **64** ✓

b. 75

c. 80

d. 53

The interquartile range (IQR) is defined as the difference between the third quartile (Q_3) and the first quartile (Q_1), i.e., $\text{IQR} = Q_3 - Q_1$. We are given that the IQR is 22 marks. We are also given that the third quartile $Q_3 = 86$ marks. Substituting these values into the formula, we get: $22 = 86 - Q_1$ To find Q_1, we rearrange the equation: $Q_1 = 86 - 22$ $Q_1 = 64$ Therefore, the first quartile is 64 marks. Common errors might include misinterpreting the median as Q_1 or Q_3, or inco.

27 [Foundational] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

The histogram below shows the distribution of scores for a class of students on a recent quiz. The total number of students who took the quiz is 50. The frequency for the score interval [70, 80) is x. Determine the value of x.

a. 5

b. **10** ✓

c. 15

d. 20

The histogram represents the frequency distribution of quiz scores. The total number of students is given as 50. The frequencies for the score intervals are as follows: [50, 60): 8 [60, 70): 12 [70, 80): x [80, 90): 7 [90, 100]: 8 The sum of the frequencies must equal the total number of students: 8 + 12 + x + 7 + 8 = 50 Combine the known frequencies: 35 + x = 50 Solve for x: x = 50 - 35 x = 15 Therefore, the frequency for the score interval [70, 80) is 15.

© ClipSAT — clipsat.org | All Rights Reserved

28 [Foundational] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

The bar chart below shows the number of students who chose each of four different sports as their favorite. What is the total number of students surveyed?

a. 50

b. 60

c. **70** ✓

d. 80

The bar chart shows the following counts for each sport: Sport 1: 14 students Sport 2: 22 students Sport 3: 18 students Sport 4: 26 students To find the total number of students surveyed, we sum the number of students for each sport: $14 + 22 + 18 + 26$. $14 + 22 = 36$ $18 + 26 = 44$ $36 + 44 = 80$. Therefore, the total number of students surveyed is 80.

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

The following bar chart shows the number of students who achieved different grades in a mathematics test. The grades are A, B, C, and D. The data is as follows: Grade A: 14 students, Grade B: 22 students, Grade C: 18 students, Grade D: 26 students. What is the total number of students who took the test?

a. 70

b. **80** ✓

c. 78

d. 60

To find the total number of students who took the test, we need to sum the number of students who achieved each grade. Number of students with Grade A = 14 Number of students with Grade B = 22 Number of students with Grade C = 18 Number of students with Grade D = 26 Total number of students = 14 + 22 + 18 + 26 = 80. Therefore, 80 students took the test.

30 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

The bar chart below shows the number of students who achieved a certain score on a test. What is the total number of students who scored 6 or 8?

a. 30

b. **28** ✓

c. 26

d. 32

From the bar chart, the number of students who scored 6 is 18 and the number of students who scored 8 is 12. The total number of students who scored 6 or 8 is the sum of these two values: $18 + 12 = 30$.

© ClipSAT — clipsat.org | All Rights Reserved

31 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

Find the standard deviation of the data set: $12, 19, 14, 22, 24$, correct to 2 decimal places.

a. 4.98

b. **4.58** ✓

c. 4.18

d. 3.78

Working through this statistics step: Mean $= \frac{12+19+14+22+24}{5} = 18.20$. $\sigma = \sqrt{\frac{\sum(x-\bar{x})^2}{n}} \approx 4.58$.

32 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

The following two-way table shows the results of a survey on favourite sports among a group of students. Calculate the total number of students surveyed.

a. 50

b. 45

c. **60** ✓

d. 55

The total number of students surveyed is the sum of all entries in the two-way table.

Total $= 15 + 10 + 20 + 15 = 60$.

33 [Intermediate] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7)

A bag contains 8 red and 9 blue balls. Two balls are drawn at random without replacement. Find $P(\text{both red})$.

a. $\frac{64}{289}$

b. $\frac{8}{17}$

c. $\frac{7}{34}$ ✓

d. $\frac{56}{289}$

Working through this statistics step: $P(\text{both red}) = \frac{8}{17} \times \frac{7}{16} = \frac{7}{34}$.

34 [Intermediate] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

A bag contains 7 red and 8 blue balls. A ball is drawn, replaced, then a second ball is drawn. Find $P(\text{both red})$.

a. $\frac{49}{225}$ ✓

b. $\frac{112}{225}$

c. $\frac{8}{15}$

d. $\frac{7}{15}$

Working through this statistics step: $P(\text{both red}) = \left(\frac{7}{15}\right)^2 = \frac{49}{225}$.

© ClipSAT — clipsat.org | All Rights Reserved

35 [Intermediate] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

Find the median of the data set: $8, 30, 32, 34, 48, 53, 54$.

a. **34** ✓

b. 33

c. 32

d. 48

Working through this statistics step: Sorted data: $8, 30, 32, 34, 48, 53, 54$. The middle value is 34.

36 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

A bag contains 4 blue balls, 8 yellow balls, and 3 green balls. A ball is drawn at random. Find $P(\text{yellow})$.

a. $\frac{8}{15}$ ✓

b. $\frac{1}{5}$

c. $\frac{7}{15}$

d. $\frac{4}{15}$

Working through this statistics step: $P(\text{yellow}) = \frac{8}{15} = \frac{8}{15}$.

37 [Intermediate] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

A bag contains 3 red balls, 5 yellow balls, and 8 blue balls. A ball is drawn at random. Find $P(\text{yellow})$.

a. $\frac{11}{16}$

b. $\frac{1}{2}$

c. $\frac{5}{16}$ ✓

d. $\frac{3}{16}$

Working through this statistics step: $P(\text{yellow}) = \frac{5}{16} = \frac{5}{16}$.

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 1.01 Number & Algebra

A bag contains 4 red balls, 3 green balls, and 4 yellow balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{3}{11}$ ✓

b. $\frac{2}{11}$

c. $\frac{4}{11}$

d. $\frac{8}{11}$

Working through this statistics step: $P(\text{green}) = \frac{3}{11} = \frac{3}{11}$.

© ClipSAT — clipsat.org | All Rights Reserved

39 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 2.01 Functions

A bag contains 3 green balls, 7 red balls, and 4 yellow balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{2}{7}$

b. $\frac{1}{2}$

c. $\frac{11}{14}$

d. $\frac{3}{14}$ ✓

Working through this statistics step: $P(\text{green}) = \frac{3}{14} = \frac{3}{14}$.

40 [Challenging] • Target: Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14) • Cross-Lesson: 3.01 Geometry & Trigonometry

A bag contains 4 yellow balls, 6 blue balls, and 3 green balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{6}{13}$

b. $\frac{3}{13}$ ✓

c. $\frac{4}{13}$

d. $\frac{10}{13}$

Working through this statistics step: $P(\text{green}) = \frac{3}{13} = \frac{3}{13}$.

41 [Challenging] • Target: Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20) • Cross-Lesson: 5.01 Calculus

A bag contains 8 yellow balls, 8 red balls, and 4 blue balls. A ball is drawn at random. Find $P(\text{yellow})$.

a. $\frac{9}{20}$

b. $\frac{2}{5}$ ✓

c. $\frac{3}{5}$

d. $\frac{1}{5}$

Working through this statistics step: $P(\text{yellow}) = \frac{8}{20} = \frac{2}{5}$.

42 [Challenging] • Target: Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28) • Cross-Lesson: 6.01 AI SL — Specific Topics

A bag contains 6 yellow balls, 2 green balls, and 3 blue balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{9}{11}$

b. $\frac{6}{11}$

c. $\frac{3}{11}$

d. $\frac{2}{11}$ ✓

Working through this statistics step: $P(\text{green}) = \frac{2}{11} = \frac{2}{11}$.

© ClipSAT — clipsat.org | All Rights Reserved

43 [Challenging] • Target: Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37) • Cross-Lesson: 1.01 Number & Algebra

A bag contains 3 blue balls, 7 green balls, and 2 red balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{1}{4}$

b. $\frac{7}{12}$ ✓

c. $\frac{1}{6}$

d. $\frac{5}{12}$

Working through this statistics step: $P(\text{green}) = \frac{7}{12} = \frac{7}{12}$.

44 [Challenging] • Target: Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45) • Cross-Lesson: 2.01 Functions

A bag contains 6 yellow balls, 4 green balls, and 3 red balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{3}{13}$

b. $\frac{9}{13}$

c. $\frac{4}{13}$ ✓

d. $\frac{6}{13}$

Working through this statistics step: $P(\text{green}) = \frac{4}{13} = \frac{4}{13}$.

45 [Challenging] • Target: Free-Response Questions (FRQs): Foundational Fluency (1–7) • Cross-Lesson: 3.01 Geometry & Trigonometry

A bag contains 8 green balls, 7 yellow balls, and 6 blue balls. A ball is drawn at random. Find $P(\text{green})$.

a. $\frac{13}{21}$

b. $\frac{2}{7}$

c. $\frac{1}{3}$

d. $\frac{8}{21}$ ✓

Working through this statistics step: $P(\text{green}) = \frac{8}{21} = \frac{8}{21}$.

© ClipSAT — clipsat.org | All Rights Reserved