Radian Measure and Arc Length

ClipSAT Printable Practice Suite (clipsat.org) • Topic 2.02

ANSWER KEY — COMPLETE WORKED SOLUTIONS

Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 **[Foundational]** • **Target: Converting between degrees and radians**

Convert each angle to radians, in exact form:

a $45° = \frac{\pi}{4}$ **b** $120° = \frac{2\pi}{3}$ **c** $270° = \frac{3\pi}{2}$ **d** $330° = \frac{11\pi}{6}$

(a) $= \frac{\pi}{4}$; (b) $= \frac{2\pi}{3}$; (c) $= \frac{3\pi}{2}$; (d) $= \frac{11\pi}{6}$

2 **[Foundational]** • **Target: Converting between degrees and radians**

Convert each angle to degrees. Round to one decimal place where needed:

a $\frac{\pi}{6} = 30°$ **b** $\frac{5\pi}{4} = 225°$ **c** $\frac{7\pi}{3} = 420°$ **d** 3.5 rad $\approx 200.5°$

(a) $= 30°$; (b) $= 225°$; (c) $= 420°$; (d) $\approx 200.5°$

3 **[Foundational]** • **Target: Arc length and sector area**

A circle has radius 6 cm. Find the exact length of the arc subtended by a central angle of $\frac{2\pi}{3}$.

$s = r\theta = 6 \cdot \frac{2\pi}{3} = 4\pi \approx 12.57$ cm.

4 **[Foundational]** • **Target: Arc length and sector area**

A sector of a circle of radius 10 cm has central angle $\frac{\pi}{5}$. Find its exact area.

$A = \frac{1}{2}r^2\theta = \frac{1}{2}(100)\frac{\pi}{5} = 10\pi \approx 31.42$ cm^2.

5 **[Foundational]** • **Target: Arc length and sector area**

A wheel of radius 30 cm rolls along the ground without slipping.

a Through what angle, in radians, does the wheel turn when it travels 1.5 m? $\theta = \frac{s}{r} = \frac{150}{30} = 5$ radians.

b Express this angle to the nearest tenth of a degree. $5 \times \frac{180°}{\pi} \approx 286.5°$.

(a) $\theta = \frac{s}{r} = \frac{150}{30} = 5$ radians.; (b) $5 \times \frac{180°}{\pi} \approx 286.5°$.

6 **[Foundational]** • **Target: Arc length and sector area**

A pendulum of length 80 cm swings through an angle of $25°$. Find the length of the arc traced by its tip, to the nearest tenth of a centimetre.

$\theta = 25° = \frac{25\pi}{180}$ rad, so $s = 80 \cdot \frac{25\pi}{180} \approx 34.9$ cm.

© ClipSAT — clipsat.org | All Rights Reserved

7 [Foundational] • Target: Angular and linear speed

A carousel horse is 4 m from the center and completes one full rotation every 20 seconds.

a Find the angular speed of the horse, in radians per second. $\omega = \frac{2\pi}{20} = \frac{\pi}{10}$ rad/s.

b Find the linear speed of the horse, in m/s. $v = r\omega = 4 \cdot \frac{\pi}{10} = \frac{2\pi}{5} \approx 1.26$ m/s.

(a) $\omega = \frac{2\pi}{20} = \frac{\pi}{10}$ rad/s.; (b) $v = r\omega = 4 \cdot \frac{\pi}{10} = \frac{2\pi}{5} \approx 1.26$ m/s.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

8 [Intermediate] • Target: Angular and linear speed

A car's wheel has radius 35 cm and spins at 400 revolutions per minute. Find the car's speed in km/h, to one decimal place.

$v = r\omega$, with $\omega = 400 \times 2\pi$ rad/min $= 800\pi$ rad/min. $v = 0.35 \times 800\pi \approx 879.6$ m/min $= 52{,}776$ m/h ≈ 52.8 km/h.

9 [Intermediate] • Target: Angular and linear speed

Two gears mesh together: the smaller gear has radius 5 cm and spins at 300 rpm. If the larger gear has radius 15 cm, find its rotational speed in rpm (the linear speed at the point of contact is the same for both gears).

$v = r_1\omega_1 = r_2\omega_2$: $5(300) = 15\,\omega_2$, so $\omega_2 = 100$ rpm.

10 [Intermediate] • Target: Coterminal angles and reference angles

Find a positive coterminal angle (between 0 and 2π) for each:

a $\frac{9\pi}{4} = \frac{\pi}{4}$ **b** $-\frac{\pi}{3} = \frac{5\pi}{3}$ **c** $\frac{13\pi}{6} = \frac{\pi}{6}$

(a) $= \frac{\pi}{4}$; (b) $= \frac{5\pi}{3}$; (c) $= \frac{\pi}{6}$

11 [Intermediate] • Target: Coterminal angles and reference angles

Find the reference angle for each angle:

a $\frac{5\pi}{6} = \frac{\pi}{6}$ **b** $\frac{7\pi}{4} = \frac{\pi}{4}$ **c** $\frac{4\pi}{3} = \frac{\pi}{3}$

(a) $= \frac{\pi}{6}$; (b) $= \frac{\pi}{4}$; (c) $= \frac{\pi}{3}$

© ClipSAT — clipsat.org | All Rights Reserved

12 [Intermediate] • Target: The radian definition itself

State the definition of one radian in terms of arc length and radius, and explain why radians are considered a 'natural' unit for angles.

One radian is the angle subtended at the center of a circle by an arc whose length equals the radius. Radians are natural because the arc length formula $s = r\theta$ has no extra conversion constant when θ is in radians, unlike degrees.

13 [Intermediate] • Target: The radian definition itself

Explain why the full circle corresponds to 2π radians, using the circumference formula $C = 2\pi r$.

Since $s = r\theta$ and the full circumference is $s = 2\pi r$, setting $r\theta = 2\pi r$ gives $\theta = 2\pi$ — the full circle sweeps exactly 2π radians.

14 [Intermediate] • Target: Combined arc length and sector area problems

A sprinkler sprays water over a sector of radius 8 m and central angle $\frac{\pi}{3}$. Find both the area watered and the length of the outer arc of the watered region.

Area: $A = \frac{1}{2}(8)^2\left(\frac{\pi}{3}\right) = \frac{32\pi}{3} \approx 33.5$ m^2. Arc length: $s = 8\left(\frac{\pi}{3}\right) = \frac{8\pi}{3} \approx 8.4$ m.

Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 [Challenging] • Target: Combined arc length and sector area problems • Cross-Lesson: 7.01 Exponential Functions and Their Graphs

A slice of pizza (a sector) has radius 12 cm and arc length 10 cm along its crust. Find the central angle in radians and the area of the slice.

$\theta = \frac{s}{r} = \frac{10}{12} = \frac{5}{6}$ rad. Area $= \frac{1}{2}r^2\theta = \frac{1}{2}(144)\left(\frac{5}{6}\right) = 60$ cm^2.

16 [Challenging] • Target: Common angles in both units (reference table) • Cross-Lesson: 7.02 Logarithms and Their Properties

Complete a conversion table for the common angles $0°, 30°, 45°, 60°, 90°, 180°, 360°$ in radians.

$0° = 0$; $30° = \frac{\pi}{6}$; $45° = \frac{\pi}{4}$; $60° = \frac{\pi}{3}$; $90° = \frac{\pi}{2}$; $180° = \pi$; $360° = 2\pi$.

17 [Challenging] • Target: Satellite and orbital speed applications • Cross-Lesson: 7.03 Exponential and Logarithmic Equations

A satellite orbits Earth at a radius of 7000 km, completing one orbit every 2 hours. Find its angular speed in rad/h and its linear speed in km/h, to the nearest whole number.

$\omega = \frac{2\pi}{2} = \pi$ rad/h. $v = r\omega = 7000\pi \approx 21{,}991$ km/h.

© ClipSAT — clipsat.org | All Rights Reserved

18 [Challenging] • Target: Satellite and orbital speed applications • Cross-Lesson: 7.04 Applications: Growth and Decay

Earth rotates once every 24 hours. Find the linear speed (due to rotation alone) of a point on the equator, using Earth's radius ≈ 6371 km, to the nearest km/h.

$\omega = \frac{2\pi}{24} = \frac{\pi}{12}$ rad/h. $v = 6371 \cdot \frac{\pi}{12} \approx 1668$ km/h.

19 [Challenging] • Target: Negative angles and full-turn arcs • Cross-Lesson: 8.01 Rational Functions and Asymptotes

Find the exact arc length swept by a radius-4 circle for a rotation of $-\frac{3\pi}{2}$ radians (clockwise), and interpret the sign.

Arc length uses the magnitude of θ: $s = 4\left(\frac{3\pi}{2}\right) = 6\pi$. The negative sign in the angle indicates clockwise rotation, but arc length itself is always a positive (unsigned) distance.

20 [Challenging] • Target: Arc length and sector area • Cross-Lesson: 8.02 Graphing Rational Functions

In a circle of radius 10 cm, chord AB subtends an angle of $\theta = \frac{2\pi}{3}$ radians at the centre O. Find: (a) the exact length of chord AB, (b) the exact area of the minor segment bounded by chord AB and minor arc AB, in terms of π and $\sqrt{3}$.

(a) Use the cosine rule in triangle OAB with $OA = OB = 10$ and $\angle AOB = \frac{2\pi}{3} = 120^\circ$: $AB^2 = 10^2 + 10^2 - 2(10)(10)\cos\left(\frac{2\pi}{3}\right) = 100 + 100 - 200\left(-\frac{1}{2}\right) = 200 + 100 = 300$ Thus $AB = \sqrt{300} = 10\sqrt{3}$ cm. (b) Area of sector OAB: $A_{\text{sector}} = \frac{1}{2}r^2\theta = \frac{1}{2}(10^2)\left(\frac{2\pi}{3}\right) = \frac{100\pi}{3}$ cm^2 Area of triangle OAB.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

21 [Foundational] • Target: Converting between degrees and radians

The diagram shows a sector of a circle with centre O. The arc length of the sector is given by 6π cm and the area of the sector is 27π cm^2. The radius of the circle is r cm and the angle of the sector is θ radians. Find the value of r and θ.

a. **r = 9, $\theta = \frac{2\pi}{3}$** ✓

b. r = 6, $\theta = \pi$

c. r = 3, $\theta = 2\pi$

d. r = 18, $\theta = \frac{\pi}{3}$

The formula for the arc length of a sector is $L = r\theta$, where L is the arc length, r is the radius, and θ is the angle in radians. We are given that $L = 6\pi$. So, we have the equation: $r\theta = 6\pi$ (1) The formula for the area of a sector is $A = \frac{1}{2}r^2\theta$, where A is the area, r is the radius, and θ is the angle in radians. We are given that $A = 27\pi$. So, we have the equation: $\frac{1}{2}r^2\theta = 27\pi$ (2) We can rewrite equation (2) as $r(r\theta) = 54\pi$. Sub.

© ClipSAT — clipsat.org | All Rights Reserved

22 [Foundational] • Target: Arc length and sector area

The diagram shows a sector of a circle with radius r. The angle at the centre is θ radians. If the arc length is 6 cm and the radius is 3 cm, find the angle θ in radians.

a. $\frac{1}{2}$ radians

b. 2 **radians** ✓

c. $\frac{2}{3}$ radians

d. $\frac{3}{2}$ radians

The formula relating arc length (s), radius (r), and angle (θ) in radians is $s = r\theta$. We are given $s = 6$ cm and $r = 3$ cm. Substituting these values into the formula, we get $6 = 3\theta$. To find θ, we divide both sides by 3: $\theta = \frac{6}{3} = 2$ radians. Therefore, the angle at the centre is 2 radians.

23 [Foundational] • Target: Angular and linear speed

A sector has radius 6 cm and angle 0.3 rad. Find the arc length.

a. **9/5** ✓

b. 63/10

c. 14/5

d. 27/5

Applying Circular Measure principles, $s = r\theta = 6 \times 0.3 = 9/5$ cm Therefore, the verified correct option is 9/5.

24 [Foundational] • Target: Coterminal angles and reference angles

Convert $\frac{\pi}{3}$ radians to degrees.

a. 300°

b. 60° ✓

c. 90°

d. 30°

$\frac{\pi}{3} \times \frac{180}{\pi} = 60^\circ$.

© ClipSAT — clipsat.org | All Rights Reserved

25 [Foundational] • Target: The radian definition itself

Consider a particle whose position in the complex plane at time t (in seconds) is given by $z(t) = 3e^{i(\frac{\pi}{6}t + \frac{\pi}{3})}$. (a) Find the initial position of the particle at $t = 0$ in polar coordinates (r, θ), where $r \geq 0$ and $0 \leq \theta < 2\pi$. (b) Determine the angular velocity of the particle in radians per second. (c) If the particle starts at a point A and moves along a circular path, find the arc length traveled by the particle from $t = 0$ to $t = 4$ seconds.

a. (a) $(3, \frac{\pi}{3})$, (b) $\frac{\pi}{6}$ rad/s, (c) 2π

b. (a) $(3, \frac{\pi}{6})$, (b) $\frac{\pi}{3}$ rad/s, (c) $\frac{4\pi}{3}$

c. **(a)** $(3, \frac{2\pi}{3})$**, (b)** $\frac{\pi}{6}$ **rad/s, (c)** $\frac{2\pi}{3}$ ✓

d. (a) $(3, \frac{\pi}{3})$, (b) $\frac{\pi}{3}$ rad/s, (c) $\frac{8\pi}{3}$

The position of the particle is given by $z(t) = 3e^{i(\frac{\pi}{6}t + \frac{\pi}{3})}$. This is in the polar form $z(t) = re^{i\theta(t)}$, where $r = 3$ and $\theta(t) = \frac{\pi}{6}t + \frac{\pi}{3}$. (a) The initial position at $t = 0$ is $z(0) = 3e^{i(\frac{\pi}{6}(0) + \frac{\pi}{3})} = 3e^{i\frac{\pi}{3}}$. In polar coordinates (r, θ), the radius is $r = 3$. The angle is $\theta(0) = \frac{\pi}{3}$. Since $0 \leq \frac{\pi}{3} < 2\pi$, the initial position is $(3, \frac{\pi}{3})$. (b) The angular velocit.

26 [Foundational] • Target: Combined arc length and sector area problems

Find the arc length of a sector with radius 5 and angle 2 radians.

a. 25

b. π

c. 5

d. 10 ✓

Applying Trigonometric Functions principles, Arc length $= r\theta = 5 \cdot 2 = 10$. Therefore, the verified correct option is 10.

27 [Foundational] • Target: Common angles in both units (reference table)

In triangle ABC, angle ABC = θ degrees, and angle ACB = 2θ degrees. The lengths of sides AB and AC are 10 cm and 15 cm respectively. The length of BC is denoted by x cm. Given that the area of triangle ABC is 60 cm^2, find the value of x, to 3 significant figures.

a. 13.7

b. 11.4

c. **12.8** ✓

d. 14.9

Let the angles of triangle ABC be denoted as follows: $\angle ABC = \theta$, $\angle ACB = 2\theta$. By the sine rule, $\frac{AC}{\sin(\theta)} = \frac{AB}{\sin(2\theta)}$. Substituting the given values, $\frac{15}{\sin(\theta)} = \frac{10}{\sin(2\theta)}$. Using the double angle identity, $\sin(2\theta) = 2\sin(\theta)\cos(\theta)$, we get $\frac{15}{\sin(\theta)} = \frac{10}{2\sin(\theta)\cos(\theta)}$. Since $\sin(\theta) eq 0$ for a non-degenerate triangle, we can cancel $\sin(\theta)$ from both sides.

© ClipSAT — clipsat.org | All Rights Reserved

28 [Foundational] • Target: Satellite and orbital speed applications

Let the complex number z_1 be represented in polar form as $z_1 = r_1(\cos\theta_1 + i\sin\theta_1)$, where $r_1 = 4$ and $\theta_1 = \frac{\pi}{3}$. Let another complex number, z_2, be represented as $z_2 = 2(\cos\theta_2 + i\sin\theta_2)$. If $z_1 \cdot z_2 = 8(\cos\alpha + i\sin\alpha)$ and $\alpha = \frac{5\pi}{6}$, find the value of θ_2 in radians, given that $0 \le \theta_2 < 2\pi$.

a. $\frac{\pi}{2}$ ✓

b. $\frac{2\pi}{3}$

c. $\frac{5\pi}{6}$

d. $\frac{\pi}{6}$

We are given two complex numbers in polar form: $z_1 = r_1(\cos\theta_1 + i\sin\theta_1)$ and $z_2 = r_2(\cos\theta_2 + i\sin\theta_2)$. We are provided with $r_1 = 4$ and $\theta_1 = \frac{\pi}{3}$. For z_2, we are given $r_2 = 2$. The product of two complex numbers in polar form is given by $z_1 \cdot z_2 = r_1 r_2(\cos(\theta_1 + \theta_2) + i\sin(\theta_1 + \theta_2))$. We are given that $z_1 \cdot z_2 = 8(\cos\alpha + i\sin\alpha)$, where $\alpha = \frac{5\pi}{6}$. Equating the magnitudes, we.

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 [Intermediate] • Target: Negative angles and full-turn arcs

Consider a point $P(\cos\theta, \sin\theta)$ on the unit circle in the Cartesian plane. If the distance from the origin to the point $Q(x, y)$ is 5, and the angle between the positive x-axis and the line segment OQ is α, such that $\cos\theta = \frac{3}{5}$ and $\sin\theta = \frac{4}{5}$. If $x = \cos\alpha$ and $y = \sin\alpha$, find the maximum possible value of the product of the x-coordinates of P and Q, i.e., $\cos\theta \cdot \cos\alpha$.

a. 4/5

b. 3/5

c. **1** ✓

d. 5/3

The point P is on the unit circle, so its coordinates are $(\cos\theta, \sin\theta)$. We are given $\cos\theta = \frac{3}{5}$ and $\sin\theta = \frac{4}{5}$. The point Q has coordinates (x, y) and its distance from the origin is 5. The angle between the positive x-axis and OQ is α. Thus, $x = 5\cos\alpha$ and $y = 5\sin\alpha$. We are asked to find the maximum possible value of the product of the x-coordinates of P and Q, which is $\cos\theta \cdot x$. Substituting the given values, we h.

© ClipSAT — clipsat.org | All Rights Reserved

30 [Intermediate] • Target: Converting between degrees and radians

Consider an arithmetic sequence where the 5th term is 19 and the 11th term is 43. If the first term is denoted by a_1 and the common difference by d, find the value of $a_1 + d$.

a. 6

b. **7 ✓**

c. 8

d. 9

Let the arithmetic sequence be defined by $a_n = a_1 + (n-1)d$. We are given that $a_5 = 19$ and $a_{11} = 43$. From the given information, we can set up a system of two linear equations: 1. $a_1 + (5-1)d = 19 \Rightarrow a_1 + 4d = 19$ 2. $a_1 + (11-1)d = 43 \Rightarrow a_1 + 10d = 43$ Subtract equation (1) from equation (2): $(a_1 + 10d) - (a_1 + 4d) = 43 - 19$ $6d = 24$ $d = \frac{24}{6} = 4$ Now substitute the value of d back into equation (1): $a_1 + 4(4) = 19$ $a_1 + 16 = 19$ $a_1 = 19 - 16 = 3$ We are aske.

31 [Intermediate] • Target: Arc length and sector area

Consider the arithmetic sequence where the first term is $a_1 = 5$ and the common difference is $d = 3$. What is the 4th term of this sequence, a_4?

a. 11

b. **14 ✓**

c. 17

d. 20

An arithmetic sequence is defined by the formula $a_n = a_1 + (n-1)d$, where a_n is the nth term, a_1 is the first term, and d is the common difference. In this problem, we are given $a_1 = 5$ and $d = 3$. We need to find the 4th term, so $n = 4$. Substituting these values into the formula: $a_4 = 5 + (4-1) \times 3$ $a_4 = 5 + (3) \times 3$ $a_4 = 5 + 9$ $a_4 = 14$ Alternatively, we can list the terms: $a_1 = 5$ $a_2 = a_1 + d = 5 + 3 = 8$ $a_3 = a_2 + d = 8 + 3 = 11$ $a_4 = a_3 + d = 11 + 3 = 14$ Therefore.

32 [Intermediate] • Target: Angular and linear speed

Consider the arithmetic sequence where the first term is $a_1 = 5$ and the common difference is $d = 3$. What is the fourth term of this sequence, a_4?

a. **14 ✓**

b. 11

c. 17

d. 8

The formula for the n-th term of an arithmetic sequence is $a_n = a_1 + (n-1)d$. Given $a_1 = 5$ and $d = 3$, we want to find a_4. Substituting $n = 4$ into the formula: $a_4 = 5 + (4-1) \times 3$ $a_4 = 5 + (3) \times 3$ $a_4 = 5 + 9$ $a_4 = 14$. Therefore, the fourth term of the sequence is 14. Common errors might occur from miscalculating $(n-1)$ or incorrectly applying the addition.

© ClipSAT — clipsat.org | All Rights Reserved

33 [Intermediate] • Target: Coterminal angles and reference angles

Find the angle between $\mathbf{u} = \langle 2, 1 \rangle$ and $\mathbf{v} = \langle 4, 2 \rangle$ (to one decimal place).

a. **0.0°** ✓

b. 20.0°

c. 180.0°

d. 10.0°

$\cos\theta = \frac{\mathbf{u}\cdot\mathbf{v}}{|\mathbf{u}||\mathbf{v}|} = \frac{10}{2.24 \times 4.47} \approx 1.0$, so $\theta \approx 0.0°$.

34 [Intermediate] • Target: The radian definition itself

Using the limit definition $f'(x) = \lim_{h\to 0} \frac{f(x+h)-f(x)}{h}$, find $f'(x)$ for $f(x) = 4x^2 - x + 4$.

a. $4x^2 - 1$

b. $8x + 4$

c. $4x - 1$

d. $8x - 1$ ✓

Expand $f(x+h) - f(x) = 4(2xh + h^2) - 1h = (8x-1)h + 4h^2$. Dividing by h: $\frac{f(x+h)-f(x)}{h} = 8x - 1 + 4h$. Letting $h \to 0$: $f'(x) = 8x - 1$.

35 [Intermediate] • Target: Combined arc length and sector area problems

Find the 4th term of the arithmetic sequence with $a_1 = 6$ and common difference $d = -5$.

a. -4

b. -19

c. -14

d. **-9** ✓

Applying Sequences & Series principles, $a_n = a_1 + (n-1)d = 6 + (4-1)(-5) = -9$. Therefore, the verified correct option is -9.

36 [Intermediate] • Target: Common angles in both units (reference table)

Find a positive coterminal angle with $400°$ in $[0°, 360°)$.

a. 30

b. **40** ✓

c. 31

d. 400

Applying Trigonometry principles, $400° - 360° = 40°$. Therefore, the verified correct option is 40.

© ClipSAT — clipsat.org | All Rights Reserved

37 [Intermediate] • Target: Satellite and orbital speed applications

A sector OAB has radius 8 cm and angle $\angle AOB = 1.2$ rad. Find the area of the minor segment cut off by chord AB, leaving your answer in exact form.

a. $32(1.2 - \sin(1.2))$ ✓

b. $64(1.2 - \sin(1.2))$

c. $33(1.2 - \sin(1.2))$

d. $32(1.2 + \sin(1.2))$

Segment $=$ Sector $-$ Triangle $= \frac{1}{2}r^2\theta - \frac{1}{2}r^2\sin\theta = 32(1.2 - \sin(1.2))$ cm^2

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Negative angles and full-turn arcs • Cross-Lesson: 7.04 Applications: Growth and Decay

Find the reference angle of 120°.

a. 120

b. **60** ✓

c. 31

d. 30

Applying Trigonometry principles, Reference angle $= 60^\circ$. Therefore, the verified correct option is 60.

39 [Challenging] • Target: Converting between degrees and radians • Cross-Lesson: 8.01 Rational Functions and Asymptotes

Find the angle between $\mathbf{u} = \langle 1, 0\rangle$ and $\mathbf{v} = \langle 0, 1\rangle$.

a. 135°

b. 60°

c. 90° ✓

d. 120°

$\cos\theta = \frac{\mathbf{u}\cdot\mathbf{v}}{|\mathbf{u}||\mathbf{v}|} = 0 \Rightarrow \theta = 90^\circ$.

© ClipSAT — clipsat.org | All Rights Reserved

40 [Challenging] • Target: Arc length and sector area • Cross-Lesson: 8.02 Graphing Rational Functions

For the ellipse $\frac{x^2}{81} + \frac{y^2}{9} = 1$, find the length of the major axis and the foci.

a. Major axis 18, foci $(\pm\sqrt{72}, 0)$ ✓

b. Major axis 9, foci $(\pm\sqrt{72}, 0)$

c. Major axis 18, foci $(0, \pm\sqrt{72})$

d. Major axis 6, foci $(\pm\sqrt{72}, 0)$

$a = 9 > b = 3$, so major axis $= 2a = 18$ (along x-axis). $c^2 = a^2 - b^2 = 72$, foci at $(\pm\sqrt{72}, 0)$.

41 [Challenging] • Target: Angular and linear speed • Cross-Lesson: 8.03 Introduction to Limits

Find the common ratio of the geometric sequence $2, 6, 18, 54, \ldots$.

a. 3 ✓

b. 2

c. -3

d. 4

Applying Sequences principles, $r = \frac{6}{2} = 3$. Therefore, the verified correct option is 3.

42 [Challenging] • Target: Coterminal angles and reference angles • Cross-Lesson: 9.01 Arithmetic Sequences and Series

Find the area of the polar region bounded by $r = 2$ for $0 \le \theta \le \frac{\pi}{6}$.

a. $\frac{2\pi}{3}$

b. $\frac{\pi}{3}$ ✓

c. $\frac{\pi}{6}$

d. $\frac{4\pi}{3}$

$A = \frac{1}{2}r^2\theta = \frac{1}{2}(2)^2(\frac{\pi}{6}) = \frac{\pi}{3}$.

43 [Challenging] • Target: The radian definition itself • Cross-Lesson: 9.02 Geometric Sequences and Series

A sector has radius 10 cm and angle 1.7 rad. Find its area.

a. 170

b. 95

c. 85 ✓

d. 17

$A = \frac{1}{2}r^2\theta = \frac{1}{2}(10)^2(1.7) = 85$ cm^2

© ClipSAT — clipsat.org | All Rights Reserved

44 [Challenging] • Target: Combined arc length and sector area problems • Cross-Lesson: 9.03 The Binomial Theorem

A geometric sequence has common ratio $r = 3$ and $a_4 = 54$. Find a_1.

a. 5

b. 54

c. **2** ✓

d. 18

$a_1 = \frac{a_4}{r^3} = \frac{54}{3^3} = 2$.

45 [Challenging] • Target: Common angles in both units (reference table) • Cross-Lesson: 1.01 Functions and Function Notation

For the ellipse $\frac{x^2}{25} + \frac{y^2}{9} = 1$, what is the length of the major axis?

a. 6

b. **10** ✓

c. 5

d. 8

$a^2 = 25 \Rightarrow a = 5$; major axis $= 2a = 10$.

© ClipSAT — clipsat.org | All Rights Reserved