
Vectors and Embeddings in AI

Section 1: Vector Spaces and Linear Combinations

1 Given word vectors $\mathbf{u} = [2, -1, 3]^T$ and $\mathbf{v} = [1, 4, -2]^T$, compute the linear combination $3\mathbf{u} - 2\mathbf{v}$.



a. $[4, -11, 13]^T$

b. $[4, 5, 5]^T$

c. $[8, -11, 5]^T$

d. $[7, 5, 13]^T$

2 Two embedding vectors $\mathbf{a} = [k, 2, -1]^T$ and $\mathbf{b} = [3, -4, 2]^T$ are orthogonal in semantic space. Find the value of k.

a. $k = \frac{10}{3}$

b. $k = 2$

c. $k = -\frac{10}{3}$

d. $k = 5$

3 Determine whether the three feature vectors $\mathbf{v}_1 = [1, 0]^T$, $\mathbf{v}_2 = [0, 1]^T$, and $\mathbf{v}_3 = [3, -2]^T$ are linearly independent in $\mathbf{R}^2$.

a. Linearly dependent (any set of > 2 vectors in $\mathbf{R}^2$ is dependent)

b. Linearly independent

c. Orthogonal set

d. Cannot be determined without a matrix inversion

Section 2: Vector Norms and Sparsity

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 Calculate the Manhattan (L_1) norm and Euclidean (L_2) norm of the embedding error vector $\mathbf{e} = [3, -4, 12]^T$.

a. $\|\mathbf{e}\|_1 = 19, \quad \|\mathbf{e}\|_2 = 13$

b. $\|\mathbf{e}\|_1 = 13, \quad \|\mathbf{e}\|_2 = 19$

c. $\|\mathbf{e}\|_1 = 11, \quad \|\mathbf{e}\|_2 = 13$

d. $\|\mathbf{e}\|_1 = 19, \quad \|\mathbf{e}\|_2 = 169$


5 Find the Chebyshev (L_∞) norm of the parameter vector $\mathbf{w} = [-7, 2, 5, -9, 4]^T$.

a. 9

b. 7

c. 27

d. 5



6 In Lasso regression, why does adding an L_1 penalty to the loss function induce exact feature sparsity (weights driven to 0)?

a. The diamond-shaped L_1 contour has sharp corners on the coordinate axes

b. The L_1 norm is continuously differentiable everywhere

c. The L_1 penalty penalizes large weights much more heavily than L_2

d. The L_1 norm computes the determinant of the covariance matrix

---

Section 3: Dot Products, Cosine Similarity and Angular Distance

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 Two query embeddings are $\mathbf{u} = [1, 1, 0]^T$ and $\mathbf{v} = [0, 2, 2]^T$. Compute the cosine similarity $\cos\theta$ between them.

a. 0.500

b. 0.707

c. 0.250

d. 1.000

8 If two unit-normalized embeddings have cosine similarity $\cos\theta = -1.0$, what is their semantic relationship?

a. Diametrically opposite in direction (180° apart)

b. Identical in meaning

c. Orthogonal / unrelated

d. Collinear in the exact same direction


9 Compute the cosine distance $D_{\cos}(\mathbf{u}, \mathbf{v}) = 1 - \cos\theta$ for two orthogonal feature vectors.

a. 1.0

b. 0.0

c. 2.0


d. -1.0



Section 4: High-Dimensional Geometry and Projections

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

10 Find the scalar projection of vector $\mathbf{a} = [4, 2]^T$ onto the direction of unit vector $\mathbf{u} = [0.6, 0.8]^T$.



a. 4.0

b. 5.0

c. 2.4

d. 3.2

11 In high-dimensional embedding spaces ($D = 1024$), why are two randomly selected vectors almost certainly nearly orthogonal?

a. The surface area of a high-dimensional hypersphere concentrates near the equator relative to any pole

b. All high-dimensional vectors have zero norm

c. The dot product of random Gaussian variables always equals 1

d. The cosine function approaches infinity as dimension increases

12 Normalize the feature vector $\mathbf{x} = [3, 0, 4]^T$ to have unit Euclidean norm.

a. $[0.6, 0, 0.8]^T$

b. $[0.75, 0, 1]^T$

c. $[1, 0, 1]^T$

d. $[0.3, 0, 0.4]^T$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647