PCA and Dimensionality Reduction

Section 1: Data Centering and the Covariance Matrix


1 Why must data features be centered (subtracting feature means) prior to performing Principal Component Analysis?

a. So the first principal component aligns with the axis of maximum variance rather than the offset from origin

b. To make the data non-linear

c. To ensure all eigenvalues are equal to 1

d. To reduce the matrix dimension



2 For a zero-mean data matrix $\mathbf{X} \in \mathbf{R}^{N \times D}$, state the sample covariance matrix $\mathbf{\Sigma}$.

a. $\mathbf{\Sigma} = \frac{1}{N-1}\mathbf{X}^T\mathbf{X}$

b. $\mathbf{\Sigma} = \frac{1}{D-1}\mathbf{X}\mathbf{X}^T$

c. $\mathbf{\Sigma} = \mathbf{X}^T\mathbf{X}$

d. $\mathbf{\Sigma} = \frac{1}{N}\mathbf{X}^2$

3 What is the mathematical symmetry property of the covariance matrix $\mathbf{\Sigma}$?

a. It is symmetric and positive semi-definite (all eigenvalues $\lambda_i \geq 0$)

b. It is skew-symmetric

c. It is an orthogonal matrix

d. It has negative trace

Section 2: Eigenvalues, Eigenvectors and Variance Maximization

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 A 2×2 covariance matrix has entries $\Sigma_{11} = 4$, $\Sigma_{22} = 1$, and covariance $\Sigma_{12} = \Sigma_{21} = 2$. Find its eigenvalues.

a. $\lambda_1 = 5, \quad \lambda_2 = 0$

b. $\lambda_1 = 4, \quad \lambda_2 = 1$

c. $\lambda_1 = 3, \quad \lambda_2 = 2$

d. $\lambda_1 = 2, \quad \lambda_2 = 2$


5 What is the geometric meaning of the first eigenvector $\mathbf{v}_1$ corresponding to largest eigenvalue λ_1?

a. The direction along which the data exhibits maximum variance

b. The direction perpendicular to all data points

c. The direction with the lowest information content

d. The vector of class centroids



6 If $\mathbf{v}$ is an eigenvector of $\mathbf{\Sigma}$ with eigenvalue λ, what is the variance of data projected onto $\mathbf{v}$?

a. λ

b. $\sqrt{\lambda}$

c. λ^2

d. 1

Section 3: Explained Variance and Scree Analysis

7 A dataset has eigenvalues $\lambda_1 = 70, \lambda_2 = 20, \lambda_3 = 10$. What percentage of total variance is explained by the first two principal components?

a. 90%

b. 70%

c. 80%

d. 20%

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

8 How is an 'elbow' in a scree plot (eigenvalues plotted against component index) used in practice?

a. To select the optimal cutoff k where adding further components yields diminishing returns

b. To detect outliers in the target labels

c. To determine the learning rate for gradient descent

d. To test for multicollinearity


9 If a dataset has 100 features and you retain the top 10 principal components, by what percentage is feature dimensionality reduced?

a. 90%

b. 10%

c. 50%

d. 80%



Section 4: SVD Connection and Applications


10 Given the Singular Value Decomposition $\mathbf{X} = \mathbf{USV}^T$, how are the principal component directions $\mathbf{v}_i$ directly obtained?

a. As the right singular vectors (columns of $\mathbf{V}$)

b. As the left singular vectors (columns of $\mathbf{U}$)

c. As the diagonal elements of $\mathbf{S}$

d. As the eigenvalues of $\mathbf{U}$



© ClipSAT: By Mr. Mohamed Abdallah +966597688647

11 What is the mathematical relation between singular values s_i of $\mathbf{X}$ and eigenvalues λ_i of covariance matrix $\mathbf{\Sigma}$?

a. $\lambda_i = \frac{s_i^2}{N-1}$

b. $\lambda_i = s_i$

c. $\lambda_i = \sqrt{s_i}$

d. $\lambda_i = s_i(N-1)$

12 Name a major practical application of PCA in image processing and computer vision.

a. Eigenfaces: facial compression and feature representation in lower-dimensional eigenspace

b. Image sharpening through edge convolution

c. Color palette inversion

d. Bilinear image interpolation

© ClipSAT: By Mr. Mohamed Abdallah +966597688647