
Nonhomogeneous Equations & Undetermined Coefficients

Section 1: General Solution Structure

1 What is the structure of the general solution to a linear nonhomogeneous equation $L[y] = g(x)$?



a. $y(x) = y_c(x) + y_p(x)$, where y_c is the complementary solution and y_p is any particular solution

b. $y(x) = y_c(x) \cdot y_p(x)$

c. $y(x) = y_p(x)$ only

d. $y(x) = c_1 y_p(x) + c_2 y_c(x)$

2 Find a particular solution y_p to $y'' - 5y' + 6y = 12$.

a. $y_p = 2$

b. $y_p = 12$

c. $y_p = 2x$

d. $y_p = 6$

3 What is the trial particular solution y_p for $y'' + 4y = 3x^2 - 1$?

a. $y_p = Ax^2 + Bx + C$

b. $y_p = Ax^2$

c. $y_p = Ax^2 + B$

d. $y_p = Ax^3 + Bx^2$

Section 2: Exponential and Sinusoidal Forcing Functions

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 Find a particular solution to $y'' + y = 4e^{2x}$.

a. $y_p = \frac{4}{5}e^{2x}$

b. $y_p = 4e^{2x}$

c. $y_p = 2e^{2x}$

d. $y_p = \frac{1}{2}e^{2x}$


5 What is the correct undetermined coefficients form for $y'' + 9y = 5\cos(2x)$?

a. $y_p = A\cos(2x) + B\sin(2x)$

b. $y_p = A\cos(2x)$

c. $y_p = Ax\cos(2x)$

d. $y_p = A\sin(2x)$



6 Find y_p for $y'' + 9y = 5\cos(2x)$.

a. $y_p = \cos(2x)$

b. $y_p = 5\cos(2x)$

c. $y_p = \frac{5}{13}\cos(2x)$

d. $y_p = \sin(2x)$

Section 3: The Modification Rule (Resonance / Overlap)

7 When must a trial particular solution y_p be multiplied by x (or x^2)?

a. When any term in the trial solution is already a solution of the associated homogeneous equation

b. Whenever the forcing function is a polynomial

c. When the coefficient of y' is negative

d. Only when the right-hand side is zero

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

8 For $y'' - 4y' + 4y = 3e^{2x}$, the homogeneous solution is $y_c = (c_1 + c_2 x)e^{2x}$. What is the correct trial form for y_p?

a. $y_p = Ax^2 e^{2x}$


b. $y_p = Ae^{2x}$

c. $y_p = Axe^{2x}$

d. $y_p = Ax^3 e^{2x}$

9 What physical phenomenon occurs when an undamped mechanical system is driven at its natural resonance frequency ($y'' + \omega_0^2 y = F_0 \cos(\omega_0 t)$)?

a. Resonance, where amplitude grows linearly with time ($y_p = Ct\sin(\omega_0 t)$)




b. Immediate decay to zero

c. Critical damping

d. Chaotic motion

Section 4: The Principle of Superposition

10 If y_{p1} solves $L[y] = g_1(x)$ and y_{p2} solves $L[y] = g_2(x)$, what solves $L[y] = 2g_1(x) - 3g_2(x)$?



a. $2y_{p1} - 3y_{p2}$

b. $y_{p1}y_{p2}$

c. $\frac{y_{p1}}{y_{p2}}$

d. $y_{p1} + y_{p2}$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

11 Can the Method of Undetermined Coefficients be used to solve $y'' + y = \tan x$?

a. No, because derivatives of $\tan x$ produce infinitely many non-terminating forms; Variation of Parameters is required

b. Yes, with trial solution $A\tan x + B\sec x$

c. Yes, with trial solution $A\sin x/\cos x$

d. Only if x is small

12 Find a particular solution to $y'' + 4y = 8x + 12$.

a. $y_p = 2x + 3$

b. $y_p = 2x$

c. $y_p = 4x + 3$

d. $y_p = 2x^2 + 3x$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647