
Variation of Parameters

Section 1: The Wronskian Determinant

1 What is the definition of the Wronskian $W(y_1, y_2)$ of two differentiable functions $y_1(x)$ and $y_2(x)$?



a. $W(y_1, y_2) = y_1 y_2' - y_1' y_2 = \det[[y_1, y_2], [y_1', y_2']]$

b. $W(y_1, y_2) = y_1 y_2 + y_1' y_2'$

c. $W(y_1, y_2) = y_1' y_2 - y_1 y_2'$

d. $W(y_1, y_2) = \frac{y_1}{y_2}$

2 Compute the Wronskian of $y_1 = \cos x$ and $y_2 = \sin x$.

a. 1

b. 0

c. -1

d. $\cos(2x)$

3 What does $W(y_1, y_2) \neq 0$ guarantee about solutions y_1 and y_2 to a homogeneous linear ODE?

a. They are linearly independent and form a fundamental set of solutions

b. They are orthogonal

c. They are periodic

d. One must be zero

Section 2: The Variation of Parameters Formula

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 For the second-order equation $y'' + P(x)y' + Q(x)y = g(x)$, what is the variation of parameters formula for y_p?

a. $y_p = -y_1 \int \frac{y_2 g}{W} dx + y_2 \int \frac{y_1 g}{W} dx$

b. $y_p = y_1 \int \frac{y_2 g}{W} dx - y_2 \int \frac{y_1 g}{W} dx$

c. $y_p = \int \frac{g(x)}{W} dx$

d. $y_p = -y_2 \int \frac{y_1}{W} dx$


5 What must the coefficient of y'' be before applying the variation of parameters formula?

a. Exactly 1 (must be in standard monic form)

b. Any non-zero constant

c. Zero

d. Positive



6 For $y'' + y = \sec x$, where $y_1 = \cos x, y_2 = \sin x$, what is $u_1' = -\frac{y_2 g}{W}$?

a. $-\tan x$

b. $-\sin x$

c. -1

d. $-\cos x$

Section 3: Solving Advanced Nonhomogeneous Equations

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

7 Find $u_1 = \int(-\tan x)dx$ for the problem $y'' + y = \sec x$.

a. $\ln|\cos x|$

b. $-\ln|\cos x|$

c. $\sec^2 x$

d. $-\cos x$

8 Find $u_2' = \frac{y_1 g}{W}$ for $y'' + y = \sec x$ and evaluate u_2.

a. $u_2 = x$

b. $u_2 = \sin x$

c. $u_2 = \cos x$

d. $u_2 = \tan x$


9 Assemble the particular solution $y_p = u_1 y_1 + u_2 y_2$ for $y'' + y = \sec x$.

a. $y_p = \cos x \ln|\cos x| + x\sin x$

b. $y_p = \cos x + x\sin x$

c. $y_p = \ln|\cos x| + x$

d. $y_p = \sin x \ln|\cos x| + x\cos x$



Section 4: Comparison of Nonhomogeneous Methods

© ClipSAT: By Mr. Mohamed Abdallah +966597688647


10 What is the major advantage of Variation of Parameters over Undetermined Coefficients?

a. It works for any continuous forcing function $g(x)$, not just polynomials, exponentials, or sines/cosines

b. It never requires any integration

c. It works without knowing the homogeneous solution y_c

d. It always produces algebraic rather than transcendental answers



11 What is the main drawback of Variation of Parameters compared to Undetermined Coefficients?

a. The integrals $\int \frac{y_i g}{W} dx$ may be difficult or impossible to evaluate in closed elementary form

b. It only works on first-order equations

c. It gives only approximate numerical solutions

d. It cannot handle initial value problems

12 What is Abel's formula for the Wronskian of solutions to $y'' + P(x)y' + Q(x)y = 0$?

a. $W(x) = Ce^{-\int P(x)dx}$

b. $W(x) = Ce^{\int Q(x)dx}$

c. $W(x) = C(P(x) + Q(x))$

d. $W(x) = Cx$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647