The Dot Product

Section 1: Algebraic Definition and Properties


1 Find the dot product $\mathbf{u} \cdot \mathbf{v}$ for $\mathbf{u} = \langle 4, -3 \rangle$ and $\mathbf{v} = \langle 2, 5 \rangle$.

a. -7

b. 23

c. $\langle 8, -15 \rangle$

d. 1



2 What is the value of $\mathbf{v} \cdot \mathbf{v}$ for any vector $\mathbf{v}$?

a. $\|\mathbf{v}\|^2$

b. $\|\mathbf{v}\|$

c. 0

d. 1

3 Which property does NOT hold for the dot product?

a. $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$ (Commutative)

b. $c(\mathbf{u} \cdot \mathbf{v}) = (c\mathbf{u}) \cdot \mathbf{v}$

c. $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$

d. $\mathbf{u} \cdot (\mathbf{v} \cdot \mathbf{w}) = (\mathbf{u} \cdot \mathbf{v}) \cdot \mathbf{w}$ (Associative)

Section 2: Geometric Definition and Angle Between Vectors

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

4 State the formula for the cosine of the angle θ between two non-zero vectors $\mathbf{u}$ and $\mathbf{v}$.

a. $\cos\theta = \frac{\mathbf{u}\cdot\mathbf{v}}{\|\mathbf{u}\|\|\mathbf{v}\|}$

b. $\cos\theta = \frac{\|\mathbf{u}\|\|\mathbf{v}\|}{\mathbf{u}\cdot\mathbf{v}}$

c. $\cos\theta = \mathbf{u}\cdot\mathbf{v}$

d. $\cos\theta = \frac{\mathbf{u}+\mathbf{v}}{\|\mathbf{u}+\mathbf{v}\|}$


5 Find the angle between $\mathbf{u} = \langle 1, 1\rangle$ and $\mathbf{v} = \langle 0, 2\rangle$.

a. 45°

b. 90°

c. 30°

d. 60°



6 If the dot product $\mathbf{u}\cdot\mathbf{v} < 0$, what type of angle is θ?

a. Obtuse ($90^\circ < \theta \leq 180^\circ$)

b. Acute ($0^\circ \leq \theta < 90^\circ$)

c. Right (90°)

d. Straight (180°)

Section 3: Orthogonality and Parallel Vectors

7 Two non-zero vectors $\mathbf{u}$ and $\mathbf{v}$ are orthogonal (perpendicular) if and only if:

a. $\mathbf{u}\cdot\mathbf{v} = 0$

b. $\mathbf{u}\cdot\mathbf{v} = 1$

c. $\mathbf{u}\times\mathbf{v} = \mathbf{0}$

d. $\|\mathbf{u}\| = \|\mathbf{v}\|$

© ClipSAT: By Mr. Mohamed Abdallah +966597688647

8 Find k such that $\mathbf{u} = \langle 3, k \rangle$ is orthogonal to $\mathbf{v} = \langle 4, 6 \rangle$.

a. -2

b. 2

c. $-1/2$

d. 12


9 Which pair of vectors is parallel?

a. $\langle 2, -3 \rangle$ and $\langle -6, 9 \rangle$

b. $\langle 2, 3 \rangle$ and $\langle 3, 2 \rangle$

c. $\langle 1, 0 \rangle$ and $\langle 0, 1 \rangle$

d. $\langle 4, 1 \rangle$ and $\langle -1, 4 \rangle$



Section 4: Vector Projections and Work


10 State the formula for the vector projection of $\mathbf{u}$ onto $\mathbf{v}$ ($\text{proj}_{\mathbf{v}}\mathbf{u}$).

a. $\left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|^2}\right)\mathbf{v}$

b. $\left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|^2}\right)\mathbf{u}$

c. $\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{v}\|}$

d. $(\mathbf{u} \cdot \mathbf{v})\mathbf{v}$



© ClipSAT: By Mr. Mohamed Abdallah +966597688647

11 Find $\text{proj}_{\mathbf{v}}\mathbf{u}$ for $\mathbf{u} = \langle 6, 2\rangle$ and $\mathbf{v} = \langle 4, 0\rangle$.

a. $\langle 6, 0\rangle$

b. $\langle 4, 0\rangle$

c. $\langle 0, 2\rangle$

d. $\langle 24, 0\rangle$

12 A force $\mathbf{F} = \langle 30, 40\rangle$ N moves a cart along the displacement $\mathbf{d} = \langle 5, 0\rangle$ m. Calculate the work done.

a. 150 J

b. 200 J

c. 350 J

d. 250 J

© ClipSAT: By Mr. Mohamed Abdallah +966597688647