

Advanced Plane Geometry

ANSWER KEY



Circle theorems: tangent-chord and secant angles

- 1 Two secants are drawn from an external point to a circle, intercepting arcs of 100° and 30° . Find the measure of the angle formed at the external point, using the formula (difference of arcs) $\div 2$.

$$\text{Angle} = \frac{100^\circ - 30^\circ}{2} = \frac{70^\circ}{2} = 35^\circ.$$

- 2 Two chords intersect inside a circle, intercepting arcs of 80° and 40° . Find the measure of the angle formed at the intersection, using the formula (sum of arcs) $\div 2$.

$$\text{Angle} = \frac{80^\circ + 40^\circ}{2} = \frac{120^\circ}{2} = 60^\circ.$$

Power of a point (secant and tangent segment relationships)

- 3 A tangent segment from an external point has length 8, and a secant from the same point passes through the circle with external segment length 4. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

$$8^2 = 4 \times (\text{whole secant}), \text{ so } 64 = 4w, \text{ giving } w = 16.$$

Area of regular polygons

- 4 Find the area of a regular pentagon with side length 6 and apothem 4.13, using $A = \frac{1}{2} \times \text{perimeter} \times \text{apothem}$.

$$\text{Perimeter} = 5(6) = 30. \text{ Area} = \frac{1}{2}(30)(4.13) = 61.95.$$

Similar solids and scale factor in 3D

- 5 Two similar cylinders have a scale factor of $\frac{2}{3}$. If the smaller cylinder has volume 40, find the volume of the larger cylinder.

$$\text{Volumes of similar solids scale by the CUBE of the linear scale factor: } \left(\frac{3}{2}\right)^3 = \frac{27}{8}. \text{ Larger volume: } 40 \times \frac{27}{8} = 135.$$

Composite 3D solids

- 6 A solid consists of a cylinder of radius 3 and height 8, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

Cylinder volume: $\pi r^2 h = \pi(9)(8) = 72\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(27) = 18\pi$. Total volume: $72\pi + 18\pi = 90\pi$.

Triangle centers: centroid and circumcenter

- 7 Find the centroid of the triangle with vertices (2, 3), (8, 3), (5, 9), using the formula (average of the vertex coordinates).

$$\text{Centroid} = \left(\frac{2 + 8 + 5}{3}, \frac{3 + 3 + 9}{3} \right) = \left(\frac{15}{3}, \frac{15}{3} \right) = (5, 5).$$

Heron's formula

- 8 A triangle has sides 7, 8, and 9. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

$$s = \frac{7 + 8 + 9}{2} = 12. \text{ Area} = \sqrt{12(12-7)(12-8)(12-9)} = \sqrt{12(5)(4)(3)} = \sqrt{720} = 12\sqrt{5}.$$

A multi-part circle geometry problem

- 9 This question has two parts. A circle has radius 10. A chord is drawn 6 units from the center. (a) Find the length of the chord, using the relationship between the radius, the distance from center to chord, and half the chord length (a right triangle relationship). (b) Find the area of the minor segment cut off by this chord, to the nearest whole number, given that the central angle subtended by the chord is approximately 106.3° and $\sin(106.3^\circ) \approx 0.960$.

(a) Half the chord length: $\sqrt{10^2 - 6^2} = \sqrt{100 - 36} = \sqrt{64} = 8$. Full chord length: 16. (b) Segment area = sector area – triangle area. Sector area: $\frac{106.3}{360} \times \pi(10)^2 \approx 0.2953 \times 314.16 \approx 92.8$. Triangle area: $\frac{1}{2}(10)(10)\sin(106.3^\circ) \approx 50(0.960) = 48$. Segment area: $92.8 - 48 \approx 45$.

Angle bisectors and the angle bisector theorem

- 10 In triangle ABC , the angle bisector from A meets side BC at point D , dividing it such that $BD = 6$ and $DC = 9$. If $AB = 8$, find AC , using the angle bisector theorem ($\frac{BD}{DC} = \frac{AB}{AC}$).

$$\frac{6}{9} = \frac{8}{AC}, \text{ so } 6 \cdot AC = 72, \text{ giving } AC = 12.$$

The 30-60-90 and 45-45-90 triangles in composite figures

- 11** An equilateral triangle has side length 10. Find its height and area, using the fact that the height splits it into two 30-60-90 triangles.

The height corresponds to the longer leg of a 30-60-90 triangle with hypotenuse 10 and shorter leg 5:
height = $5\sqrt{3}$. Area = $\frac{1}{2}(10)(5\sqrt{3}) = 25\sqrt{3}$.

Circumscribed and inscribed circles

- 12** A square is inscribed in a circle of radius 5. Find the side length of the square, using the fact that the diagonal of the square equals the diameter of the circle.

Diameter = $2(5) = 10$. For a square, diagonal = $s\sqrt{2}$, so $s\sqrt{2} = 10$, giving $s = \frac{10}{\sqrt{2}} = 5\sqrt{2}$.

Regular polygon interior and exterior angles

- 13** Find the measure of each interior angle and each exterior angle of a regular octagon.

Sum of interior angles of an n -gon: $(n - 2)180^\circ$. For $n = 8$: $(6)(180^\circ) = 1080^\circ$. Each interior angle:
 $\frac{1080}{8} = 135^\circ$. Each exterior angle (supplementary to the interior angle): $180^\circ - 135^\circ = 45^\circ$.

A multi-part composite figure and circle problem

- 14** This question has three parts. A running track consists of a rectangle with two semicircular ends. The rectangle is 80 meters long and 40 meters wide, and the semicircles are attached to the two 40-meter-wide ends. (a) Find the radius of each semicircle. (b) Find the total perimeter (distance around) the track, in terms of π . (c) Find the total enclosed area of the track shape, in terms of π .

(a) The semicircles are attached to the 40-meter ends, so the diameter of each semicircle is 40, giving radius = 20. (b) Perimeter = $2(\text{rectangle length}) + 2(\text{semicircle arc length})$. The two semicircles together form one full circle's circumference: $2\pi(20) = 40\pi$. Total perimeter: $2(80) + 40\pi = 160 + 40\pi$ meters. (c) Area = rectangle area + combined semicircle area. Rectangle area: $80 \times 40 = 3200$. The two semicircles together form one full circle: $\pi(20)^2 = 400\pi$. Total area: $3200 + 400\pi$ square meters.

Multiple choice — easy

- 15** Two chords intersect inside a circle, intercepting arcs of 170° and 76° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 246°
- b. 133°
- c. 47°
- d. 123° ✓

$$\text{Angle} = \frac{170^\circ + 76^\circ}{2} = \frac{246^\circ}{2} = 123^\circ.$$

- 16** Two secants are drawn from an external point to a circle, intercepting arcs of 46° and 16° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 30°
- b. 25°
- c. 15° ✓
- d. 31°

$$\text{Angle} = \frac{46^\circ - 16^\circ}{2} = \frac{30^\circ}{2} = 15^\circ.$$

- 17** Two secants are drawn from an external point to a circle, intercepting arcs of 157° and 43° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 67°
- b. 114°
- c. 100°
- d. 57° ✓

$$\text{Angle} = \frac{157^\circ - 43^\circ}{2} = \frac{114^\circ}{2} = 57^\circ.$$

- 18 Two chords intersect inside a circle, intercepting arcs of 85° and 69° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 8°
- b. 154°
- c. 77° ✓
- d. 87°

$$\text{Angle} = \frac{85^\circ + 69^\circ}{2} = \frac{154^\circ}{2} = 77^\circ.$$

- 19 Two chords intersect inside a circle, intercepting arcs of 112° and 164° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 276°
- b. 148°
- c. 138° ✓
- d. 26°

$$\text{Angle} = \frac{112^\circ + 164^\circ}{2} = \frac{276^\circ}{2} = 138^\circ.$$

- 20 Two chords intersect inside a circle, intercepting arcs of 117° and 17° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 134°
- b. 50°
- c. 77°
- d. 67° ✓

$$\text{Angle} = \frac{117^\circ + 17^\circ}{2} = \frac{134^\circ}{2} = 67^\circ.$$

- 21 Two secants are drawn from an external point to a circle, intercepting arcs of 95° and 37° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 58°
- b. 29° ✓
- c. 39°
- d. 66°

$$\text{Angle} = \frac{95^\circ - 37^\circ}{2} = \frac{58^\circ}{2} = 29^\circ.$$

- 22 Two chords intersect inside a circle, intercepting arcs of 12° and 104° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 68°
- b. 58° ✓
- c. 46°
- d. 116°

$$\text{Angle} = \frac{12^\circ + 104^\circ}{2} = \frac{116^\circ}{2} = 58^\circ.$$

- 23 Two chords intersect inside a circle, intercepting arcs of 86° and 38° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 62° ✓
- b. 72°
- c. 124°
- d. 24°

$$\text{Angle} = \frac{86^\circ + 38^\circ}{2} = \frac{124^\circ}{2} = 62^\circ.$$

- 24** Two secants are drawn from an external point to a circle, intercepting arcs of 118° and 34° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 84°
- b. 42° ✓
- c. 52°
- d. 76°

$$\text{Angle} = \frac{118^\circ - 34^\circ}{2} = \frac{84^\circ}{2} = 42^\circ.$$

- 25** Two secants are drawn from an external point to a circle, intercepting arcs of 184° and 14° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 99°
- b. 170°
- c. 85° ✓
- d. 95°

$$\text{Angle} = \frac{184^\circ - 14^\circ}{2} = \frac{170^\circ}{2} = 85^\circ.$$

- 26** Two chords intersect inside a circle, intercepting arcs of 83° and 157° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 130°
- b. 240°
- c. 37°
- d. 120° ✓

$$\text{Angle} = \frac{83^\circ + 157^\circ}{2} = \frac{240^\circ}{2} = 120^\circ.$$

- 27 Two secants are drawn from an external point to a circle, intercepting arcs of 118° and 52° . Find the measure of the angle formed at the external point, using the formula $(\text{difference of arcs}) \div 2$.

- a. 43°
- b. 33° ✓
- c. 85°
- d. 66°

$$\text{Angle} = \frac{118^\circ - 52^\circ}{2} = \frac{66^\circ}{2} = 33^\circ.$$

- 28 Two chords intersect inside a circle, intercepting arcs of 98° and 92° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 190°
- b. 95° ✓
- c. 3°
- d. 105°

$$\text{Angle} = \frac{98^\circ + 92^\circ}{2} = \frac{190^\circ}{2} = 95^\circ.$$

- 29 Two chords intersect inside a circle, intercepting arcs of 127° and 31° . Find the measure of the angle formed at the intersection, using the formula $(\text{sum of arcs}) \div 2$.

- a. 89°
- b. 79° ✓
- c. 158°
- d. 48°

$$\text{Angle} = \frac{127^\circ + 31^\circ}{2} = \frac{158^\circ}{2} = 79^\circ.$$

Multiple choice — medium

- 30** A tangent segment from an external point has length 30, and a secant from the same point passes through the circle with external segment length 20. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 25
- b. 30
- c. 45 ✓
- d. 46

$30^2 = 20 \times (\text{whole secant})$, so $900 = 20w$, giving $w = 45$.

- 31** A tangent segment from an external point has length 18, and a secant from the same point passes through the circle with external segment length 12. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 28
- b. 15
- c. 27 ✓
- d. 18

$18^2 = 12 \times (\text{whole secant})$, so $324 = 12w$, giving $w = 27$.

- 32** A tangent segment from an external point has length 28, and a secant from the same point passes through the circle with external segment length 16. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 33
- b. 49 ✓
- c. 28
- d. 50

$28^2 = 16 \times (\text{whole secant})$, so $784 = 16w$, giving $w = 49$.

- 33 A tangent segment from an external point has length 24, and a secant from the same point passes through the circle with external segment length 18. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 24
- b. 14
- c. 33
- d. 32 ✓

$24^2 = 18 \times (\text{whole secant})$, so $576 = 18w$, giving $w = 32$.

- 34 A tangent segment from an external point has length 8, and a secant from the same point passes through the circle with external segment length 4. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 17
- b. 12
- c. 16 ✓
- d. 8

$8^2 = 4 \times (\text{whole secant})$, so $64 = 4w$, giving $w = 16$.

- 35 A tangent segment from an external point has length 6, and a secant from the same point passes through the circle with external segment length 2. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 6
- b. 18 ✓
- c. 19
- d. 16

$6^2 = 2 \times (\text{whole secant})$, so $36 = 2w$, giving $w = 18$.

- 36** A tangent segment from an external point has length 22, and a secant from the same point passes through the circle with external segment length 11. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 45
- b. 44 ✓
- c. 33
- d. 22

$22^2 = 11 \times (\text{whole secant})$, so $484 = 11w$, giving $w = 44$.

- 37** A tangent segment from an external point has length 6, and a secant from the same point passes through the circle with external segment length 3. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 9
- b. 6
- c. 12 ✓
- d. 13

$6^2 = 3 \times (\text{whole secant})$, so $36 = 3w$, giving $w = 12$.

- 38** A tangent segment from an external point has length 8, and a secant from the same point passes through the circle with external segment length 2. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

- a. 32 ✓
- b. 30
- c. 33
- d. 8

$8^2 = 2 \times (\text{whole secant})$, so $64 = 2w$, giving $w = 32$.

- 39 A tangent segment from an external point has length 4, and a secant from the same point passes through the circle with external segment length 2. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 8 ✓

b. 4

c. 9

d. 6

$4^2 = 2 \times (\text{whole secant})$, so $16 = 2w$, giving $w = 8$.

- 40 A tangent segment from an external point has length 15, and a secant from the same point passes through the circle with external segment length 5. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 40

b. 45 ✓

c. 15

d. 46

$15^2 = 5 \times (\text{whole secant})$, so $225 = 5w$, giving $w = 45$.

- 41 A tangent segment from an external point has length 12, and a secant from the same point passes through the circle with external segment length 4. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 12

b. 37

c. 36 ✓

d. 32

$12^2 = 4 \times (\text{whole secant})$, so $144 = 4w$, giving $w = 36$.

- 42 A tangent segment from an external point has length 6, and a secant from the same point passes through the circle with external segment length 4. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 6
b. 10
c. 5
d. 9 ✓

$6^2 = 4 \times (\text{whole secant})$, so $36 = 4w$, giving $w = 9$.

- 43 A tangent segment from an external point has length 12, and a secant from the same point passes through the circle with external segment length 9. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 16 ✓
b. 12
c. 7
d. 17

$12^2 = 9 \times (\text{whole secant})$, so $144 = 9w$, giving $w = 16$.

- 44 A tangent segment from an external point has length 20, and a secant from the same point passes through the circle with external segment length 10. Find the length of the entire secant segment, using $(\text{tangent})^2 = (\text{external secant})(\text{whole secant})$.

a. 20
b. 41
c. 30
d. 40 ✓

$20^2 = 10 \times (\text{whole secant})$, so $400 = 10w$, giving $w = 40$.

Multiple choice — hard

- 45 A triangle has sides 9, 12, and 15. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

- a. 52
- b. 54 ✓
- c. 36
- d. 56

$$s = \frac{9 + 12 + 15}{2} = 18. \text{ Area} = \sqrt{18(9)(6)(3)} = \sqrt{2916} = 54.$$

- 46 A triangle has sides 6, 8, and 10. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

- a. 26
- b. 27
- c. 24 ✓
- d. 22

$$s = \frac{6 + 8 + 10}{2} = 12. \text{ Area} = \sqrt{12(6)(4)(2)} = \sqrt{576} = 24.$$

- 47 A triangle has sides 13, 14, and 15. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

- a. 86
- b. 82
- c. 42
- d. 84 ✓

$$s = \frac{13 + 14 + 15}{2} = 21. \text{ Area} = \sqrt{21(8)(7)(6)} = \sqrt{7056} = 84.$$

- 48 A triangle has sides 10, 10, and 16. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

- a. 46
- b. 36
- c. 48 ✓
- d. 50

$$s = \frac{10 + 10 + 16}{2} = 18. \text{ Area} = \sqrt{18(8)(8)(2)} = \sqrt{2304} = 48.$$

- 49 A triangle has sides 15, 15, and 18. Find its area using Heron's formula, $A = \sqrt{s(s-a)(s-b)(s-c)}$ where s is the semi-perimeter.

- a. 106
- b. 110
- c. 48
- d. 108 ✓

$$s = \frac{15 + 15 + 18}{2} = 24. \text{ Area} = \sqrt{24(9)(9)(6)} = \sqrt{11664} = 108.$$

- 50 A solid consists of a cylinder of radius 3 and height 3, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

- a. 45π ✓
- b. 48π
- c. 27π
- d. 18π

Cylinder volume: $\pi r^2 h = \pi(9)(3) = 27\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(27) = 18\pi$. Total volume: $27\pi + 18\pi = 45\pi$.

- 51 A solid consists of a cylinder of radius 6 and height 11, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

a. 540π ✓

b. 396π

c. 546π

d. 144π

Cylinder volume: $\pi r^2 h = \pi(36)(11) = 396\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(216) = 144\pi$. Total volume: $396\pi + 144\pi = 540\pi$.

- 52 A solid consists of a cylinder of radius 6 and height 12, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

a. 144π

b. 582π

c. 576π ✓

d. 432π

Cylinder volume: $\pi r^2 h = \pi(36)(12) = 432\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(216) = 144\pi$. Total volume: $432\pi + 144\pi = 576\pi$.

- 53 A solid consists of a cylinder of radius 3 and height 5, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

a. 66π

b. 18π

c. 45π

d. 63π ✓

Cylinder volume: $\pi r^2 h = \pi(9)(5) = 45\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(27) = 18\pi$. Total volume: $45\pi + 18\pi = 63\pi$.

- 54 A solid consists of a cylinder of radius 9 and height 4, topped with a hemisphere of the same radius. Find the total volume, in terms of π .

a. 810π ✓

b. 324π

c. 819π

d. 486π

Cylinder volume: $\pi r^2 h = \pi(81)(4) = 324\pi$. Hemisphere volume: $\frac{2}{3}\pi r^3 = \frac{2}{3}\pi(729) = 486\pi$. Total volume: $324\pi + 486\pi = 810\pi$.

- 55 Two similar cylinders have a scale factor of $\frac{3}{4}$. If the smaller cylinder has volume 54, find the volume of the larger cylinder.

a. 96

b. 54

c. 128 ✓

d. 131

Volumes of similar solids scale by the CUBE of the linear scale factor: $\left(\frac{4}{3}\right)^3 = \frac{64}{27}$. Larger volume: $54 \times \frac{64}{27} = 128$.

- 56 Two similar cylinders have a scale factor of $\frac{2}{3}$. If the smaller cylinder has volume 32, find the volume of the larger cylinder.

a. 110

b. 32

c. 108 ✓

d. 72

Volumes of similar solids scale by the CUBE of the linear scale factor: $\left(\frac{3}{2}\right)^3 = \frac{27}{8}$. Larger volume: $32 \times \frac{27}{8} = 108$.

- 57 Two similar cylinders have a scale factor of $\frac{5}{8}$. If the smaller cylinder has volume 750, find the volume of the larger cylinder.

a. 3072 ✓

b. 750

c. 1920

d. 3077

Volumes of similar solids scale by the CUBE of the linear scale factor: $\left(\frac{8}{5}\right)^3 = \frac{512}{125}$. Larger volume:
 $750 \times \frac{512}{125} = 3072$.

- 58 Two similar cylinders have a scale factor of $\frac{3}{5}$. If the smaller cylinder has volume 162, find the volume of the larger cylinder.

a. 753

b. 450

c. 162

d. 750 ✓

Volumes of similar solids scale by the CUBE of the linear scale factor: $\left(\frac{5}{3}\right)^3 = \frac{125}{27}$. Larger volume:
 $162 \times \frac{125}{27} = 750$.

- 59 Two similar cylinders have a scale factor of $\frac{4}{7}$. If the smaller cylinder has volume 256, find the volume of the larger cylinder.

a. 784

b. 256

c. 1376

d. 1372 ✓

Volumes of similar solids scale by the CUBE of the linear scale factor: $\left(\frac{7}{4}\right)^3 = \frac{343}{64}$. Larger volume:
 $256 \times \frac{343}{64} = 1372$.