Simultaneous Equations and Inequalities

ClipSAT Printable Practice Suite (clipsat.org) • Topic 2.01


Part I — Free-Response Questions (FRQs): Foundational Fluency (1–7)

1 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Simultaneous Equations & Inequalities · Practice 4] If the governing relation for this system is $g(t) = 6t + 6$, evaluate $g(10)$ and find the inverse input t when $g(t) = 72$.

2 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

Find the set of values of x satisfying the inequality $2x^2 - 7x + 3 \leq 0$.



3 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Differential Equations · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

4 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises


The temperature θ of a cooling metal rod satisfies $\frac{d\theta}{dt} = -k(\theta - 20)$, where t is time in minutes and k is a constant. Given $\theta = 80$ when $t = 0$ and $\theta = 50$ when $t = 10$, find the value of t when $\theta = 30$.

5 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Differentiation & Integration · Practice 4] If the governing relation for this system is $g(t) = 6t + 6$, evaluate $g(10)$ and find the inverse input t when $g(t) = 72$.

6 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Trigonometry · Practice 4] If the governing relation for this system is $g(t) = 6t + 6$, evaluate $g(10)$ and find the inverse input t when $g(t) = 72$.



© ClipSAT — clipsat.org | All Rights Reserved

7 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

[About AS Level · Practice 2] If the governing relation for this system is $g(t) = 4t + 4$, evaluate $g(6)$ and find the inverse input t when $g(t) = 32$.

Part II — Free-Response Questions (FRQs): Intermediate Multi-Step Application (8–14)

8 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises


[Binomial Distribution · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

9 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Coordinate Geometry · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

10 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Representation of Data · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.



11 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Discrete Random Variables · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

12 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Energy, Work & Power · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

© ClipSAT — clipsat.org | All Rights Reserved

13 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Forces & Equilibrium · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

14 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

[Graphs & Transformations · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.


Part III — Free-Response Questions (FRQs): Challenging & Cross-Lesson Synthesis (15–20)

15 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 3.01 Indices, Surds and Polynomials

[Kinematics · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

16 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 4.01 Logarithmic and Exponential Functions

[Newton's Laws of Motion · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.



17 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 5.01 Trigonometric Functions and Circular Measure

[Normal Distribution · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

18 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 6.01 Permutations and Combinations

[Circular Measure · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

19 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 7.01 Differentiation and Rates of Change

[Series & the Binomial Theorem · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

© ClipSAT — clipsat.org | All Rights Reserved


20 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 8.01 Integration and Kinematics

[Statistics & Probability · Practice 3] If the governing relation for this system is $g(t) = 5t + 5$, evaluate $g(8)$ and find the inverse input t when $g(t) = 50$.

Part IV — Multiple-Choice Questions (MCQs): Foundational Concept Mastery (21–28)

21 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

The diagram shows the graphs of two linear equations. The point of intersection of the two lines represents the solution to the system of simultaneous equations. Find the coordinates of the point of intersection.



a. (1, 2)

b. (2, 1)

c. (3, 0)

d. (0, 3)

22 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

The diagram shows the graphs of the linear equation $y = 2x + c$ and the quadratic equation $y = -x^2 + 4x + 5$. The two graphs intersect at the points where $x = -1$ and $x = k$. Find the value of c and the value of k.

a. c = 3, k = 4

b. c = 5, k = 3


c. c = 3, k = 3

d. c = 5, k = 4

© ClipSAT — clipsat.org | All Rights Reserved

23 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

Find the x-coordinates of the points of intersection of $y = x^2 - 9x + 22$ and $y = 2$.



a. x=4 or x=5

b. x=-4 or x=-5

c. x=5 or x=5

d. x=4 or x=4

24 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

Solve the simultaneous equations $2x + 3y = 18$ and $3x - 4y = 10$.

a. x=7, y=2

b. x=-6, y=-2

c. x=6, y=2

d. x=2, y=6

25 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

A curve has equation $y = x\sqrt{16 - x^2}$ for $0 \le x \le 4$. A point $P(x, y)$ moves along the curve in the first quadrant, and the normal to the curve at P intersects the x-axis at the point N. Given that the x-coordinate of P is increasing at a constant rate of $\sqrt{3}$ units s^{-1}, find the rate of change of the area of triangle OPN, where O is the origin, at the instant when $x = 2$.

a. -6 units^2 s^{-1}

b. 6 units^2 s^{-1}

c. -12 units^2 s^{-1}

d. $-2\sqrt{3}$ units^2 s^{-1}

© ClipSAT — clipsat.org | All Rights Reserved

26 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

A sector OAB of a circle with centre O and radius 12 cm has angle $\angle AOB = \frac{\pi}{2}$ radians. The point C is the midpoint of the radius OA. The line passing through C perpendicular to OA intersects the circular arc AB at the point D. The straight line segment BC is drawn. Find the exact area, in cm^2, of the region bounded by the line segment CD, the circular arc BD, and the line segment BC.

a. $12\pi + 18\sqrt{3} - 36$

b. $24\pi + 18\sqrt{3} - 36$

c. $12\pi + 36\sqrt{3} - 36$

d. $18\pi + 18\sqrt{3} - 36$

27 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

In the expansion of $(1 + ax)^n$ in ascending powers of x, the coefficient of x^2 is 60 and the coefficient of x^3 is 160, where n is a positive integer and $a > 0$. A cubic polynomial $P(x)$, in which the coefficient of x^3 is 1, has $(x - a)^2$ as a factor and leaves a remainder of n when divided by $(x - 1)$. Find the remainder when $P(x)$ is divided by $(x + a)$.

a. 48

b. -48


c. -112

d. 0

© ClipSAT — clipsat.org | All Rights Reserved

28 [Foundational] • Target: Simultaneous Equations & Inequalities Practice Exercises

The line $y = 7x - 4$ is tangent to the curve $y = ax^2 + \frac{b}{x}$ at the point where $x = 2$, where a and b are constants and $xe0$. Find the coordinates of the stationary point of the curve and determine its nature.



a. $(1, 6)$, which is a local minimum

b. $(1, 6)$, which is a local maximum

c. $(-1, -2)$, which is a local minimum

d. $(2, 10)$, which is a local minimum

Part V — Multiple-Choice Questions (MCQs): Intermediate Exam Application (29–37)

29 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

The curve C has equation $y = x^3 - 3x + 2$. The tangent to C at the point $P(2, 4)$ intersects the curve again at point Q. Find the area of the finite region completely enclosed between the curve C and the tangent line.

a. 108

b. 84

c. 64

d. 12

© ClipSAT — clipsat.org | All Rights Reserved

30 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

A storage capsule consists of a right circular cylinder of radius r cm and height h cm, closed with a flat circular base of radius r cm at the bottom, and surmounted by a hemispherical dome of radius r cm at the top. The total exterior surface area of the capsule is fixed at 180π cm^2. During a heating process, the radius r expands at a constant rate of 0.4 cm s^{-1}. Find the rate of change of the total volume of the capsule, in cm^{3}s^{-1}, at the instant when the height of the cylindrical section is 16.5 cm.

a. 20π

b. 50π

c. $\frac{36\pi}{5}$

d. 11π

31 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

The variables x and y, where $x > 0$, satisfy the relationship $y = ax + \frac{b}{x}$ where a and b are positive constants. When xy is plotted on the vertical axis against x^2 on the horizontal axis, a straight line is obtained which passes through the points $(2, 11)$ and $(5, 23)$. Given further that x and y satisfy the logarithmic equation $\log_3(y - ax) + \log_9(x^2y^2) = 1 + \log_3(x + 10)$ find the sum of all possible values of x.

a. $\frac{10}{3}$

b. $\frac{8}{3}$

c. $\frac{7}{2}$

d. $\frac{13}{3}$

32 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

A group of n students, which includes Ali and Priya, are to be arranged in a straight line for a photograph. The number of different arrangements in which Ali and Priya are not standing next to each other is 3600. Find the value of the integer n.

a. 6

b. 7


c. 8

d. 9

© ClipSAT — clipsat.org | All Rights Reserved

33 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

Relative to a fixed origin O, ship A has an initial position vector of $(24\mathbf{i} + 10\mathbf{j})$ km and moves with a constant velocity of $(-2\mathbf{i} + 5\mathbf{j})$ km h^{-1}. Ship B has an initial position vector of $(-20\mathbf{i} - 18\mathbf{j})$ km and moves with a constant velocity of $(u\mathbf{i} + 12\mathbf{j})$ km h^{-1}, where u is a constant. Given that ship B intercepts ship A, find the speed of ship B.



a. 15 km h^{-1}

b. 9 km h^{-1}

c. 21 km h^{-1}

d. 34 km h^{-1}

34 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

Two parallel lines, L_1 and L_2, have n and $n + 2$ distinct points marked on them respectively, where n is an integer with $n \geq 2$. Triangles are formed by choosing 3 of these points as vertices such that they are not collinear. Quadrilaterals are formed by choosing 4 of these points as vertices such that no three are collinear. Given that the ratio of the number of quadrilaterals to the number of triangles that can be formed is $6 : 5$, find the total number of triangles that can be formed.

a. 175

b. 210

c. 220

d. 96

35 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

In triangle ABC, the side lengths are $AB = (x + 2)$ cm, $BC = (3x - 1)$ cm, and $AC = 7$ cm. Given that $\angle ABC = 60^\circ$ and $x > 0$, find the exact area of triangle ABC.

a. $10\sqrt{3}$ cm^2

b. $5\sqrt{3}$ cm^2

c. $20\sqrt{3}$ cm^2

d. 10 cm^2

© ClipSAT — clipsat.org | All Rights Reserved

36 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

Variables x and y satisfy the equation $y = Ax^k$, where A and k are constants, with $x > 0$. The graph of $\ln y$ against $\ln x$ is a straight line passing through the points $(\ln 2, \ln 72)$ and $(\ln 6, \ln 8)$. Find the value of x when $y = 18$.

a. 4

b. 2

c. 1

d. 8

37 [Intermediate] • Target: Simultaneous Equations & Inequalities Practice Exercises

The variables x and y are connected by the equation $y = \sqrt{2x + 5}$. Given that x increases at a constant rate of 6 units per second, find the rate of change of y with respect to time when $x = 2$.

a. 2 units per second

b. 1 unit per second

c. $\frac{1}{3}$ units per second

d. 12 units per second

Part VI — Multiple-Choice Questions (MCQs): Challenging & Cross-Lesson Synthesis (38–45)

38 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 6.01 Permutations and Combinations

The variables x and y satisfy the relationship $y = Ax^n$, where A and n are constants. When $\log_2 y$ is plotted against $\log_2 x$, a straight line is obtained which passes through the points $(1, 5)$ and $(4, -1)$. What is the value of y when $x = 4$?

a. 8

b. $\frac{1}{2}$


c. 3

d. 32

© ClipSAT — clipsat.org | All Rights Reserved

39 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 7.01 Differentiation and Rates of Change

The polynomial $P(x) = 2x^3 + ax^2 + bx - 6$ has a factor of $x - 2$ and leaves a remainder of -24 when divided by $x + 1$, where a and b are constants. Find the remainder when $P(x)$ is divided by $x - 3$.



a. 12

b. -30

c. 60

d. -150

40 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 8.01 Integration and Kinematics

A particle moves in a straight line such that its acceleration, a m s^{-2}, at time t seconds is given by $a = 6t - 18$. The particle is at instantaneous rest when $t = 1$. Given that the displacement of the particle from a fixed point O is $s = -10$ m when $t = 2$, find the displacement of the particle, in metres, when it is next at instantaneous rest.

a. -37

b. -25

c. -13

d. -5

41 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 1.01 Functions, Modulus and Quadratics

A committee of 5 members is to be selected from a group consisting of n men and 6 women, where $n \geq 4$. Given that the number of possible committees containing exactly 3 men is equal to the number of possible committees containing exactly 4 men, find the value of n.

a. 13

b. 10

c. 8

d. 7

© ClipSAT — clipsat.org | All Rights Reserved

42 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 3.01 Indices, Surds and Polynomials

In the expansion of $(1+ax)^n$ in ascending powers of x, where n is a positive integer and a is a constant, the coefficient of x is -12 and the coefficient of x^2 is 60. Find the coefficient of x^3.

a. -160

b. 160

c. -40

d. -240

43 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 4.01 Logarithmic and Exponential Functions

A curve has equation $y = 6x - x^2$. The straight line $y = mx$, where $0 < m < 6$, intersects the curve at the origin and at the point P. Given that the area of the region enclosed between the curve and the line is $\frac{32}{3}$, find the value of the constant m.

a. 2

b. 4

c. 3

d. 1

44 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 5.01 Trigonometric Functions and Circular Measure

A particle moves in a straight line such that its velocity, v m s^{-1}, at time t seconds is given by $v = 6t^2 - 18t + k$, where k is a constant. Given that the particle is momentarily at rest when $t = 2$, find the total distance traveled by the particle in the first 2 seconds.

a. 6 m

b. 4 m

c. 5 m

d. 8 m

45 [Challenging] • Target: Simultaneous Equations & Inequalities Practice Exercises • Cross-Lesson: 6.01 Permutations and Combinations

The variables x and y satisfy the equation $y = Ab^x$, where A and b are positive constants. When $\log_2 y$ is plotted against x, a straight line is obtained that passes through the points $(2, 7)$ and $(5, 13)$. Find the value of y when $x = 4$.

a. 2048

b. 1024

c. 11

d. 256

© ClipSAT — clipsat.org | All Rights Reserved