

THE IRON FIX

A Woman's Guide to Understanding
Iron Deficiency, Reclaiming Your
Energy, and Feeling Like Yourself
Again

Hannah Wells

THE IRON FIX

A Woman's Guide to Understanding Iron Deficiency, Reclaiming Your Energy, and Feeling Like Yourself Again

Copyright © 2026 Hannah Wells

All rights reserved.

ISBN: 978-0-000-00000-0

Published by Pink Mango Publishing

To every woman who has been told she is —just
tired,
and to every doctor who never thought to check her
ferritin —
this is the conversation we should have been having
all along.

Contents

01. foreword

02. ch01

03. ch02

04. ch03

05. ch04

06. ch05

07. ch06

08. ch07

09. ch08

10. ch09

11. ch10

12. ch11

13. resources

foreword

I SPENT THREE YEARS BEING TOLD I WAS TIRED BECAUSE I WAS BUSY.

I was a woman in my thirties with a full life — work, family, exercise, social obligations — and every doctor I saw nodded knowingly when I described the exhaustion. "You're doing too much," they said. "Try cutting back. Try sleeping more. Try magnesium."

I tried all of it. I went to bed earlier. I cut out caffeine after noon. I took magnesium glycinate and ashwagandha and a dozen other supplements recommended by wellness blogs. I reduced my social calendar and said no to things and practiced the kind of gentle self-care that is marketed to tired women like a solution to a problem that runs deeper than any of it can reach.

Nothing changed. The fatigue was not the kind that sleep fixes. It was the kind that lives underneath everything — a constant, low-grade heaviness that made every task feel like it required more energy than I had. I forgot words. I lost my train of speech mid-sentence. My hair was coming out in clumps in the shower. I was pale in a way that people noticed. "Are you feeling okay?" they asked, and I didn't know how to say: I don't think I've felt okay in years, but I can't remember what okay feels like, so maybe this is just what being a woman my age is supposed to be.

After three years, four doctors, and a growing suspicion that I was imagining the whole thing, a nurse practitioner ran a ferritin test on a whim. Ferritin is a measure of your body's iron stores — not the iron circulating in your blood right now, but the iron you have saved for a rainy day.

My ferritin was 12 ng/mL. The lab's "normal" range started at 15. Technically, I was within range. Clinically, I was running on empty.

Iron deficiency is the most common nutritional deficiency in the world, and women are disproportionately affected. One in three premenopausal women in the United States has low iron stores. Among women with heavy periods, the number rises to one in two. And yet it is routinely missed, dismissed, or treated as an afterthought — because the symptoms (fatigue, brain fog, hair loss, mood changes, dizziness, cold hands and feet) overlap with almost every other condition in women's health, and because the "normal" range for ferritin is set so low that millions of women are walking around technically "normal" while their bodies are starving for iron.

This book is the conversation I wish I had been part of from the beginning. It covers the full picture: what iron actually does in your body, why women lose so much of it, the hidden symptoms that go beyond simple fatigue, how to get the right tests and interpret them honestly, how to supplement without destroying your digestive system, how to eat to build your stores, how to manage heavy

periods, what pregnancy and postpartum do to your iron levels, and how iron affects your brain, your mood, your hair, and your sense of yourself.

Every chapter ends with a toolkit you can use immediately. The tone is warm, direct, and grounded in real experience — because I have been where you are, and I know what it feels like to be told you are fine when you know you are not.

One honest note before we begin: I am not a doctor. I am a health writer who has spent years researching iron deficiency and living with its consequences. This book is a guide, not a prescription. Talk to your doctor before starting any supplement, especially iron — because iron is not a vitamin you can pee out if you take too much. It is a mineral that accumulates in your body, and getting the dose right matters.

You are not lazy. You are not making this up. You are not weak. You may be iron deficient — and if you are, there is a path back to feeling like yourself again.

This is where it starts.

Toolkit Summary

- **Iron deficiency is the most common nutritional deficiency in women** — 1 in 3 premenopausal women have low stores. It is routinely missed.

- **"Normal" ferritin does not mean optimal ferritin** — a level of 15 ng/mL may be "in range" but is far too low for energy, hair growth, and brain function.
- **Your symptoms are real** — fatigue, brain fog, hair loss, mood changes, dizziness, and cold extremities are all predictable effects of low iron.
- **This is a practical guide** — every chapter ends with tools you can use immediately.
- **I am not a doctor** — this book is a companion, not a prescription. Talk to your doctor before starting supplements.
- **You are not alone** — millions of women are navigating this same silent deficiency. Knowledge is the first step toward reclaiming your energy.

ch01

LET'S START WITH A NUMBER THAT MIGHT STOP YOU IN YOUR TRACKS: ONE IN THREE PREMENOPAUSAL WOMEN IN THE UNITED STATES HAS LOW IRON STORES. ONE IN THREE. AMONG WOMEN WITH HEAVY PERIODS, THAT NUMBER CLIMBS TO ONE IN TWO. IF YOU ARE READING THIS BOOK, THE ODDS ARE BETTER THAN GOOD THAT YOU ARE ONE OF THEM — OR THAT YOU LOVE SOMEONE WHO IS.

These numbers are not pulled from a small study or a fringe theory. They come from the Centers for Disease Control and Prevention and are backed by decades of global research. The World Health Organization has called iron deficiency the most common nutritional deficiency in the world, and women bear the brunt of it. And yet most women who are iron deficient have no idea. They are not told. They are not tested. They are not even asked.

This is the silent epidemic in women's health — a condition so widespread that it touches nearly every family, every friend group, every workplace — and so routinely overlooked that millions of women spend years, sometimes decades, feeling terrible without ever knowing why.

Why Women Lose Iron

If you have ever wondered why iron deficiency is so much more common in women than in men, the answer starts with biology.

The average woman loses between 30 and 80 milliliters of blood during each menstrual period. That does not sound like much until you do the math: over the course of a year, that blood loss adds up to a significant drain on your body's iron reserves. One milliliter of blood contains roughly half a milligram of iron. A woman with an average period loses about 15 to 25 milligrams of iron per cycle. A woman with heavy periods — and nearly a third of women meet the medical definition of heavy menstrual bleeding — can lose 50 milligrams or more per cycle, month after month.

Your body does not have an unlimited supply of iron to give. It has to rebuild those stores from food and supplements between periods. And for many women, the gap between what they lose and what they can recoup is too wide to close.

Pregnancy makes it worse. During pregnancy, your body produces roughly 50 percent more blood to support the growing baby. That extra blood volume requires a massive increase in iron — about 1,000 milligrams of total iron over the course of the pregnancy. Some of that goes to the baby, some to the placenta, some to the increase in your own blood volume. After birth, blood loss during delivery can drain another significant portion. It is not

uncommon for a woman to enter pregnancy with adequate iron stores and emerge from the postpartum period severely depleted.

Then there is diet. Even with the best intentions, getting enough iron from food alone is genuinely difficult for many women, especially if you do not eat red meat or if you follow a vegetarian or vegan diet. The type of iron found in plant foods — non-heme iron — is absorbed less efficiently than the iron found in animal foods. You can eat bowls of spinach and lentils and still fall short if your body cannot absorb enough of what is passing through your digestive system. We will talk about how to fix this in the food chapter, but for now, know that diet alone often is not enough for women with higher iron losses.

Absorption issues add another layer. Conditions like celiac disease, inflammatory bowel disease, Crohn's disease, ulcerative colitis, and even less severe gut issues can impair your body's ability to absorb iron from food and supplements. If you have any of these conditions, your risk of iron deficiency is significantly higher, regardless of how much iron you consume. The same is true if you have had gastric bypass surgery or any procedure that alters your digestive tract.

And here is something that does not get talked about enough: common medications can interfere with iron absorption. Acid-reducing medications like proton pump inhibitors (omeprazole, esomeprazole, lansoprazole) and H2 blockers (famotidine, ranitidine) reduce stomach acid,

and stomach acid is necessary for absorbing non-heme iron. If you take these medications daily, you may be absorbing far less iron from your food than you think.

The Silent Epidemic

Here is the part that should make you angry: most women who are iron deficient do not know it.

This is not because the signs are invisible. It is because the signs are normalized. Fatigue is explained away by a busy life. Brain fog is blamed on perimenopause or not sleeping enough. Hair loss is dismissed as stress. Dizziness is attributed to low blood sugar or not drinking enough water. Mood changes are labeled anxiety or depression. Every symptom of iron deficiency is a symptom of something else, and the default assumption — by doctors, by friends, by the culture at large — is that women are just tired because they are doing too much.

The result is that women spend years cycling through doctors, specialists, and diagnoses. They are prescribed antidepressants when their iron stores are critically low. They are told to try yoga, meditation, and better sleep hygiene. They are advised to cut back on commitments and practice self-care. None of this is bad advice on its own, but none of it will fix a body that is starved for iron.

One study published in the *Journal of the American Board of Family Medicine* found that women with iron deficiency visited their doctors more frequently, received more prescriptions, and were more likely to be diagnosed

with depression and anxiety — all before anyone checked their ferritin. Another study found that iron-deficient women who had not yet developed anemia reported significantly lower quality of life, more fatigue, and more cognitive difficulties than women with normal iron stores, but fewer than half had ever been tested.

The silence is built into the system. Routine blood work often does not include ferritin testing — the key measure of your iron stores. Standard complete blood counts (CBCs) check hemoglobin, which tells you if you have anemia, but hemoglobin can be perfectly normal while your iron stores are already running on fumes. By the time hemoglobin drops, you have been iron deficient for months, possibly years.

The Spectrum: Iron Deficiency vs. Iron Deficiency Anemia

One of the most important things to understand — and one of the least explained by doctors — is that iron deficiency and iron deficiency anemia are not the same thing. They exist on a spectrum, and most symptomatic women are somewhere along it long before they reach the anemia end.

Think of it like a bank account. Iron stores — measured as ferritin — are your savings account. When you lose iron through bleeding or use it for daily bodily functions, your body withdraws from savings first. If you replenish

those savings through food and supplements, everything stays balanced. But if you keep withdrawing more than you deposit, your savings account eventually runs low.

Stage one is **iron depletion**. Your ferritin is low — typically under 30 ng/mL — but your hemoglobin is still normal. Your body has enough iron to keep your red blood cells working, but it has no backup reserves. This is like having your checking account fully funded but your savings account empty. You can function, but you have no buffer. Your body is running on what it can get from day to day, with nothing saved for a rainy day.

Stage two is **iron-deficient erythropoiesis**. Your ferritin is very low, and your body is starting to struggle to make enough healthy red blood cells. Your hemoglobin may still be in the normal range, but it is trending downward. Your body is now prioritizing iron for red blood cell production over other functions — your hair, your skin, your brain, your immune system. This is when symptoms like hair loss, brittle nails, and brain fog often appear.

Stage three is **iron deficiency anemia**. Your ferritin is critically low, your hemoglobin has dropped below the normal range, and your red blood cells are small and pale. This is the point at which your body can no longer transport enough oxygen to meet even basic needs. The fatigue becomes profound. You may be short of breath walking up a flight of stairs. You may feel dizzy when you stand up. Your heart may beat faster to compensate for the lack of oxygen.

The critical insight here is that most women with iron deficiency never reach stage three. They spend months or years in stages one and two — symptomatic, struggling, but not technically anemic. And because anemia is what most doctors look for, they are told their iron is fine while their ferritin sits at 18 and their hair is falling out in handfuls.

If you have been told your iron is normal but you do not feel normal, this is why. Normal is not the same as optimal. We will talk about what optimal actually looks like in the testing chapter, but for now, know this: ferritin under 30 is absolute iron deficiency, and many experts now argue that ferritin under 50 can cause symptoms in physically active women. The conventional lab range — often as low as 15 to 150 — was never designed to identify optimal health. It was designed to rule out severe disease.

Who Is at Risk

The list of women who are at higher risk for iron deficiency is long, and if you see yourself here, you should pay attention.

Women with heavy periods. If your periods are heavy enough that you change pads or tampons every hour, pass clots larger than a quarter, or bleed for more than seven days, you are losing significantly more iron than the

average woman. Heavy menstrual bleeding is the single most common cause of iron deficiency in women of reproductive age.

Postpartum women. Pregnancy, childbirth, and breastfeeding drain enormous amounts of iron. Many women enter the postpartum period already depleted, and the months of healing and nursing can keep iron stores low for years if not addressed.

Vegetarians and vegans. Plant-based iron is harder to absorb than the iron found in meat. Vegetarian and vegan women need to be more intentional about iron intake and absorption. Studies consistently show lower ferritin levels in women who do not eat red meat.

Athletes. Exercise increases your body's need for iron. Endurance athletes, runners, and anyone who trains intensely loses iron through sweat, through foot-strike hemolysis (the breakdown of red blood cells in the feet during running), and through increased red blood cell turnover. Female athletes are at particularly high risk.

Women with gastrointestinal conditions. Celiac disease, Crohn's disease, ulcerative colitis, and other conditions that affect the digestive tract can impair iron absorption. Even if you eat plenty of iron, your body may not be able to absorb it. Some of these conditions go undiagnosed for years — and occult gastrointestinal bleeding is another hidden source of iron loss.

Women taking certain medications. As mentioned, acid-reducing medications, oral contraceptives (which can cause breakthrough bleeding in some women), and NSAIDs (which can cause gastrointestinal bleeding) can all affect iron status.

Blood donors. Regular blood donation removes iron from your body. Women who donate blood several times a year are at increased risk for iron depletion, especially if they have other risk factors.

Women with endometriosis, fibroids, or adenomyosis. These conditions can cause heavy menstrual bleeding, chronic blood loss, and in some cases, internal bleeding that further depletes iron stores.

If any of these describe you, consider this an early warning. Your risk is higher. Your need for awareness — and possibly for testing and supplementation — is higher too.

The Cost of Undiagnosed Deficiency

Iron deficiency does not stay in your body. It radiates outward into every part of your life.

The lost productivity is staggering. Studies have found that iron deficiency costs billions of dollars annually in reduced work performance, missed workdays, and healthcare utilization. Women with untreated iron deficiency report significantly lower energy levels, more

difficulty concentrating, and reduced physical capacity — all of which affect their ability to show up fully at work and at home.

But the human cost is harder to quantify. It is the promotion you did not pursue because you were too tired to prepare for the interview. The workout you skipped because you could not muster the energy. The social event you declined because the effort of getting ready and making conversation felt overwhelming. The days you spent feeling like a lesser version of yourself, wondering what was wrong with you.

And then there is the misdiagnosis. Iron deficiency is a master mimicker. Its symptoms overlap with depression, anxiety, chronic fatigue syndrome, thyroid disorders, ADHD, and perimenopause. Women are diagnosed with these conditions — and treated for them — without anyone ever checking their ferritin. Antidepressants are prescribed. Therapy is recommended. Lifestyle changes are suggested. And some of these interventions are genuinely helpful, but none of them will work if the underlying problem is iron.

One study found that women with iron deficiency were twice as likely to be diagnosed with depression as women with normal iron stores. Another found that iron supplementation improved depressive symptoms in women with low ferritin, even when they had not responded to antidepressant medication. The implication is profound: how many women are being treated for depression when what they actually need is iron?

This is not to say that every case of fatigue or low mood is really iron deficiency. But it is to say that iron deficiency should be on the short list of possibilities — and right now, for most women, it is not even considered.

The Path Forward

Here is the good news: iron deficiency is treatable. It is not a mystery. It is not something you have to live with. It is a well-understood condition with well-established solutions — testing, supplementation, dietary changes, and when necessary, medical interventions to address the underlying cause of blood loss.

But the first step is knowledge. You cannot fix what you do not know is broken. And the most common reason women go untreated for iron deficiency is not that treatment is difficult — it is that they never know they have it.

This chapter is your introduction to the problem. The chapters that follow will walk you through exactly what iron does in your body, the full range of symptoms to watch for, how to get the right tests and interpret them honestly, how to supplement effectively, how to eat to build your stores, how to manage heavy periods, how pregnancy affects your iron, and how to navigate the emotional and cognitive effects of deficiency.

You are here because something feels off. Maybe you have been told it is all in your head. Maybe you have been told you are fine. Maybe you have been told to drink more water, sleep more, or stress less.

Maybe it is time to get your iron checked.

Toolkit Summary

- **One in three premenopausal women has low iron stores** — and most do not know it. Heavy periods, pregnancy, diet, and absorption issues are the main drivers of loss.
- **Iron deficiency and iron deficiency anemia are different** — deficiency comes first (low ferritin, normal hemoglobin), anemia comes later. Most women are in the deficiency stage and are not tested.
- **Iron deficiency is a master mimicker** — it is frequently misdiagnosed as depression, anxiety, chronic fatigue, or thyroid disorders. Ask your doctor to check your ferritin, not just your hemoglobin.
- **Risk factors include** heavy periods, postpartum status, vegetarian/vegan diet, athletic training, GI conditions, blood donation, and medications that reduce stomach acid.

- **Undiagnosed iron deficiency has real costs** — lost productivity, reduced quality of life, misdiagnosis, and years of unnecessary suffering. But the condition is treatable, and recovery is possible.
- **Talk to your doctor** — if you have any of the risk factors or symptoms described in this chapter, a simple blood test can tell you whether iron is part of your picture.

ch02

IRON HAS A PR PROBLEM. WHEN MOST PEOPLE HEAR THE WORD, THEY THINK OF CAST-IRON SKILLETS, RUSTY PIPES, AND HEAVY METAL. IT DOES NOT SOUND LIKE SOMETHING YOU WANT INSIDE YOUR BODY. BUT IRON IS NOT JUST SOME MINERAL YOUR DOCTOR MENTIONS IN PASSING. IT IS ONE OF THE MOST ESSENTIAL SUBSTANCES YOUR BODY NEEDS TO FUNCTION, AND IT TOUCHES NEARLY EVERY SYSTEM YOU HAVE — FROM THE AIR YOU BREATHE TO THE THOUGHTS YOU THINK.

Understanding what iron actually does is the key to understanding why low iron makes you feel the way you do. Every symptom of iron deficiency — the fatigue, the brain fog, the hair loss, the cold hands, the racing heart, the shortness of breath — is a direct consequence of your body not having enough iron to do its job. When you know what that job is, the symptoms stop being mysterious. They become logical. They become predictable. And that knowledge is the first step toward fixing them.

Oxygen Transport: The Main Event

This is iron's most famous role, and for good reason. Your body needs oxygen to survive. Every cell in your body uses oxygen to produce energy, and the only way

oxygen gets from your lungs to your cells is through your blood. And the only way your blood carries oxygen is with iron.

Here is how it works. Your red blood cells contain a protein called hemoglobin. Hemoglobin is made of four subunits, and each subunit contains a heme group — a molecular structure with an iron atom at its center. That iron atom is what actually grabs hold of oxygen molecules as they pass through your lungs. One hemoglobin molecule can carry up to four oxygen molecules at a time, each one latched onto an iron atom.

When you breathe in, oxygen enters your lungs and binds to the iron in your hemoglobin. Your heart pumps that oxygen-rich blood through your arteries to every part of your body. When the blood reaches your tissues — your muscles, your brain, your organs — the oxygen releases from the iron and diffuses into the cells that need it. Then your blood, now carrying carbon dioxide, returns to your lungs to pick up more oxygen.

This cycle repeats thousands of times a day, and it depends entirely on having enough iron to build enough hemoglobin to carry enough oxygen.

When your iron stores are low, your body cannot make enough hemoglobin. Your red blood cells become smaller and paler. They carry less oxygen per cell. Your tissues get less oxygen than they need. And because every system in your body depends on oxygen, every system is affected.

The fatigue you feel is not psychological. It is your cells literally not getting enough oxygen to produce energy. The shortness of breath is not anxiety. It is your body trying to compensate for low oxygen by making you breathe faster. The dizziness is not dehydration. It is your brain — the most oxygen-hungry organ in your body — not getting enough fuel.

Energy Production: The Mitochondrial Connection

You have probably heard of mitochondria — they are often called the power plants of your cells. What you may not know is that mitochondria are absolutely dependent on iron.

Inside every mitochondrion, a process called the electron transport chain produces the energy your cells need to function. This energy is stored in a molecule called ATP, and without ATP, your cells cannot do anything — not contract, not divide, not repair themselves, not communicate with each other. Several of the key proteins in the electron transport chain contain iron. Without iron, the chain breaks, and energy production drops.

This is part of why iron deficiency fatigue is so different from ordinary tiredness. Ordinary tiredness is what you feel after a long day or a bad night's sleep. It improves with rest. Iron deficiency fatigue is a cellular energy

crisis. Your mitochondria are literally producing less ATP. Rest does not fix it because the machinery is broken, not overheated.

This is also why women with iron deficiency often describe their energy as a limited resource that runs out faster than it should. You are not lazy. Your cells are running at reduced capacity.

Immune Function

Your immune system relies on iron to fight off infections. Iron is essential for the proliferation and activity of immune cells — particularly lymphocytes and natural killer cells, which are your body's first line of defense against pathogens.

When you are iron deficient, your immune response is impaired. Your body has a harder time fighting off infections, and you may notice that you get sick more often, or that it takes you longer to recover from colds and flus. Some research has also linked iron deficiency to an increased risk of recurrent infections, particularly respiratory and gastrointestinal infections.

There is an interesting evolutionary nuance here, too. Your body deliberately lowers iron levels in the blood during an active infection — this is known as the iron-withholding defense. Many bacteria need iron to grow, so your body pulls iron from circulation to starve the invaders. This is why iron levels can drop during illness and why you should never supplement iron during an

active infection without talking to your doctor first. It is not that your body needs less iron when you are sick — it is that your body is strategically restricting access to keep the infection from getting worse.

Brain Function and Neurotransmitter Synthesis

The brain is the most metabolically active organ in your body. It accounts for only about 2 percent of your body weight but consumes roughly 20 percent of your oxygen. And as we already established, oxygen delivery depends on iron.

But iron's role in the brain goes beyond oxygen transport. Iron is also essential for synthesizing neurotransmitters — the chemical messengers that allow your brain cells to communicate with each other. Dopamine, serotonin, and norepinephrine all require iron-dependent enzymes for their production.

Dopamine is involved in motivation, focus, pleasure, and movement. Serotonin regulates mood, appetite, and sleep. Norepinephrine is involved in alertness and the fight-or-flight response. When your brain does not have enough iron to produce these neurotransmitters, the effects can mimic depression, anxiety, and ADHD — and that is exactly what happens in many women with undiagnosed iron deficiency.

Iron is also required for the production of myelin, the insulating sheath that surrounds your nerve cells and allows signals to travel quickly. Without enough iron, your neural communication slows down. This contributes to the brain fog, slow processing speed, and difficulty finding words that so many iron-deficient women report.

And here is something that might surprise you: iron levels in the brain are tightly regulated, and they do not always reflect iron levels in the rest of your body. Your brain can hold onto its iron stores even when the rest of you is depleted — but eventually, prolonged deficiency catches up. When brain iron drops, it can take significantly longer to replenish than iron in your blood or muscles. This is why cognitive and mood symptoms can persist even after your hemoglobin and ferritin have started to recover.

Temperature Regulation

Have you noticed that you are always cold? That your hands and feet feel like ice even when the rest of you is warm? That you reach for sweaters and blankets while everyone else is comfortable?

This, too, is iron at work.

Iron is involved in thyroid hormone synthesis and metabolism. Thyroid hormones help regulate your body's metabolic rate and temperature. When iron is low, your thyroid may not function optimally, and your body's ability to generate and retain heat is compromised.

But there is a more direct mechanism, too. Oxygen is required for your cells to produce heat as a byproduct of metabolism. Without enough oxygen — because your blood cannot carry enough oxygen — your body produces less heat. Your blood vessels in your hands and feet constrict to conserve heat for your core organs, leaving your extremities cold and sometimes pale or bluish.

Cold intolerance is one of the most common but least recognized symptoms of iron deficiency. If you have been told you just have poor circulation, it is worth checking your ferritin.

Hair and Nail Growth

Your hair and nails are made of keratin, a protein that requires iron-dependent enzymes for its production. When iron is low, your body prioritizes iron for more essential functions — red blood cell production, brain function, immune response — and hair and nail growth get deprioritized.

This is why hair loss is so common in iron deficiency. It is typically a diffuse thinning — hair falling out evenly across your scalp rather than in patches. You may notice more hair in your brush, in the shower drain, on your pillow. Your ponytail may feel thinner. Your part may look wider.

Nail changes can be more subtle. Nails may become brittle, thin, and prone to splitting. They may develop vertical ridges. In more severe cases, nails can become

spoon-shaped (koilonychia) — flattened with raised edges, forming a shallow depression that can hold a drop of water.

These changes are reversible. When your iron stores are restored, your hair and nails will recover. But it takes time — hair growth cycles mean that improvement in hair shedding may not be visible for three to six months after your iron normalizes.

The Iron Cycle: How Your Body Manages Its Supply

Your body does not have an effective way to excrete excess iron. Unlike some vitamins that you can pee out if you take too much, iron is a mineral that your body holds onto tightly and recycles obsessively. Understanding how this cycle works helps explain why deficiency develops so easily and why supplementation has to be careful.

Absorption in the duodenum. Iron from food and supplements is absorbed in the duodenum, the first section of your small intestine. The cells lining your duodenum — enterocytes — have special proteins that pull iron from your digestive tract into your bloodstream. This is the only place where iron enters your body, and it is tightly regulated.

Transport via transferrin. Once iron enters your bloodstream, it binds to a protein called transferrin, which acts like a taxi service. Transferrin carries iron

through your blood to wherever it is needed — your bone marrow to make red blood cells, your muscles to make energy, your brain to make neurotransmitters.

Storage as ferritin. Iron that is not immediately needed is stored inside a protein called ferritin. Ferritin acts like a closet — it tucks iron away safely for future use. Most of your body's iron stores are held in your liver, spleen, and bone marrow. The amount of ferritin in your blood correlates closely with the amount of iron in your stores, which is why a ferritin blood test is the best measure of whether you have enough iron saved up.

Recycling by macrophages. Here is where your body's efficiency shines. When your red blood cells reach the end of their lifespan — about 120 days — they are broken down by macrophages, a type of immune cell. Macrophages recycle the iron from old red blood cells and release it back into your bloodstream, where transferrin can carry it to new red blood cells or to storage. Your body recycles about 90 percent of the iron it needs this way. Only about 10 percent comes from your diet each day.

This recycling system is brilliant, but it has a vulnerability. If you lose blood — through menstruation, childbirth, injury, or gastrointestinal bleeding — you lose the iron inside that blood. And that iron is gone. Your body cannot recycle what it no longer has. You have to replace it through absorption, and absorption is limited.

Heme vs. Non-Heme Iron

Not all iron is created equal, and the difference matters for how you eat and supplement.

Heme iron comes from animal sources — red meat, poultry, fish, and shellfish. It is called heme iron because it is attached to the heme protein found in hemoglobin and myoglobin. Heme iron is absorbed efficiently, and its absorption is less affected by other foods you eat at the same meal. About 25 to 30 percent of heme iron is absorbed.

Non-heme iron comes from plant sources — beans, lentils, spinach, fortified cereals, nuts, seeds — and also from supplements. Non-heme iron is absorbed less efficiently. Depending on what else is in your digestive tract, you may absorb only 1 to 10 percent of the non-heme iron you consume. Its absorption is heavily influenced by other dietary factors — vitamin C can boost it, while tannins, phytates, and calcium can block it.

This difference is why vegetarians and vegans need higher total iron intake than meat-eaters, and why the timing of meals matters. We will cover this in detail in the food chapter.

The Hepcidin Problem: Why You Cannot Just Absorb More When You Are Low

Here is one of the most frustrating things about iron biology: your body does not automatically increase iron absorption when you are deficient. It should — logically, the lower your iron stores, the more iron your body should pull from your food. And to some extent, it does. But the process is regulated by a hormone called hepcidin, and hepcidin often works against you.

Hepcidin is produced by your liver and acts as the master regulator of iron absorption. When hepcidin levels are high, your body blocks iron absorption at the duodenum. When hepcidin levels are low, your body lets more iron in.

Here is the cruel irony: inflammation raises hepcidin. Many women with iron deficiency also have some level of chronic inflammation — from their heavy periods, from gut issues, from the stress of living month after month with a body that is not working properly. And inflammation-driven hepcidin blocks iron absorption even when your stores are critically low.

This is also why every-other-day iron dosing works better than daily dosing. High doses of oral iron cause a spike in hepcidin that lasts about 24 hours. If you take iron every day, hepcidin stays elevated, and your absorption efficiency drops over time. If you take iron every other day, hepcidin has time to fall, and you absorb more total iron across the week.

Understanding how hepcidin works helps you understand why some people can eat iron-rich foods and take iron supplements and still remain deficient. It is not that you are doing something wrong. Your body's regulation system is just complicated.

What Happens When Stores Run Out

When your body detects that iron stores are getting low — low ferritin — it tries to compensate. It increases absorption. It increases recycling efficiency. It reduces the amount of iron it allocates to non-essential functions like hair growth.

But if losses continue to outpace intake, the cascade is predictable.

First, **ferritin drops**. Your iron savings account runs low. You still have enough to function day to day, but your buffer is gone.

Next, **transferrin saturation drops**. Your iron taxi service has fewer passengers. Your blood has less iron circulating to the tissues that need it.

Then, **hemoglobin starts to fall**. Your bone marrow cannot make enough hemoglobin for new red blood cells. Your red blood cells become smaller (microcytic) and paler (hypochromic). This is the beginning of anemia.

Finally, **full anemia develops**. Your hemoglobin drops below the normal range. Your tissues are chronically under-oxygenated. Every system in your body is affected

— your heart beats faster to compensate, your lungs work harder, your brain slows down, your muscles weaken, your immune system falters.

This cascade takes months, sometimes years, to unfold. And at any point along the way, supplementation and dietary changes can reverse it. But the longer you wait, the more iron you need to replace, and the longer recovery takes.

Understanding what iron does in your body changes how you think about your symptoms. They stop being mysterious signs that something is wrong with you. They become predictable consequences of a biological problem — a problem that has a solution.

Your body is full of iron-dependent processes that are running on empty right now. The goal is to fill that tank. And the chapters ahead will show you exactly how.

Toolkit Summary

- **Iron is essential for oxygen transport** — hemoglobin, the protein that carries oxygen in your red blood cells, is built around iron. Low iron means less oxygen reaches your tissues.
- **Iron powers your mitochondria** — your cells' energy factories depend on iron to produce ATP. Without it, energy production drops, and no amount of rest will fix it.

- **Your brain needs iron to think and feel** — iron is required to synthesize dopamine, serotonin, and norepinephrine, and to build myelin. Low iron mimics depression, anxiety, and ADHD.
- **Your body recycles iron efficiently but cannot replace blood loss** — about 90 percent of your iron is recycled from old red blood cells. Menstrual bleeding, childbirth, and other blood losses bypass this recycling and must be replaced through diet or supplements.
- **Heme vs. non-heme matters** — iron from animal sources is absorbed 2-3 times more efficiently than iron from plants. Vegetarians and vegans need to be more intentional about iron intake.
- **Hepcidin regulates absorption** — inflammation raises hepcidin and blocks iron absorption. This is why daily iron dosing is less effective than every-other-day dosing for many people.
- **Iron deficiency is a cascade** — low ferritin comes first, then low transferrin saturation, then falling hemoglobin, then full anemia. The earlier you catch it, the faster recovery will be. Talk to your doctor about testing.

ch03

IF YOU ASK MOST WOMEN WHAT IRON DEFICIENCY FEELS LIKE, THE FIRST WORD OUT OF THEIR MOUTHS WILL BE "TIRED." AND THEY ARE NOT WRONG — FATIGUE IS THE MOST COMMON SYMPTOM AND OFTEN THE MOST DEBILITATING. BUT FATIGUE ALONE DOES NOT CAPTURE THE FULL PICTURE. IRON DEFICIENCY DOES NOT JUST MAKE YOU TIRED. IT CHANGES THE WAY YOU THINK, THE WAY YOU FEEL, THE WAY YOU MOVE THROUGH THE WORLD. IT SHOWS UP IN YOUR HAIRBRUSH, IN YOUR NAILS, IN YOUR HANDS AND FEET, IN YOUR CHEST WHEN YOU CLIMB A FLIGHT OF STAIRS, IN THE MIDDLE OF THE NIGHT WHEN YOUR LEGS WILL NOT STOP MOVING.

Many of these symptoms do not seem connected to iron at first glance. Restless legs? That does not sound like a mineral problem. Ice cravings? That is just a quirk. Tinnitus? That is probably stress. Anxiety? That is a mental health issue.

But all of them trace back to the same root: your body does not have enough iron to do what it needs to do. And when you know the full list, the seemingly random collection of complaints that has been bothering you for years suddenly forms a coherent picture.

The Full Symptom List

Here is what iron deficiency can cause. Not every woman experiences every symptom, and severity varies wildly from person to person. But if you recognize yourself in more than a few of these, it is time to pay attention.

Fatigue that rest does not fix. This is the hallmark. Not the tiredness of a late night, but a deep, cellular exhaustion that does not improve with sleep. You wake up tired. You are tired all day. Coffee does not help. Naps do not help. You are not physically incapable of doing things, but everything costs more energy than it should.

Brain fog. Difficulty concentrating. Trouble finding words mid-sentence. Forgetting why you walked into a room. Reading the same paragraph three times and still not absorbing it. Feeling like your thinking is slow, like there is a layer of cotton between you and clear thought. This is one of the most distressing symptoms for many women, and one of the most commonly dismissed.

Hair loss. Diffuse thinning across your entire scalp. More hair in your brush, on your pillow, in the shower drain. Your ponytail feels thinner. Your part looks wider. Hair loss from iron deficiency is rarely patchy — it is a general reduction in density, and it can be subtle at first. The shedding often starts about three months after your iron drops, because hair growth cycles have a built-in delay.

Pica — particularly ice cravings. This is one of the stranger symptoms, and if you have it, you probably already suspect something is off. Pica is the medical term for craving non-food substances. In iron deficiency, the most common craving is ice — pagophagia. Women with low iron report an almost irresistible urge to chew ice, sometimes consuming trays of it per day. No one knows exactly why this happens, but the leading theory is that chewing ice increases blood flow to the brain, temporarily relieving the cognitive effects of low oxygen. Other pica cravings include clay, dirt, paper, cornstarch, and laundry detergent. If you find yourself craving any of these, mention it to your doctor — it is a strong signal that your iron is low.

Restless legs syndrome. An overwhelming urge to move your legs, especially at night when you are trying to sleep. It can feel like crawling, tingling, aching, or an electric sensation deep in your legs. Movement provides temporary relief, but the sensation returns when you stop. Restless legs syndrome is strongly linked to iron deficiency, particularly low brain iron levels. Iron is needed for dopamine production, and dopamine dysfunction in the brain is a key mechanism in restless legs.

Shortness of breath. Feeling winded from activities that used to be easy — climbing stairs, walking uphill, carrying groceries. Your body is compensating for low oxygen by making you breathe faster and harder. This is

often misattributed to being out of shape, but if your fitness level has not changed and your breathing has, iron could be the culprit.

Dizziness and lightheadedness. Feeling woozy when you stand up too fast (orthostatic hypotension). Your brain is sensitive to changes in oxygen delivery, and when your iron is low, your blood may not be able to supply enough oxygen when you change position.

Pale skin. A general pallor — your skin may look washed out, especially in your face. This is most visible in areas with less pigment: the inside of your lower eyelid (the conjunctiva), your nail beds, your palms, and the lining of your mouth. If your lower eyelid is pale instead of a healthy pink, that is a sign of reduced red blood cells.

Cold hands and feet. Poor circulation to your extremities as your body prioritizes oxygen for your core organs. Your hands and feet may feel cold to the touch even in warm environments. Some women report that their hands and feet go numb or tingle.

Brittle or spoon-shaped nails. Your nails may become thin, brittle, and prone to splitting. They may develop vertical ridges. In more advanced deficiency, nails can become koilonychia — flattened with raised edges, forming a concave shape that can hold a drop of water. This is less common now than it was historically, but it is a classic sign of long-standing iron deficiency.

Heart palpitations. A racing, pounding, or irregular heartbeat. Your heart works harder to pump oxygen-depleted blood to your tissues. You may notice your heart beating fast with minimal exertion, or you may feel it pounding in your chest when you are lying in bed at night.

Anxiety. Feeling on edge, restless, or unreasonably worried. Your body's stress response system is affected by low iron in several ways: reduced oxygen triggers a sympathetic nervous system response, and impaired neurotransmitter production affects mood regulation. Many women report that their anxiety decreases significantly once their iron is corrected.

Depression. Low mood, lack of motivation, loss of interest in things you used to enjoy. The overlap between iron deficiency and depression is significant — and frequently missed. Some research suggests that women with iron deficiency are twice as likely to be diagnosed with depression, and that treating the deficiency can improve depressive symptoms even in women who have not responded to antidepressants.

Tinnitus. Ringing, buzzing, or whooshing sounds in your ears that are not caused by an external source. Reduced oxygen delivery to the inner ear and auditory nerve can cause or worsen tinnitus. Some women report that their tinnitus improves or disappears once their iron is corrected.

Frequent infections. Getting sick more often than you used to, or taking longer to recover from colds, flus, and other illnesses. Iron is essential for immune cell function, and deficiency impairs your body's ability to fight off pathogens.

Poor exercise recovery. Feeling wiped out after workouts that used to feel manageable. Taking longer to bounce back. Your muscles need oxygen to repair themselves after exercise, and with low iron, that repair process is slower and less efficient.

Why Women Normalize These Symptoms

Here is the thing about iron deficiency that makes it so insidious: it comes on slowly. You do not wake up one day dramatically different. You wake up a little more tired than last month. You lose a little more hair. You feel a little more forgetful. The changes are gradual, and gradual changes are easy to dismiss.

You adapt. You drink more coffee. You sit down more often. You stop climbing the stairs two at a time. You stop expecting yourself to have the energy you used to have. You tell yourself you are getting older, or busier, or that this is just what being a woman in her thirties or forties feels like.

And the women around you — your friends, your coworkers, your family — are going through the same thing. So when you mention being tired, they nod in recognition. "Me too," they say. "This is just how it is."

You stop asking whether there is a solution because you stop believing there is a problem. You have normalized feeling bad.

This is the trap that keeps women undiagnosed for years. You do not know that other people do not feel this way because everyone around you is also low on iron, and the collective exhaustion becomes a shared baseline. You lose the ability to distinguish between how you feel and how you could feel.

The "I Thought Everyone Felt This Way" Trap

Every woman I have ever spoken to about iron deficiency has said some version of this sentence. It is the moment of recognition — the shift from "this is normal" to "maybe this is not normal after all."

Maybe it happens when a friend shares her ferritin number and it is three times yours. Maybe it happens when you start supplementing and realize, weeks later, that you have not felt this good in years. Maybe it happens when you read a description of iron deficiency and feel like someone has been following you around with a notebook.

The trap exists because iron deficiency is so common that it has become normalized not just individually, but culturally. One in three women. Among heavy bleeders, one in two. These numbers mean that your social circle,

your workplace, your family — many of the women in those circles are also running on empty. The shared experience of exhaustion becomes invisible because it is everywhere.

But here is the truth that women discover when they correct their deficiency: feeling good is not a luxury. Having energy is not a privilege reserved for people who are less busy or less stressed than you. Clear thinking, steady mood, restful sleep, warm hands, healthy hair, a heart that does not pound when you stand up — these are not aspirational goals. They are the biological defaults of a body that has enough iron.

If you have been living below that default for so long that you cannot remember what it feels like, that is not a personal failure. It is a biological problem, and it has a biological solution.

Symptom Severity and Ferritin Levels

Here is where it gets complicated — and where you need to be patient with yourself and your body.

There is not a universal ferritin number at which every woman develops symptoms. Some women feel terrible at ferritin 40. Others function reasonably well at ferritin 20 and only crash when they drop to single digits. Genetics, overall health, activity level, age, and individual biochemistry all play a role in how sensitive you are to low iron.

That said, there are general patterns that can help you understand where you might fall on the spectrum.

Ferritin above 100 ng/mL. Generally considered optimal for most women. You likely have adequate iron stores. Symptoms from iron are unlikely at this level.

Ferritin 70 to 100 ng/mL. Good stores. Most women feel well in this range. Some experts suggest this as the target range for symptomatic women after supplementation.

Ferritin 50 to 70 ng/mL. Suboptimal. You probably have some iron reserves, but they are not generous. You may not have obvious symptoms, or you may have mild symptoms that you have normalized. Some women, particularly athletes or those with heavy periods, start experiencing symptoms in this range.

Ferritin 30 to 50 ng/mL. Low. Your stores are limited, and you may be experiencing symptoms. Many experts now consider ferritin under 50 to be a potential contributor to symptoms in women, especially when combined with heavy menstrual bleeding or athletic training.

Ferritin under 30 ng/mL. Absolute iron deficiency. Your iron stores are depleted. At this level, symptoms are common, and they can be significant — fatigue, brain fog, hair loss, restless legs, cold extremities, shortness of breath, mood changes. Even if your hemoglobin is still normal, you are iron deficient and you need to address it.

Ferritin under 12 to 15 ng/mL. Severe deficiency. You are almost certainly symptomatic. You may be bordering on or already in iron deficiency anemia. Your energy, cognition, and immune function are significantly compromised. You need aggressive treatment — talk to your doctor about the best approach for your situation.

The important takeaway: ferritin below 30 is not normal, even if your lab report says it is. And ferritin below 50 may be causing symptoms for you, especially if you exercise, have heavy periods, or have other risk factors. Your symptoms matter more than the number. If you feel bad and your ferritin is below 50, your symptoms deserve attention.

The Symptom Journal: Your Personal Evidence File

One of the most frustrating aspects of undiagnosed iron deficiency is that the symptoms are invisible to everyone but you. You cannot show someone your fatigue or your brain fog. You cannot hand them a sample of your cognitive slowness. When a doctor asks you how you feel, you have to translate a complex, months-long experience into a few sentences, and in translation, the severity is often lost.

A symptom journal changes that. It gives you a written record of what you have been experiencing — not a vague recollection, but day-by-day, week-by-week evidence. It

helps you identify patterns. It gives you concrete information to bring to your doctor. And it gives you a way to track your progress once you start treatment.

Here is how to start one.

Choose your format. A simple notebook, a spreadsheet, or a notes app on your phone — whatever you will actually use. Consistency matters more than elegance.

Track the basics each day. Date. How many hours you slept. How you felt when you woke up. A brief energy rating on a scale of 1 to 10.

Track specific symptoms. Which ones are bothering you today? Use the list from this chapter as a starting point. Note severity (mild, moderate, severe) and whether the symptom is better or worse than yesterday.

Track your cycle. If you menstruate, note where you are in your cycle. Iron symptoms often worsen during and immediately after your period, and having this documented helps you and your doctor understand the pattern.

Note anything that affects your iron. What did you eat today? Did you take any supplements? Did you drink coffee or tea with meals? Are you taking any medications that affect absorption — acid reducers, antacids, oral contraceptives? This information becomes extremely useful when you start working on treatment.

Bring it to your doctor. A symptom journal is not whining. It is data. It is the difference between sitting in a doctor's office and saying "I think I might be tired" versus handing them a month of daily records showing that your energy levels average 3 out of 10, that your hair has been falling out for weeks, that you have been craving ice every afternoon, and that all of this gets worse the week after your period. Data gets taken seriously.

You do not have to keep the journal forever. But keeping one for a few weeks, especially before your first doctor's appointment about iron, can make the difference between being dismissed and being heard.

The Bigger Picture

Reading this list of symptoms may feel overwhelming. You may recognize yourself in more of them than you expected, and that recognition can bring up a lot of emotions — relief that there is a name for what you are experiencing, sadness for how long you have been struggling, and maybe some anger that no one connected the dots sooner.

All of those feelings are valid. Let yourself feel them. And then let them fuel your next step, which is testing — the subject of the next chapter.

Because symptoms alone can point you in the right direction, but only blood work can confirm what is happening. And when you know what is happening, you can fix it.

Toolkit Summary

- **Iron deficiency causes a wide range of symptoms** beyond fatigue — brain fog, hair loss, ice cravings, restless legs, shortness of breath, dizziness, pale skin, cold hands and feet, brittle nails, heart palpitations, anxiety, depression, and tinnitus. If you have several of these, iron could be the common thread.
- **Women normalize these symptoms because they develop gradually** and because iron deficiency is so common that many women share the same experience. "I thought everyone felt this way" is the most common refrain among women who recover.
- **Ferritin levels correlate with symptoms, but the relationship is individual.** Ferritin under 30 is absolute deficiency. Ferritin under 50 can cause symptoms in many women. Ferritin over 70 is generally considered optimal.
- **A symptom journal creates objective data** you can bring to your doctor. Track energy, specific symptoms, your menstrual cycle, and diet. A few weeks of data is more persuasive than a verbal description.
- **Your symptoms are real, and they have a biological cause.** You are not lazy, not making it up, and not weak. Iron deficiency is a treatable condition. Talk to your doctor about testing.

ch04

YOU HAVE RECOGNIZED YOURSELF IN THE SYMPTOMS. YOU SUSPECT IRON DEFICIENCY MIGHT BE THE MISSING PIECE OF A PUZZLE YOU HAVE BEEN TRYING TO SOLVE FOR MONTHS OR YEARS. NOW YOU NEED THE DATA — AND THAT MEANS GETTING THE RIGHT BLOOD TESTS.

This should be straightforward. It is not. Iron testing is one of the most confusing and frustrating parts of the journey, not because the tests themselves are complicated, but because how they are interpreted — and how they are offered — can vary wildly depending on your doctor, your lab, and what your insurance covers.

Some women walk into a doctor's office, ask for a simple iron test, and walk out with a complete picture of their iron status. Others ask for the same test and receive only a hemoglobin result — which tells them nothing about their stores. Others are told their iron is "normal" based on a ferritin of 12, because the lab's reference range starts at 15.

This chapter is going to arm you with everything you need to navigate this process. You will learn what each test measures, what optimal versus normal actually means, why the reference ranges are so problematic, how to ask your doctor for the tests you need, and what to do if you get a result that does not match how you feel.

The Complete Iron Panel

A complete iron panel includes several tests, each giving you a different piece of the puzzle. Your doctor may not order all of them at once, but knowing what each one measures helps you understand your results.

Ferritin. This is the single most important test for diagnosing iron deficiency. Ferritin measures your body's stored iron — the iron you have saved for future use. It is the closest thing you have to a fuel gauge for your iron reserves. Low ferritin is the earliest sign of iron depletion, appearing long before your hemoglobin drops. If you can only get one test, make it ferritin.

Hemoglobin. This measures the amount of hemoglobin protein in your red blood cells. Hemoglobin is what carries oxygen, and its levels directly reflect your blood's oxygen-carrying capacity. Low hemoglobin means anemia. But — and this is critical — hemoglobin can be perfectly normal even when your ferritin is critically low. Hemoglobin is a late-stage indicator. By the time hemoglobin drops, you have been iron deficient for a long time.

Hematocrit. This measures the percentage of your blood volume that is made up of red blood cells. It is closely related to hemoglobin — when hemoglobin is low, hematocrit is usually low too. Like hemoglobin, hematocrit is a measure of anemia, not of iron stores.

Serum iron. This measures the amount of iron circulating in your blood right now, at this moment. It fluctuates throughout the day and is affected by what you have eaten recently. It is useful as part of the full picture but unreliable on its own — a single serum iron level can be normal in an iron-deficient person depending on when the blood was drawn.

Total iron-binding capacity (TIBC). This measures how much iron your blood could carry if it were fully saturated. Transferrin is the protein that carries iron in your blood, and TIBC is an indirect measure of transferrin levels. When your body is low on iron, it produces more transferrin to try to capture more iron from your diet. So high TIBC is a sign of iron deficiency. Low TIBC can indicate iron overload or chronic inflammation.

Transferrin saturation. This is calculated from your serum iron and your TIBC — what percentage of your iron-carrying capacity is actually being used. Low transferrin saturation (below 15 to 20 percent) suggests that not enough iron is available for your body's needs. This is a more sensitive indicator of iron deficiency than serum iron alone, because it accounts for how much transport capacity your body has created.

RDW (red cell distribution width). This measures the variation in size of your red blood cells. In iron deficiency, your body produces a mix of normal-sized red blood cells and smaller-than-normal cells, which increases

the RDW. A high RDW is often one of the earliest signs of iron deficiency on a standard complete blood count — sometimes showing up before anemia develops.

What Optimal vs. Normal Actually Means

Here is the most important thing to understand about iron lab results: the ranges printed on your lab report are not designed to identify optimal health. They are designed to flag severe disease. And for ferritin, that distinction matters enormously.

Most labs in the United States use a ferritin reference range that starts somewhere between 10 and 30 ng/mL and ends somewhere between 150 and 300 ng/mL. A ferritin of 12 is "normal" by some lab standards. A ferritin of 15 is "normal" by many. A ferritin of 20 is comfortably "normal" on most lab reports.

But "normal" is not the same as "optimal." And in the case of ferritin, the gap between the two is wide enough to drive a truck through.

Leading experts in women's health and iron metabolism have been arguing for years that the reference ranges need to be updated. The World Health Organization defines ferritin under 15 as iron deficiency in adults. Many hematologists consider ferritin under 30 to be absolute iron deficiency. Some researchers now argue that ferritin under 50 can impair physical performance and cognitive function in women, particularly athletes and those with heavy menstrual bleeding.

Here is how to interpret your ferritin in a way that reflects what your body actually needs:

Ferritin under 30 ng/mL. This is iron deficiency, period. Regardless of what your lab's reference range says, a ferritin under 30 means your iron stores are depleted. You need to address this. If your doctor tells you your iron is fine with a ferritin under 30, they are using an outdated or overly permissive standard.

Ferritin 30 to 50 ng/mL. This is borderline low. Many women experience symptoms in this range, especially if they exercise, have heavy periods, or have other iron losses. You may not need aggressive supplementation, but you should be aware that your stores are limited and work on improving them through diet and targeted supplementation.

Ferritin 50 to 70 ng/mL. Suboptimal but not alarming. Some women feel fine at this level. Others — particularly athletes or women with ongoing blood loss — may still experience symptoms. If you are symptomatic, it is worth working toward a higher level.

Ferritin 70 to 100+ ng/mL. Generally considered optimal for menstruating women. At this level, most women have adequate stores to function well, handle monthly blood loss, and support their hair, skin, and cognitive function. This is the range you are aiming for with treatment.

The same principle applies to transferrin saturation. Optimal transferrin saturation is typically between 20 and 35 percent. Below 15 to 20 percent is a sign that iron availability is limited, even if your ferritin looks okay.

The Normal-Range Controversy

Why are the lab ranges set so low? The answer is complicated, but it comes down to a combination of outdated data, population-based statistics, and a lack of sex-specific standards.

Lab reference ranges are typically calculated by testing a large population of healthy people and drawing cutoffs at the 2.5th and 97.5th percentiles. Everyone who falls within the middle 95 percent of that population is considered "normal." The problem is that the reference population for many iron studies includes people who are themselves iron deficient.

Think about that for a moment. One in three premenopausal women has low iron stores. If you include those women in your reference population, the 2.5th percentile for ferritin is going to include people who are genuinely deficient. The cutoff ends up being set at a level that is "normal" only in the statistical sense of being common — not in the medical sense of being adequate.

This is not a conspiracy. It is a methodological problem that has been slow to change because updating reference ranges requires large-scale studies, consensus among professional organizations, and buy-in from commercial

labs. In the meantime, millions of women are walking around with ferritin levels that fall within the "normal" range but are far too low to support optimal health.

Different organizations use different cutoffs for iron deficiency. The World Health Organization uses ferritin below 15 for adults. Many hematologists use below 30. The American Society of Hematology acknowledges that ferritin under 30 is suggestive of iron deficiency, but does not have a universal standard. The CDC has published data showing that substantial percentages of women have ferritin levels below 20. And leading researchers in the field have called for raising the cutoff for iron deficiency in women to 30 or even 50 ng/mL.

The bottom line: if your ferritin is under 30, you are iron deficient. If your ferritin is under 50 and you have symptoms, you may be iron deficient in a way that matters for your quality of life. Do not let a "normal" label on your lab report convince you otherwise.

How to Ask Your Doctor for the Right Tests

Knowing what to ask for is half the battle. Here is how to have that conversation with your doctor.

Be specific about what you want. Do not say "Can you check my iron?" because that can mean different things to different providers. Say: "I would like a complete iron panel including ferritin, hemoglobin,

hematocrit, serum iron, TIBC, and transferrin saturation." If you are concerned about anemia, you can also ask for a complete blood count (CBC) with RDW.

Explain why. Help your doctor understand why you are asking. You can say something like: "I have been experiencing persistent fatigue, brain fog, hair loss, and ice cravings. I have read that these can be symptoms of iron deficiency, and I would like to be tested. I know that ferritin is the most sensitive test for iron stores, and I would like that included."

Bring your symptom journal. If you have been tracking your symptoms, bring your journal. Hand your doctor a month of daily data showing low energy, specific symptoms, and their relationship to your menstrual cycle. Data speaks louder than vague complaints.

If your doctor says no, ask why. Some doctors may resist testing because they do not think it is necessary, because your hemoglobin was normal on a recent CBC, or because they follow guidelines that only recommend testing in certain populations. If this happens, ask: "I understand your reasoning, but I would like to rule this out because my symptoms are affecting my quality of life. Could we at least check my ferritin?" A single ferritin test is inexpensive and low-risk. Most doctors will agree if you present it clearly.

Consider a direct-to-consumer lab. If you are unable to get your doctor to order the tests, or if you want to do your own initial investigation, you can order your own lab work through companies like Walk-In Lab, Ulta Lab Tests,

Request A Test, or Jason Health. You pay out of pocket, and the cost for an iron panel plus ferritin is typically between \$30 and \$80. You take the lab requisition to a local blood draw center, and you get your results online. This is a legitimate option for women who are struggling to be heard. Bring those results to your doctor for interpretation and follow-up.

What to Do If Your Doctor Says "Your Iron Is Fine"

This is one of the most disheartening moments in the iron deficiency journey. You have gathered your courage, made the appointment, described your symptoms carefully, and waited for the results. And then the message comes back: "Your iron is fine. All your labs are normal."

But you do not feel fine. And you need to know whether "normal" means normal by the right standard.

Here is how to navigate this situation.

Ask for the actual numbers. Do not accept "everything looks normal" as a full answer. Ask for a copy of your lab results. You are entitled to them under HIPAA, and most patient portals make them easily accessible. Look at the actual ferritin number. If your ferritin is under 30 ng/mL, your iron is not fine — and you can say so.

Ask which ranges the lab uses. Different labs use different reference ranges. A ferritin of 18 may be flagged as low at one lab and normal at another. Ask your doctor which cutoff your lab uses for ferritin and whether they are aware that many experts consider ferritin under 30 to be iron deficiency.

Bring information to the conversation. It is not your job to educate your doctor, but sometimes a gentle nudge helps. You can say: "I understand that my ferritin of 18 is within the lab's reference range, but I have done a lot of reading, and I know that ferritin under 30 is considered absolute iron deficiency by many experts. I would like to discuss how to improve my iron stores, because my symptoms are significantly affecting my quality of life."

Ask about a different approach. If your ferritin is low but your hemoglobin is normal, ask about trying oral iron supplementation for a few months to see if it helps your symptoms. A trial of iron is low-risk when done under medical supervision, and many doctors will agree to a monitored trial even if they are not convinced you are deficient by the numbers alone.

Consider a second opinion. If your doctor is unwilling to engage with the evidence or to explore the possibility that your symptoms are related to low iron, it may be time for a second opinion. A functional medicine practitioner, a hematologist, or a women's health

specialist may take a more nuanced view of ferritin than a general practitioner who is following population-level guidelines.

Use the lab tracker. Below, you will find a lab tracker that helps you organize your results over time. Having multiple data points — your ferritin this month versus three months ago versus before your period — can reveal trends that a single lab draw cannot.

The Lab Tracker: Your Personal Iron Record

One of the most valuable tools in your iron journey is a simple tracker that records your lab results over time. Iron status changes with your cycle, with supplementation, with pregnancy, and with changes in your health. A single ferritin number from one day is useful. A year of ferritin numbers is powerful.

Here is what to track each time you are tested:

- Date of blood draw
- Where you were in your menstrual cycle (day of cycle, or phase if you do not track days)
- Ferritin
- Hemoglobin
- Hematocrit
- Serum iron
- TIBC

- Transferrin saturation
- RDW
- Any supplements you were taking (type, dose, frequency)
- Any symptoms you were experiencing (brief notes)

You can track this in a spreadsheet, a notebook, or a notes app. The important thing is to have a record you can refer to over time. When you see your ferritin moving from 12 to 25 to 45 over the course of six months of supplementation, you have objective evidence that the treatment is working — even before you feel the full benefit in your symptoms.

And when you switch doctors or see a specialist, having a timeline of your results saves you from having to explain your entire history from scratch.

The Bottom Line

Testing is your gateway to treatment. Without the right tests, you are guessing. With the right tests interpreted correctly, you have a map.

The most important test is ferritin. The most important number is not whether your ferritin falls within the reference range — it is whether your ferritin is high enough to support your body's needs. For most women, that means a ferritin of 70 or above.

If your ferritin is under 30, you are iron deficient, regardless of what your lab report says. If your ferritin is under 50 and you have symptoms, your deficiency may be contributing to how you feel. Trust your body over the reference range.

And if you have already been tested and your doctor said everything is fine, but you are still struggling — go back and look at the actual numbers. You might find that "fine" means something very different than you thought.

Toolkit Summary

- **Ferritin is the most important test** — it measures your stored iron. Ferritin under 30 is absolute iron deficiency, even if your lab report calls it normal. Ferritin under 50 can cause symptoms in many women.
- **Hemoglobin only detects anemia, not deficiency** — normal hemoglobin does not mean normal iron stores. You can be iron deficient with normal hemoglobin for months or years before anemia develops.
- **A complete iron panel includes** ferritin, hemoglobin, hematocrit, serum iron, TIBC, transferrin saturation, and RDW. If you can only get one test, make it ferritin.

- **The "normal range" for ferritin is set too low** — lab ranges are calculated from populations that include iron-deficient women. Do not accept "your iron is normal" without seeing the actual numbers.
- **Ask your doctor specifically for the tests you need** — bring your symptom journal, be clear about what you want, and do not be afraid to push back or seek a second opinion.
- **Track your results over time** — a lab tracker gives you a record of your progress and helps you identify trends that a single test cannot reveal. Talk to your doctor about getting tested.

ch05

YOU HAVE YOUR TEST RESULTS. YOUR FERRITIN IS LOW. YOUR DOCTOR HAS AGREED THAT SUPPLEMENTATION MAKES SENSE. NOW COMES THE PRACTICAL QUESTION: WHAT DO YOU ACTUALLY TAKE, AND HOW DO YOU TAKE IT IN A WAY THAT WORKS WITH YOUR BODY INSTEAD OF AGAINST IT?

Iron supplementation is not as simple as buying a bottle and swallowing a pill. Different forms of iron have different absorption rates, different side effect profiles, and different effectiveness for different people. The dose matters. The timing matters. What you eat with it matters. And for some women, oral supplements are not enough — they need intravenous iron to get their stores back where they need to be.

This chapter covers all of it. You will learn about the different types of oral iron, how to dose and time them for maximum absorption and minimum side effects, how long recovery takes, when to consider IV iron, and how to talk to your doctor about the approach that is right for you.

Types of Oral Iron

Not all iron supplements are created equal. The form of iron — and the amount of elemental iron it delivers — makes a significant difference in how well you tolerate it and how much you absorb.

Ferrous sulfate is the most common and most widely studied form of oral iron. It is inexpensive, readily available, and effective. Most over-the-counter iron supplements and many prescriptions are ferrous sulfate. The standard dose is 325 mg, which provides about 65 mg of elemental iron. The downside: ferrous sulfate is the most likely to cause gastrointestinal side effects — nausea, constipation, stomach cramping, and that metallic aftertaste. If you have a sensitive stomach, this form may be hard to tolerate.

Ferrous fumarate is another common form, similar to ferrous sulfate in absorption and side effects. It contains a higher percentage of elemental iron per milligram — about 33 percent — which means a smaller pill can deliver the same amount of iron. A standard 200 mg tablet of ferrous fumarate provides about 65 mg of elemental iron. The side effect profile is similar to ferrous sulfate.

Ferrous bisglycinate is a chelated form of iron, meaning the iron is bound to the amino acid glycine. This form is generally better tolerated than ferrous sulfate — less constipation, less nausea, less stomach upset. The iron is absorbed through a different pathway in the digestive tract, which may reduce the gastrointestinal distress that many women experience with standard iron pills. Ferrous bisglycinate typically contains about 20 percent elemental iron. Some studies suggest it is absorbed as well as or better than ferrous sulfate when taken at equivalent doses of elemental iron, but the

evidence is mixed. Many women who cannot tolerate ferrous sulfate find ferrous bisglycinate much more comfortable.

Heme iron polypeptide is a different category entirely. It is derived from animal hemoglobin and contains iron in its heme form — the same form found in meat. Heme iron is absorbed through a separate pathway from non-heme iron, and it is not affected by many of the dietary inhibitors that block non-heme iron absorption. It is also much less likely to cause gastrointestinal side effects. The downsides: heme iron supplements are more expensive than ferrous sulfate, they are harder to find, and they are not always covered by insurance. For women who cannot tolerate other forms of oral iron, heme iron polypeptide can be a game-changer.

Slow-release or extended-release iron is designed to release iron gradually through the digestive tract, theoretically reducing side effects. In practice, slow-release iron formulations often release the iron past the duodenum — the only place where iron is efficiently absorbed — which means you absorb less of the dose. They are generally not recommended for treating iron deficiency. Standard immediate-release formulations are preferred.

What Works Best with Fewest Side Effects

There is no single "best" iron supplement for everyone. The best supplement is the one you can tolerate consistently enough to actually take for the weeks or months needed to rebuild your stores.

That said, here is a general framework based on the evidence and the experiences of thousands of women:

If you have no history of GI sensitivity, **ferrous sulfate** is a perfectly reasonable starting point. It is the most studied form, it is cheap, and it works. Take it with a small amount of food if needed — but not with calcium-rich foods, tea, or coffee.

If you have a sensitive stomach or you tried ferrous sulfate and could not tolerate it, **ferrous bisglycinate** is usually the next step. It is gentler and produces fewer side effects for most women. It is slightly more expensive but well worth it if it means you can take it consistently.

If you have significant GI sensitivity or you have tried multiple forms and none work, **heme iron polypeptide** is worth considering. It is the most expensive option and requires a prescription in some countries, but it is extremely well tolerated and bypasses many of the barriers that make non-heme iron absorption difficult.

If you cannot tolerate any oral iron, or if oral iron is not raising your ferritin despite consistent use, **IV iron** may be the right option — more on that below.

Dosing and Timing: The Science of Taking Iron Well

How you take iron matters as much as what form you take. Here are the key principles.

Every other day is better than daily. Remember hepcidin from the second chapter? High-dose oral iron triggers a spike in hepcidin that lasts about 24 hours and blocks further iron absorption during that window. When you take iron every day, hepcidin stays elevated, and your absorption efficiency drops over time. When you take iron every other day — even at the same total weekly dose — hepcidin has time to fall between doses, and you absorb more of the iron you take. A landmark study published in *The Lancet* found that every-other-day iron dosing resulted in similar or better absorption than daily dosing, with fewer side effects. If you are taking moderate to high doses of oral iron, try every other day.

Take with vitamin C. Vitamin C (ascorbic acid) significantly enhances non-heme iron absorption. It works by keeping iron in a form that is easier for your body to absorb and by helping release iron from food components that would otherwise block it. Taking your iron supplement with a glass of orange juice, a vitamin C supplement, or a meal that includes vitamin C-rich foods can double or triple absorption. Aim for at least 250 mg of vitamin C with your iron dose.

Avoid absorption inhibitors. For at least one hour before and two hours after taking iron, avoid coffee, tea (black, green, and herbal varieties that contain tannins), calcium supplements, calcium-rich foods (dairy, fortified plant milks, yogurt), and high-fiber foods. These all interfere with iron absorption. If you must have your morning coffee, take your iron at a different time of day.

Take on an empty stomach if possible. Iron is best absorbed on an empty stomach — ideally at least one hour before a meal or two hours after. This is often the hardest rule to follow because iron on an empty stomach can cause nausea, especially with ferrous sulfate. If you need to take it with food, keep the meal small and avoid the inhibitors listed above. A small amount of food that includes vitamin C (like a few strawberries or a glass of orange juice) is ideal.

Consider the time of day. Many women prefer to take iron at bedtime to avoid the interaction with meals and the potential for nausea during the day. This works well, but be aware that iron can cause stomach upset in some people even at night. Start with a lower dose and see how you tolerate it before committing to a specific schedule.

Managing Side Effects

Iron supplements are famous for their side effects, and if you have had a bad experience with them before, you are not alone. But most side effects can be managed with the right approach.

Constipation is the most common complaint. Iron slows bowel movements, and for some women, it can cause significant discomfort. To manage it: increase your water intake, add a fiber supplement like psyllium husk (taken at least two hours away from your iron dose), increase your intake of magnesium-rich foods or consider a magnesium supplement, and consider switching to ferrous bisglycinate if constipation is severe.

Nausea and stomach cramping are usually related to taking iron on an empty stomach or taking too high a dose too quickly. Try taking iron with a very small amount of food (a cracker or two, a few bites of fruit) and gradually increase the dose. If nausea persists, switch to a gentler form like ferrous bisglycinate.

Black stool is normal. It is not a sign that something is wrong. The unabsorbed iron passes through your digestive tract and turns your stool dark green or black. This is harmless and expected. The only time you should be concerned is if your stool is black and tarry — like black road tar — or accompanied by abdominal pain, because that can indicate gastrointestinal bleeding. Normal iron-related black stool is a uniform dark color without a sticky, tarry consistency.

Metallic taste is common with liquid iron supplements and some tablets. If you experience it, try a different form (bisglycinate is less likely to cause this), rinse your mouth after taking the supplement, or take it with a small amount of citrus.

Diarrhea is less common than constipation but can occur, particularly with high doses or liquid iron. If it happens, reduce your dose and consider switching to a different form.

If side effects are severe enough that you are skipping doses or dreading taking your supplement, talk to your doctor about a different form of iron or a different dosing strategy. You should not have to suffer to treat a condition that treatment is supposed to help.

How Long Until You Feel Better

This is the question every woman wants answered, and the honest answer is: it depends. For many women, symptom improvement begins within two weeks — a lightening of the fatigue, clearer thoughts, less afternoon slump. But feeling better is not the same as being better. Hair shedding often decreases around three to four months (hair cycles are slow, and the hair already in the shedding phase must complete its cycle first). Full restoration of iron stores takes three to six months of consistent supplementation, sometimes longer if your deficiency is severe or you have ongoing blood loss. Your ferritin level is the best measure: a typical response to oral iron is a ferritin increase of about 5-10 ng/mL per month, but this varies widely based on dose, form, absorption, and ongoing losses. If you are not seeing

improvement in ferritin after three months of consistent supplementation, talk to your doctor — you may need a higher dose, a different form, or IV iron.

IV Iron: Who Needs It and What to Expect

For some women, oral iron is not enough. This can happen when:

- Your iron deficiency is severe (ferritin under 10, hemoglobin critically low)
- You cannot tolerate oral iron due to side effects
- You have a condition that impairs absorption (celiac disease, Crohn's, ulcerative colitis, gastric bypass)
- You have ongoing heavy blood loss that outstrips what oral iron can replace
- You have been taking oral iron consistently for months with little to no improvement

In these situations, intravenous (IV) iron can be a faster and more effective option. IV iron delivers iron directly into your bloodstream, bypassing the digestive tract entirely. It is not absorbed — it is already in your system the moment it is infused. This means no GI side effects and no absorption issues.

There are several types of IV iron, and they differ primarily in how much iron they deliver per infusion and how quickly they can be administered.

Iron dextran (INFeD, Dexferrum) is an older formulation that requires a test dose because of a small risk of allergic reaction. It can be given in large doses, meaning fewer infusions, but the risk of reaction makes it less commonly used today.

Ferric gluconate (Ferrlecit) is a newer formulation with a better safety profile. It is typically given in smaller doses over multiple sessions. It is commonly used for people with chronic kidney disease or other conditions requiring regular IV iron.

Iron sucrose (Venofer) is similar to ferric gluconate — small doses, multiple sessions, good safety profile. It is also commonly used in kidney disease and in pregnancy.

Ferumoxytol (Feraheme) can be infused more quickly than older formulations — sometimes in as little as 15 minutes. It delivers a larger dose per infusion, so you may need only one or two treatments. It has a good safety profile, though there is a small risk of allergic reaction.

Ferric carboxymaltose (Injectafer) is one of the most commonly used IV iron formulations today. It delivers a high dose — up to 750 mg per infusion, sometimes 1,000 mg or more depending on your weight and deficiency — in about 15 to 30 minutes. Most people need one or two infusions total. It is well tolerated and has a low risk of serious side effects.

Iron isomaltoside (Monoferric) is similar to ferric carboxymaltose — high-dose, short infusion time, good safety profile. It can be given in doses up to 1,000 mg or more per infusion.

Here is what to expect with IV iron. First, you will have a consultation with your doctor to determine whether IV iron is appropriate. If it is, you will be scheduled for an infusion, usually at a hospital outpatient clinic, an infusion center, or your doctor's office. A nurse or doctor will insert an IV line and administer the iron over 15 minutes to an hour, depending on the formulation. You will be monitored for a short time afterward to make sure you do not have a reaction. Most women tolerate IV iron very well.

After the infusion, you may experience some mild side effects: headaches, muscle aches, a metallic taste in your mouth, and — this is common — a temporary drop in blood pressure that can make you feel lightheaded. These usually resolve within a day or two. Serious allergic reactions are rare with modern formulations.

The benefits of IV iron can be dramatic. Many women report feeling significantly better within days to a week after their infusion — much faster than with oral iron. Your ferritin will rise sharply, and your body will have the iron it needs to function properly almost immediately.

Important note on infection: IV iron can temporarily suppress your immune system's ability to fight certain infections, because it overwhelms the body's iron-withholding defense system. If you have an active

infection, most doctors will delay IV iron until the infection is resolved. Always tell your doctor about any signs of infection before your infusion.

The Iron Supplement Comparison Table

Here is a quick-reference guide to the most common oral iron options. Share this with your doctor to help decide what is right for you.

Ferrous sulfate — 325 mg tablet provides 65 mg elemental iron. Absorption: moderate. GI side effects: common (constipation, nausea, cramping). Cost: very low. Available: over the counter.

Ferrous fumarate — 200 mg tablet provides 65 mg elemental iron. Absorption: moderate. GI side effects: common. Cost: low. Available: over the counter.

Ferrous bisglycinate — Varies by brand; check label for elemental iron. Typically about 20% elemental iron. Absorption: moderate to good. GI side effects: mild to none for most people. Cost: moderate. Available: over the counter.

Heme iron polypeptide (Proferrin, OptiFer) — 11-12 mg elemental iron per capsule. Absorption: high. GI side effects: minimal. Cost: higher. Available: over the counter or by prescription depending on country.

Talking to Your Doctor About IV Iron

If you think IV iron might be right for you, here is how to bring it up with your doctor.

Start with your experience: "I have been taking oral iron for [X months] at [dose], and my ferritin has only gone from [X] to [Y]. I am still experiencing symptoms, and I would like to explore IV iron as an option."

If you cannot tolerate oral iron: "I have tried ferrous sulfate, ferrous fumarate, and ferrous bisglycinate, and I cannot tolerate any of them at an effective dose. I need to discuss alternatives including IV iron."

If your deficiency is severe: "My ferritin is [X] and I am symptomatic. I would like to discuss whether IV iron could help me recover more quickly."

Some doctors are more open to IV iron than others. If your doctor is hesitant, ask what threshold they use for considering IV iron — some follow guidelines that only recommend it for women with hemoglobin below a certain level, which may not capture your situation if you are iron deficient without anemia. A referral to a hematologist can be helpful in these cases.

A Word of Caution

Iron is not a vitamin. It is a mineral that your body stores and accumulates. Unlike water-soluble vitamins that you excrete when you take too much, excess iron can build up in your tissues and cause damage over time. This

is why you should never supplement iron without knowing your ferritin level, and why you should stop supplementing when your ferritin reaches a healthy range.

Always work with your doctor on your supplementation plan. The iron supplement comparison table and the dosing guidance in this chapter are starting points for a conversation, not prescriptions. Your doctor can help you determine the right dose, form, and duration for your specific situation — and monitor your progress to make sure you are getting the benefits without the risks.

Toolkit Summary

- **Different forms of oral iron have different side effect profiles** — ferrous sulfate is the cheapest but hardest on the stomach; ferrous bisglycinate is gentler; heme iron polypeptide is the gentlest but most expensive. The best supplement is the one you can take consistently.
- **Every-other-day dosing is often more effective than daily** — hepcidin spikes after a dose and blocks absorption for about 24 hours. Every-other-day dosing lets hepcidin fall and improves overall absorption.
- **Take iron with vitamin C and away from coffee, tea, calcium, and high-fiber foods** — vitamin C boosts absorption; the others block it. At least one hour before and two hours after.

- **Symptoms often improve in 2 weeks; hair shedding may decrease in 3-4 months; full stores take 3-6 months** — patience is essential. Track your ferritin every 2-3 months to monitor progress.
- **IV iron is an option when oral iron fails** — due to intolerance, absorption issues, severe deficiency, or ongoing blood loss that oral iron cannot keep up with. Modern formulations are safe and effective. Talk to your doctor about whether IV iron is right for you.
- **Never supplement iron without knowing your ferritin** — iron overload is a real risk. Work with your doctor to determine the right dose and duration for your situation.

ch06

THERE IS A QUIET FRUSTRATION THAT COMES FROM EATING A HEALTHY DIET AND STILL BEING IRON DEFICIENT. YOU ARE EATING YOUR SPINACH. YOU ARE COOKING WITH LENTILS. YOU STARTED YOUR DAY WITH FORTIFIED CEREAL. AND YET YOUR FERRITIN IS SITTING AT 18 AND YOUR HAIR IS STILL FALLING OUT. WHAT GIVES?

The answer is that food iron is complicated. The amount of iron in a food is only half the story. The other half — the half that determines whether your body actually uses that iron — is about absorption, about what else you eat with it, and about the intricate dance of enhancers and inhibitors that plays out in your digestive tract with every meal.

This chapter is going to teach you how to eat for your iron stores — not by obsessing over every bite, but by understanding the principles that make your food work for you rather than against you.

Heme Iron Sources: The Gold Standard

Heme iron — the type found in animal foods — is absorbed roughly two to three times more efficiently than non-heme iron from plants. About 25 to 30 percent of the heme iron you eat makes it into your bloodstream,

compared to roughly 1 to 10 percent of non-heme iron. This difference is significant enough that it shapes the entire conversation about iron in food.

Red meat is the most concentrated source of heme iron. Beef, lamb, and bison are all excellent. A three-ounce serving of beef sirloin provides about 2 to 3 milligrams of iron. Grass-fed and conventionally raised beef have similar iron content. The cut matters less than you might think — lean cuts contain nearly as much iron as fattier ones, so you do not need to seek out marbled meats to get the benefit.

Organ meats are the undisputed champions of iron density. Beef liver is one of the most iron-rich foods on the planet — a three-ounce serving contains about 5 to 6 milligrams of iron, and because it is heme iron, almost all of it is usable. Chicken liver is even more concentrated. If you can incorporate liver into your diet — even in small amounts, like a few ounces once a week — it is one of the most efficient ways to support your iron stores. Liver pâté is an approachable option if the texture of whole liver is not appealing.

Poultry provides moderate amounts of heme iron. Dark meat chicken and turkey contain more iron than white meat — about 1 to 1.5 milligrams per three-ounce serving for dark meat, compared to 0.5 to 1 milligram for white meat. Chicken thighs, legs, and wings are better choices than breast meat if you are eating for iron.

Fish and shellfish are excellent sources of heme iron. Clams are exceptional — a three-ounce serving of canned clams can contain 20 milligrams or more of iron. Oysters, mussels, sardines, and anchovies are also very high. Canned sardines (with bones) provide about 2 to 3 milligrams per serving, plus calcium, which is both a bonus and a complication — more on calcium below. Tuna, salmon, and mackerel provide moderate amounts.

If you eat animal foods, including a serving of heme iron several times per week is one of the most effective dietary strategies for maintaining healthy iron stores.

Non-Heme Iron Sources: The Plant Powerhouses

If you do not eat meat, or if you eat it sparingly, you can still build your iron stores — but you need to be more strategic. Non-heme iron is harder to absorb, but the sheer quantity available in plant foods means you can compensate with volume and with the right meal combinations.

Beans and legumes are foundational. Lentils are a standout — one cup of cooked lentils provides about 6.5 milligrams of iron. Chickpeas, kidney beans, black beans, and pinto beans all provide between 3 and 5 milligrams per cup. Soybeans and edamame are also excellent sources. The catch: beans and legumes are also high in

phytates, which block non-heme iron absorption. Soaking, sprouting, and cooking them reduces phytate content and improves absorption — more on that below.

Dark leafy greens are the foods most people think of when they think of iron. Spinach, kale, collard greens, and Swiss chard all contain iron — about 3 to 4 milligrams per cup of cooked spinach. But here is the catch: spinach is also high in oxalates, which bind to iron and reduce absorption. Cooking spinach reduces oxalate content and improves iron availability. And the common belief that spinach is a super source of iron is partly a myth — it contains iron, but you absorb very little of the non-heme iron from raw spinach because the oxalates block it. Cooked spinach is better than raw, and pairing it with vitamin C helps significantly.

Fortified cereals can be a practical source of non-heme iron, especially for women who are early in their recovery. Many breakfast cereals are fortified with iron — check the label for at least 5 to 10 milligrams per serving. The iron used in fortification is non-heme and not always well absorbed, but fortified cereals can still contribute meaningfully, especially when eaten with vitamin C (strawberries, orange juice) and without milk (calcium blocks absorption).

Pumpkin seeds, sesame seeds, and hemp seeds are concentrated sources of iron. A quarter-cup of pumpkin seeds provides about 2.5 milligrams of iron. Tahini

(sesame seed paste) provides about 2.5 milligrams per two tablespoons. Seeds are also good sources of zinc and magnesium, which support overall health.

Other good sources: Tofu and tempeh (about 3 to 6 milligrams per half-cup), quinoa (about 2.5 milligrams per cup cooked), dried apricots and raisins (about 1 to 2 milligrams per quarter-cup), blackstrap molasses (about 3.5 milligrams per tablespoon), and dark chocolate (about 3 to 4 milligrams per ounce of 70 percent or higher cocoa).

Absorption Enhancers

The single most powerful thing you can do to improve non-heme iron absorption is to pair iron-rich foods with **vitamin C**. Vitamin C converts iron into a form that is easier for your body to absorb and counteracts the effects of absorption inhibitors.

Adding a vitamin C source to an iron-rich meal can increase non-heme iron absorption by two to six times. The effect is substantial enough that it can transform a meal that barely contributes any usable iron into a meal that meaningfully supports your stores.

Good vitamin C pairings: - Squeeze lemon juice over spinach, lentils, or beans - Drink a small glass of orange juice with your fortified cereal - Add bell peppers, tomatoes, or broccoli to a bean or lentil dish - Eat strawberries or kiwi as a dessert after an iron-rich meal - Add sauerkraut or kimchi to a grain bowl with beans

The meat factor is a phenomenon where small amounts of heme iron (from meat, poultry, or fish) enhance the absorption of non-heme iron from the rest of the meal. Adding a small portion of meat — even just an ounce or two — to a plant-based meal can significantly increase the iron you absorb from the vegetables, beans, or grains in the same meal. This is good news for flexitarians who eat small amounts of meat and want to maximize their iron absorption.

Soaking, sprouting, and fermenting reduce the phytate content of grains, beans, nuts, and seeds. Phytates are natural compounds that bind to minerals — including iron — and prevent absorption. Soaking beans overnight before cooking, sprouting grains and seeds, and choosing fermented foods like tempeh over unfermented soy can all improve iron bioavailability. Sourdough bread has lower phytate content than standard bread because the fermentation process breaks down phytates.

Absorption Inhibitors

Just as some foods boost iron absorption, others block it. Understanding these inhibitors — and how to work around them — is essential to getting the most from your diet.

Phytates are found in whole grains, beans, nuts, and seeds. They are the most potent inhibitors of non-heme iron absorption. Soaking, sprouting, cooking, and fermenting reduce phytate content. If you eat a whole-

foods, plant-heavy diet, you are eating phytates regularly — and that is fine for your overall health, but it matters for iron timing.

Tannins are found in tea (black, green, and some herbal teas), coffee, red wine, and certain spices. The tannins in tea can reduce iron absorption by up to 60 percent when consumed with a meal. If you drink tea or coffee with your meals, you are significantly reducing the amount of iron your body absorbs from that meal. The solution is timing: drink tea and coffee between meals, not with them. A gap of one to two hours is enough to avoid the inhibitory effect.

Calcium from dairy products, calcium-fortified foods, and calcium supplements competes with iron for absorption. A glass of milk with your fortified cereal can reduce the iron you absorb from that cereal by 50 percent or more. The solution is the same as with tea and coffee: separate calcium from iron by at least one to two hours. If you take a calcium supplement, take it at a different time of day than your iron supplement and your iron-rich meals.

Oxalates are found in spinach, rhubarb, beets, Swiss chard, and some other vegetables. They bind to iron and reduce absorption. Cooking reduces oxalate content, which is why cooked spinach delivers more usable iron than raw spinach. Pairing high-oxalate vegetables with vitamin C also helps counteract the inhibition.

Polyphenols are found in many plant foods — berries, cocoa, spices, some herbs, and certain vegetables. They can inhibit iron absorption. The effect is generally moderate compared to tannins and phytates, but it is worth noting if your diet is exceptionally high in polyphenol-rich foods (berries, dark chocolate, turmeric, etc.). As with other inhibitors, separation is the key strategy.

The Meal-Timing Strategy

The practical takeaway from all of this is simple: separate your iron-rich foods and supplements from your inhibitors.

Here is a framework you can use:

Breakfast: If you eat iron-rich foods at breakfast (fortified cereal, eggs, spinach), avoid having coffee, tea, or milk at the same meal. Drink your coffee or tea one to two hours after breakfast. If you want to boost absorption, include a vitamin C source — a glass of orange juice, a handful of strawberries, or sliced bell peppers.

Lunch and dinner: When you eat iron-rich foods — beans, lentils, meat, dark leafy greens — pair them with vitamin C and be mindful of calcium. If you drink tea or coffee with meals, switch to drinking them between meals instead. If you take calcium supplements, take them at a different meal.

Snacks: Iron-rich snacks (pumpkin seeds, dried fruit, hummus with vegetables) are most effective when eaten away from tea, coffee, and dairy.

Supplements: Take iron supplements at least one hour before or two hours after coffee, tea, and calcium. Take them with vitamin C. Take them every other day rather than daily.

This approach does not require you to give up foods you love. It just requires you to be strategic about when you eat them.

Sample High-Iron Meal Plans

These meal plans are examples, not prescriptions. Adjust them to fit your preferences, your budget, and your lifestyle. The key is to build each meal around iron-rich foods, pair them with enhancers, and separate them from inhibitors.

Omnivore Plan

Breakfast: Two scrambled eggs with a handful of spinach and a side of bell peppers. A small glass of orange juice. Coffee one hour after breakfast.

Lunch: Grilled chicken thigh salad with chickpeas, tomatoes, and lemon vinaigrette. A handful of strawberries for dessert.

Dinner: Beef and lentil chili with tomatoes and bell peppers. Served over a small portion of rice. Avoid dairy toppings; use avocado instead.

Snack: Pumpkin seeds and an orange.

Vegetarian Plan

Breakfast: Fortified oatmeal with strawberries and a tablespoon of pumpkin seeds. A glass of orange juice. Tea one hour after breakfast.

Lunch: Lentil soup with lemon juice, served with a side of sautéed kale and bell peppers. A kiwi for dessert.

Dinner: Stir-fried tofu with broccoli, bok choy, and bell peppers in a sauce made with blackstrap molasses and tamari. Served over quinoa.

Snack: Hummus with raw vegetables and a handful of dried apricots.

Flexitarian Plan (mostly plant-based with small amounts of animal foods)

Breakfast: Smoothie with spinach, fortified plant milk, frozen berries, and a tablespoon of tahini. Coffee one hour after.

Lunch: Grain bowl with quinoa, black beans, roasted sweet potatoes, sautéed kale, and a small portion of grilled chicken or sardines. Lemon-tahini dressing.

Dinner: Stir-fried tempeh with snap peas, bell peppers, and cashews. Served over brown rice. A side of sautéed spinach with lemon.

Snack: Apple slices with almond butter and a handful of pumpkin seeds.

The "Don't Go Broke on Supplements" Rule for Food

Here is a principle that is worth repeating: supplements are a tool, not a solution. The goal is not to stay on high-dose iron supplements forever. The goal is to rebuild your iron stores to a healthy level, then maintain them through diet and — if needed — a lower maintenance dose.

This means that even if you are taking supplements now, it is worth thinking about your diet. Every meal is an opportunity to support your iron status. The better your diet supports your iron, the lower your supplement needs will be, and the sooner you can stop supplementing or drop to a low maintenance dose.

You do not need to overhaul your diet overnight. Start with one change: add a vitamin C source to one iron-rich meal per day. Move your coffee to an hour after breakfast. Add a handful of pumpkin seeds to your afternoon snack. Small, consistent changes add up.

The Cast Iron Cooking Debate

You may have heard that cooking in cast iron cookware can boost the iron content of your food. Is this true?

Yes — but the effect is variable and probably modest. Cooking acidic foods in cast iron — tomato sauce, chili, lemon-based dishes — does increase the iron content of the food. The heat and acidity leach some iron from the pan into your food. One study found that spaghetti sauce simmered in a cast iron pan contained nearly 5 milligrams of iron per serving, compared to less than 1 milligram from a non-stick pan.

However, the iron that leaches from cast iron is non-heme iron, and its absorption depends on the same factors that affect other non-heme iron — meal composition, inhibitors, enhancers. It is also hard to predict or control how much iron you get from cast iron cooking. It varies based on cooking time, acidity, and how well-seasoned the pan is.

Cast iron cooking can be a helpful addition to your iron strategy, especially if you already cook with cast iron. But it is not a reliable replacement for dietary iron sources or supplementation. Think of it as a bonus, not a foundation.

The Big Picture

Food is not a replacement for supplements when your ferritin is critically low. If your ferritin is 12, you will need oral or IV iron to rebuild your stores, and no amount of lentils and pumpkin seeds alone will get you where you need to be.

But food is what keeps you there. Once your stores are restored, a well-designed diet — one that maximizes enhancers and minimizes inhibitors — can help you maintain healthy ferritin levels without relying on high-dose supplements forever. And if your deficiency is mild (ferritin in the 30 to 50 range), dietary changes supported by targeted low-dose supplementation may be enough to bring you into the optimal range.

The best approach is an integrated one: use supplements to correct deficiency, use food to support stores, and use strategic meal timing to make every bite count. Your body does not care whether the iron comes from a pill or a plate. It just needs enough iron to do its work. Food first, supplements second, and always in conversation with your doctor about what is right for your specific situation.

Toolkit Summary

- **Heme iron from animal sources is absorbed 2-3 times better than non-heme iron from plants** — red meat, organ meats, poultry, and

shellfish are the most efficient sources. If you eat animal foods, include them regularly as part of your iron strategy.

- **Non-heme iron sources — beans, lentils, spinach, pumpkin seeds, fortified cereals — can build stores with the right approach** — pair them with vitamin C, cook your greens, and soak or sprout beans and grains to reduce phytates.
- **Vitamin C is your most powerful tool** — a squeeze of lemon, a side of bell peppers, or a glass of orange juice can double or triple the iron you absorb from a meal.
- **Separate iron from inhibitors** — drink tea and coffee between meals, not with them. Take calcium supplements and dairy at different times. A gap of one to two hours is enough.
- **Meal plans should be built around iron-rich foods, vitamin C pairings, and inhibitor separation** — whether you are omnivore, vegetarian, or flexitarian, the principles are the same.
- **Food is not a replacement for supplements when your stores are critically low** — use supplements to correct deficiency and food to maintain health. Talk to your doctor about the right approach for you.

- **Cast iron cooking adds a small, variable amount of non-heme iron to food** — a helpful bonus, but not a foundation for recovery.

ch07

IF YOU HAVE EVER STOOD IN THE BATHROOM AND WONDERED IF THIS MUCH BLOOD IS NORMAL, YOU ARE NOT ALONE — AND YOU ARE PROBABLY RIGHT TO WONDER.

Heavy menstrual bleeding is far more common than most women realize. Studies estimate that anywhere from 20 to 30 percent of women of reproductive age meet the medical criteria for heavy periods, and among them, iron deficiency is the norm rather than the exception. You cannot lose that much blood month after month without your iron stores feeling the impact.

But here is the part that does not get talked about enough: heavy bleeding is not something you have to accept. It is not just the way your body works. It is a medical condition with a name — menorrhagia — and it has treatments. Understanding what is normal, what is not, and what you can do about it is one of the most powerful steps you can take toward protecting your iron and your quality of life.

What Counts as Heavy Bleeding

The medical definition of menorrhagia is menstrual bleeding that lasts longer than seven days or involves blood loss of more than 80 milliliters per cycle. Since most women do not measure their blood loss in milliliters, doctors use practical criteria that are easier to assess.

You may have heavy bleeding if:

- You soak through a pad or tampon every one to two hours
- You need to change your protection during the night
- You pass clots larger than a quarter
- Your period lasts more than seven days
- You bleed through your clothing or onto your bedding
- You need to use both a pad and a tampon together to manage the flow
- You feel like you cannot leave the house on your heaviest days

If any of these sound familiar, it is worth taking seriously. Not because you need to compare yourself to other women — there is no prize for having the heaviest period — but because every milliliter of blood you lose is iron that your body has to replace. The average woman loses about 30 to 40 milliliters per cycle. A woman with

heavy periods can lose 80, 100, or even more. That is the difference between an iron drain you can keep up with and one that slowly depletes you month after month.

One milliliter of blood contains roughly 0.5 milligrams of iron. A normal period costs you about 15 to 25 milligrams of iron. A heavy period can cost you 50 milligrams or more per cycle. Over the course of a year, that adds up to a deficit of hundreds of milligrams — more than most women can replace through diet alone.

This is why heavy menstrual bleeding is the single most common cause of iron deficiency in women of reproductive age. It is not a side note. It is the main event.

Common Causes of Heavy Bleeding

Heavy periods are not random. They usually have an underlying cause, and identifying that cause is the first step toward effective treatment.

Uterine fibroids are noncancerous growths in the uterine muscle wall. They are incredibly common — by age 50, about 70 percent of white women and 80 percent of Black women have them — and they are a leading cause of heavy bleeding. Fibroids can distort the uterine cavity, increase the surface area of the endometrium, and interfere with contractions that help control bleeding. Black women are disproportionately affected, developing fibroids at younger ages with more severe symptoms.

Endometriosis occurs when tissue similar to the uterine lining grows outside the uterus — on the ovaries, fallopian tubes, bladder, bowel, and other pelvic structures. It affects roughly one in ten women and can cause both heavy periods and severe pelvic pain. The endometrial-like tissue responds to hormonal changes during your cycle, bleeding in places it should not, which contributes to inflammation, pain, and iron loss.

Adenomyosis is a condition where the endometrial tissue grows into the muscular wall of the uterus rather than staying in the lining. It thickens the uterine wall, increases the surface area of bleeding tissue, and can cause extremely heavy, painful periods. Adenomyosis is often underdiagnosed because it requires imaging — usually an ultrasound or MRI — to confirm, and many women live with it for years without knowing they have it.

Polycystic ovary syndrome (PCOS) is a hormonal disorder that affects ovulation and menstrual regularity. While PCOS is more commonly associated with infrequent or absent periods, some women with PCOS experience very heavy bleeding when they do menstruate. This happens because the uterine lining builds up thicker than normal during long stretches without ovulation, and when it finally sheds, the bleeding can be substantial.

Thyroid disorders — both hypothyroidism and hyperthyroidism — can affect menstrual bleeding. Hypothyroidism is particularly associated with heavy, prolonged periods. The thyroid gland regulates your metabolism, and when it is underactive, it disrupts the

hormonal balance that controls your menstrual cycle. Treating the underlying thyroid condition often resolves the heavy bleeding.

Coagulation disorders like von Willebrand disease affect your blood's ability to clot. Women with bleeding disorders often have heavy periods from their very first menstrual cycle. If you have had heavy periods since adolescence, you have a family history of bleeding problems, or you bruise easily and have frequent nosebleeds, a coagulation disorder could be the underlying cause.

Hormonal imbalances during perimenopause — the years leading up to menopause — can cause unpredictable and sometimes very heavy bleeding. As ovulation becomes irregular, estrogen levels can fluctuate dramatically, and the uterine lining may build up more than usual before shedding.

Tracking Your Cycle and Symptoms

Before you talk to your doctor about treatment, you need data. Not vague recollections of how heavy your last period was, but concrete information that helps you and your provider understand the pattern.

Here is what to track for at least two to three cycles:

- **Flow intensity** — Use a pad or tampon count. How many do you use per day on your heaviest days? How many total per cycle?

- **Clot size** — Are you passing clots smaller than a quarter, larger than a quarter, or larger than a golf ball? Size matters for diagnosis.
- **Duration** — How many days does your period last? Is it getting longer over time?
- **Pain level** — Rate your cramping on a scale of 1 to 10. Note when it peaks and what helps.
- **Breakthrough bleeding** — Do you bleed or spot between periods? When?
- **Associated symptoms** — Fatigue, dizziness, shortness of breath, anxiety — especially in the week after your period ends.
- **Iron symptoms** — How do you feel in the days after your period compared to the rest of your cycle?

A period tracker app can help, but a simple notebook or calendar works just as well. The goal is to walk into your doctor's office with a clear picture, not a vague sense that something might be off.

Talking to Your Gynecologist

The conversation about heavy periods can feel awkward. Many women normalize their bleeding because they have never known anything different, or because they assume that heavy periods are just part of being a woman. They are not. And your doctor needs to hear you say it out loud.

Here is how to start the conversation.

Use specific language. Say "I think I have heavy menstrual bleeding" or "I want to be evaluated for menorrhagia." Naming the condition signals that this is a medical concern, not casual complaining.

Bring your tracking data. "I am changing a super-plus tampon every two hours on my heaviest day, and I bleed through overnight protection. This has been happening for the past year, and it is getting worse." Specifics are harder to dismiss.

Ask about imaging. If your doctor does not mention it, ask: "Could we do an ultrasound to check for fibroids or adenomyosis?" Many causes of heavy bleeding are only visible on imaging, and you cannot treat what you cannot see.

Ask about iron testing. "I would like my ferritin checked, because I know heavy bleeding can cause iron deficiency." If your doctor has not already ordered it, this is the time.

Be persistent. If your doctor says "it is probably just perimenopause" or "some women have heavy periods" without offering a treatment plan, ask: "What are my options for reducing this bleeding?" You are entitled to treatment, not just explanation.

Treatment Options: What Is Available and When Each Makes Sense

The treatment for heavy bleeding depends on the underlying cause, your goals around fertility, your age, and your personal preferences. Here is a breakdown of the main options.

Nonsteroidal anti-inflammatory drugs (NSAIDs) like ibuprofen and naproxen reduce prostaglandin production, which can decrease menstrual flow by 20 to 30 percent. They also reduce cramping. NSAIDs are most effective when you start taking them a day or two before your period begins and continue through your heaviest days. They are not a solution for severe bleeding, but they are a useful tool, especially if your bleeding is mild to moderate or you are waiting for other treatments to take effect.

Tranexamic acid helps your blood clot more effectively. It reduces heavy menstrual bleeding by about 40 to 60 percent and is taken only during your period — typically three to five days per month. It does not affect your hormones, does not interfere with fertility, and has been used safely for decades. For many women, it is the first-line medical treatment for heavy periods, and it can be life-changing. You take it at the start of your period, and within a day or two, you notice the difference. If your doctor has not offered it and you have heavy bleeding, ask about it.

Hormonal birth control — pills, patches, rings, hormonal IUDs, implants, and shots — work by thinning the uterine lining and regulating your cycle. The hormonal IUD (Mirena, Liletta, Kyleena) is particularly effective. It releases a low dose of progestin directly into the uterus, thinning the endometrial lining significantly. Many women find their periods become much lighter — some stop entirely — within three to six months of insertion. It is one of the most effective non-surgical treatments for heavy periods and lasts for five to eight years depending on the device. Combined oral contraceptives can also reduce flow and help regulate your cycle.

Endometrial ablation is a minimally invasive procedure that removes or destroys the lining of the uterus. It is typically only recommended for women who have completed childbearing, because it significantly reduces the chances of pregnancy. The procedure takes about an hour, is done under anesthesia, and has a short recovery time. Most women experience dramatically lighter periods afterward — many stop bleeding entirely. Ablation is a good option for women who want a permanent solution but are not ready for a hysterectomy.

Myomectomy is surgery to remove fibroids while leaving the uterus intact. If fibroids are causing your heavy bleeding, removing them can restore normal flow. The approach depends on the size and location of the fibroids — hysteroscopic (through the cervix), laparoscopic (through small incisions), or open (through

an abdominal incision for large or complex fibroids). Recovery time varies. Myomectomy preserves fertility, making it a good option for women who want to become pregnant.

Uterine artery embolization blocks the blood supply to fibroids, causing them to shrink. A radiologist inserts a catheter through a small incision in your groin or wrist and guides it to the arteries feeding the fibroids, then injects tiny particles that block those vessels. The fibroids shrink over several months, and bleeding typically decreases. Recovery is usually faster than with surgery.

Hysterectomy is surgical removal of the uterus. It is the only definitive cure for heavy menstrual bleeding, but it is major surgery with a significant recovery period and permanent consequences — you will no longer be able to become pregnant. It is reserved for women who have not responded to other treatments, who have completed childbearing, or who have additional conditions that make it the best option. It should never be the first treatment offered without discussing alternatives.

The right treatment for you depends on your specific situation. There is no one-size-fits-all answer, and a good doctor will walk you through the options based on your diagnosis, your goals, and your preferences.

Emergency Iron Protocol for Heavy Bleeders

If you have heavy periods, you cannot afford to take a passive approach to your iron. You need to be proactive — because even with the best treatment for your bleeding, it takes time to bring heavy periods under control, and every cycle between now and then is still costing you iron.

Here is what a proactive iron strategy looks like for heavy bleeders.

Get tested now. Do not wait until you feel terrible. If you know you have heavy periods, ask for a full iron panel — including ferritin — at your next visit. Establish your baseline so you know where you are starting from.

Supplement proactively. If your ferritin is below 50, talk to your doctor about starting oral iron supplementation. For heavy bleeders, waiting until ferritin drops below 30 is waiting too long. The goal is to keep your stores high enough that monthly blood loss does not crash you.

Adjust your dose around your period. Some women benefit from increasing their iron intake — either through supplementation or iron-rich meals — in the days leading up to and during their period. This is a conversation to have with your doctor, but the logic is straightforward: you are about to lose blood, so front-loading iron can help buffer the hit.

Watch for post-period crashes. The week after your period ends is when iron deficiency symptoms often peak for heavy bleeders. If you notice a predictable pattern of exhaustion, brain fog, or mood changes in that window, it is a sign that your period is depleting you faster than your body can recover. That is useful information to bring to your doctor.

Consider IV iron as a bridge. If your ferritin is critically low and you are waiting for your bleeding treatment to take effect, talk to your doctor about whether IV iron could help you recover your stores more quickly. For heavy bleeders with severe depletion, IV iron can provide a faster recovery that helps you feel better while you address the underlying cause of the blood loss.

Celebrate small wins. Every cycle that is lighter than the last is a victory. Every ferritin point that goes up instead of down is progress. The goal is not perfection — it is a trend in the right direction.

The Big Picture

Heavy periods are not a life sentence. They are a medical condition with causes you can identify and treatments that work. But the first step is acknowledging that your bleeding is not normal and that you deserve help.

For many women, treating the heavy bleeding changes everything — not just their iron levels, but their quality of life, their energy, their ability to plan around their period

instead of being ruled by it. If you have been living with heavy periods, you may have forgotten what it feels like to not have to manage them. Treatment can bring that back.

And in the meantime, while you are working on the bleeding, you can protect your iron. Supplements, timing, and awareness of your cycle can keep your stores from bottoming out month after month. You do not have to choose between treating the bleeding and protecting your iron. They are two sides of the same coin, and you deserve both.

Toolkit Summary

- **Heavy menstrual bleeding (menorrhagia) affects 20 to 30 percent of women** — it is defined by periods lasting more than 7 days, soaking through protection every 1-2 hours, passing large clots, or blood loss over 80 mL per cycle. It is the leading cause of iron deficiency in women.
- **Common causes include fibroids, endometriosis, adenomyosis, PCOS, thyroid disorders, and coagulation disorders** — identifying the cause through imaging and lab work is the first step toward effective treatment.

- **Track your cycle with specific data** — pad/tampon counts, clot size, duration, pain level, and post-period iron symptoms. Concrete numbers help your doctor take your bleeding seriously.
- **Treatment options range from NSAIDs and tranexamic acid to hormonal IUDs, ablation, myomectomy, and hysterectomy** — each has a place depending on the cause, your fertility goals, and your preferences. Tranexamic acid and the hormonal IUD are first-line options for many women.
- **Heavy bleeders need a proactive iron strategy** — get tested early, supplement when ferritin is below 50, adjust dosing around your cycle, and consider IV iron as a bridge while you work on treating the bleeding. Talk to your doctor about what approach is right for your situation.

ch08

PREGNANCY IS THE MOST INTENSE PHYSICAL DEMAND YOUR IRON STORES WILL EVER FACE. IT IS NOT A GENTLE DRAIN — IT IS A SUSTAINED, MONTHS-LONG EXTRACTION THAT YOUR BODY IS DESIGNED TO HANDLE, BUT ONLY IF IT HAS ENOUGH IRON TO BEGIN WITH. AND FOR TOO MANY WOMEN, IT DOES NOT.

Here is the scale of what happens. During pregnancy, your blood volume increases by nearly 50 percent. That extra blood is not optional — it is necessary to support the placenta, the growing baby, and the changes happening in your own organs. But blood is made of hemoglobin, and hemoglobin is made of iron. Your body needs to produce roughly 500 milligrams of additional iron just to support that expanded blood volume. The baby needs another 300 to 400 milligrams of iron for its own blood supply and tissue development. The placenta and umbilical cord require their share. In total, a healthy pregnancy demands about 1,000 milligrams of iron above and beyond what your body normally uses.

That is the equivalent of emptying your iron savings account and then some.

Why Ferritin Drops in Pregnancy

If you start pregnancy with adequate iron stores — a ferritin of 70 or above — your body can draw on those reserves to meet the increased demand. Your ferritin will drop as the pregnancy progresses because you are spending your savings. That is normal. The problem is when you start with low or borderline stores.

By the second trimester, your blood volume has expanded significantly, and the fetal demand for iron is ramping up. This is when ferritin levels often take their sharpest dive. Women who entered pregnancy with ferritin in the 30 to 50 range may find themselves in single digits by the third trimester. And by that point, you are not just tired from growing a human — you are iron deficient, and your body is struggling to keep up.

The screening picture is frustrating. Many prenatal care protocols include a complete blood count (CBC) that checks hemoglobin, but not ferritin. Your hemoglobin may stay in the normal range through much of your pregnancy because your body is prioritizing red blood cell production over everything else. By the time hemoglobin drops, your iron stores have been depleted for months. Some women are never tested for ferritin at any point during their pregnancy.

If you are pregnant or planning to become pregnant, ask your provider to check your ferritin — not just your hemoglobin. The American College of Obstetricians and

Gynecologists recommends screening for anemia in pregnancy, but screening for iron deficiency specifically is not universal. You may need to advocate for this test.

Safe Iron Supplementation in Pregnancy

Prenatal vitamins typically contain iron — usually about 27 to 30 milligrams of elemental iron, which is the recommended daily allowance for pregnancy. For women who start with adequate stores, this may be enough to prevent deficiency. For women who are already low or who have additional risk factors — heavy bleeding before pregnancy, closely spaced pregnancies, a vegetarian or vegan diet — it often is not enough.

The challenge is that pregnancy can make iron supplementation harder to tolerate. Nausea and vomiting in the first trimester can make any pill difficult, and iron can worsen constipation, which is already a common complaint in pregnancy. Ferrous bisglycinate is often better tolerated than ferrous sulfate during pregnancy. Taking iron at a different time of day than your prenatal vitamin — especially if your prenatal contains calcium — can help with absorption and tolerance.

If your ferritin is low during pregnancy, talk to your doctor about whether a higher dose of oral iron is appropriate. Studies have shown that iron supplementation in pregnancy reduces the risk of maternal iron deficiency anemia and improves outcomes for both mother and baby. Higher doses should always be

guided by your lab results, not taken on your own — because iron needs are real, but iron overload during pregnancy carries its own risks.

Some women with severe deficiency during pregnancy are candidates for IV iron. Intravenous iron is considered safe in the second and third trimesters for women who cannot tolerate oral supplements or whose deficiency is too severe to correct with oral iron alone. If your ferritin is critically low, or if you are diagnosed with iron deficiency anemia during pregnancy, ask your doctor about whether IV iron is an option for you. It can restore your stores quickly and help you enter the third trimester and delivery in a stronger position.

Postpartum Hemorrhage and Iron Loss

Childbirth itself is a major iron event. A typical vaginal delivery involves about 500 milliliters of blood loss. A cesarean delivery averages about 1,000 milliliters. Postpartum hemorrhage — defined as blood loss greater than 500 milliliters for a vaginal birth or 1,000 milliliters for a cesarean — occurs in about 1 to 5 percent of deliveries and can involve much greater blood loss.

Every milliliter of blood lost during delivery is iron you will need to replace. A woman who loses 500 milliliters of blood has lost roughly 250 milligrams of iron — more than most women can absorb from oral supplements in a

month. A woman who experiences a postpartum hemorrhage can lose 1,000 to 2,000 milliliters or more, representing a staggering iron deficit.

This is why the immediate postpartum period is such a critical time for iron monitoring. If you had a difficult delivery, significant blood loss, or entered labor with low iron stores, you are at high risk for postpartum iron deficiency and should be tested within the first few weeks after delivery.

The Fourth Trimester Iron Crisis

The fourth trimester — the first three months after birth — is a period of intense physical recovery, sleep deprivation, and enormous demands on your body. And for many women, it is also when iron deficiency reaches its peak.

You may have entered pregnancy with adequate iron. You may have been tested and supplemented. You may have made it through delivery without complications. But the combination of blood loss during birth, the normal drop in ferritin that happens in the third trimester, and the demands of breastfeeding can leave you more depleted than you have ever been.

The symptoms of postpartum iron deficiency overlap heavily with the normal experience of new motherhood: exhaustion, brain fog, mood changes, difficulty concentrating, feeling overwhelmed. Every new mother is tired. But iron deficiency fatigue is different — it is the

kind of tired that does not improve even when the baby sleeps. It is the kind that makes you feel like you cannot get through the day, like your body is running on fumes.

If you are a new mother and you feel like something is wrong beyond normal exhaustion, trust that feeling. Ask for a ferritin test. The postpartum period is when women are at their most vulnerable to iron deficiency, and it is also when the condition is most likely to be dismissed as just part of being a new parent.

Baby Blues versus Iron Deficiency Overlap

Postpartum mood changes are common — the baby blues affect up to 80 percent of new mothers, typically starting a few days after birth and resolving within two weeks. Postpartum depression and anxiety are more persistent and require treatment. Iron deficiency can contribute to both.

The overlap between postpartum iron deficiency and postpartum mood disorders is significant and underrecognized. Iron is required for dopamine and serotonin synthesis — the same neurotransmitters that regulate mood, motivation, and emotional resilience. When your brain is low on iron, you are more vulnerable to mood disturbances. And the physical exhaustion of iron deficiency can make it harder to cope with the normal challenges of new motherhood.

Several studies have found that low ferritin in the postpartum period is associated with higher rates of depression and anxiety. One study found that women with postpartum iron deficiency were twice as likely to report depressive symptoms. Another found that iron supplementation in iron-deficient postpartum women improved mood and energy levels.

If you are struggling with your mood after having a baby — whether it feels like depression, anxiety, or just an overwhelming sense that something is not right — ask for a ferritin test. It is a simple blood draw, and it could change the way you understand what you are going through. Postpartum mood disorders are real and deserve treatment, but iron deficiency should be part of the conversation, not an afterthought.

Breastfeeding and Iron

Breastfeeding adds another layer to the iron story. Breast milk is naturally low in iron — it contains about 0.2 to 0.4 milligrams per liter. This is by design; high iron levels in breast milk can increase the risk of infection in newborns because many bacteria thrive on iron. But it means that breastfeeding does not directly drain your iron stores the way pregnancy and delivery do.

What breastfeeding does is delay the return of your menstrual cycle. For women who breastfeed exclusively, periods may not return for six months or longer. This is actually protective for your iron — no periods means no

monthly blood loss. But when your period does return, it may be heavy as your body readjusts, and the combination of breastfeeding fatigue, sleep deprivation, and already-depleted iron stores can hit hard.

The main risk during breastfeeding is that you enter the postpartum period with severe deficiency and do not replenish because you are focused on the baby and not on yourself. Many women do not have their iron checked after delivery unless there was a known hemorrhage. If you are breastfeeding and exhausted in a way that feels deeper than normal new-parent tiredness, ask for testing.

When to Push for IV Iron

IV iron during and after pregnancy is safe, effective, and underused. If you fall into any of these categories, it is worth having a conversation with your provider about whether IV iron is right for you.

- You are diagnosed with iron deficiency anemia during pregnancy and your ferritin is critically low
- You cannot tolerate oral iron supplements due to nausea or digestive side effects
- Your ferritin does not improve after several months of oral supplementation
- You experienced a postpartum hemorrhage
- You had a cesarean delivery and your ferritin was low going into it

- You are in the postpartum period and your ferritin is below 30, especially if you are symptomatic

The hesitation around IV iron in pregnancy is usually about safety, but modern formulations — particularly ferric carboxymaltose and iron sucrose — have strong safety records in pregnancy and postpartum. The benefits of correcting severe iron deficiency — more energy, better mood, improved ability to care for your baby — often outweigh the small risks of the infusion.

You may need to advocate for yourself on this one. Not all providers are familiar with IV iron protocols for pregnancy and postpartum. If your doctor is hesitant, ask for a referral to a hematologist or a maternal-fetal medicine specialist. Having the conversation is worth it.

The Fourth Trimester Recovery Plan

If you are postpartum or planning a pregnancy, here is a framework for protecting your iron through the entire journey.

Before pregnancy: Get your ferritin checked. If it is below 50, work on bringing it up before you conceive. Entering pregnancy with solid iron stores is one of the best things you can do for yourself and your baby.

During pregnancy: Ask for ferritin testing at least once per trimester. If your ferritin drops below 30, discuss supplementation with your doctor. If you cannot tolerate oral iron, ask about IV iron.

At delivery: If you have significant blood loss, ask for a ferritin and hemoglobin check before you leave the hospital. Do not wait for your six-week checkup.

After delivery: Get your ferritin tested at your postpartum visit. If it is low, treat it. Do not accept exhaustion as just part of new motherhood. It may be, but it does not have to be.

While breastfeeding: If you are struggling with energy, mood, or focus, check your ferritin. Breastfeeding is protective in some ways, but it does not protect you from deficiency that started before or during pregnancy.

Between pregnancies: If you are planning to have more children, make sure your iron stores are fully restored before you conceive again. Closely spaced pregnancies — less than 18 months apart — significantly increase your risk of iron deficiency in the second pregnancy because your body may not have had time to rebuild its stores.

The Big Picture

Pregnancy and the postpartum period are the most iron-intensive years of your life. Your body is doing something extraordinary, and it needs extraordinary resources to do it well. Iron is not optional in this equation — it is essential.

The good news is that the tools exist. Testing, oral supplementation, and IV iron are all available options. The hard part is often the advocacy — asking for the test, asking for the treatment, asking for the follow-up. But you are worth that effort. Your health matters, not just for your baby, but for you. You deserve to feel like yourself during one of the most transformative periods of your life.

Toolkit Summary

- **Pregnancy demands about 1,000 milligrams of extra iron** — for 50 percent more blood volume, the baby's needs, and placental development. If you start with low stores, you will be depleted by the third trimester.
- **Ferritin drops most sharply in the second and third trimesters** — standard prenatal screening often checks only hemoglobin, not ferritin. Ask for a full iron panel, especially if you have risk factors.

- **Oral iron is safe in pregnancy** — prenatal vitamins typically provide 27-30 mg, but higher doses may be needed if you are deficient. Ferrous bisglycinate is often better tolerated. Talk to your doctor about the right dose for your situation.
- **Postpartum hemorrhage and childbirth itself cause significant iron loss** — a typical delivery costs 250 mg of iron or more. Get tested after delivery if you had significant blood loss.
- **Postpartum iron deficiency masquerades as normal new-mother exhaustion** — if you feel like something is wrong beyond tiredness, trust yourself and ask for a ferritin test. The overlap with postpartum depression and anxiety is significant.
- **IV iron is a safe and effective option during pregnancy and postpartum** — especially for women with severe deficiency, intolerance to oral iron, or significant blood loss. Advocate for yourself if you think you need it. Talk to your doctor about whether IV iron is right for you.

ch09

IF YOU HAVE EVER SAT IN A THERAPIST'S OFFICE OR A PSYCHIATRIST'S OFFICE DESCRIBING YOUR SYMPTOMS — THE PERSISTENT LOW MOOD, THE RACING THOUGHTS, THE INABILITY TO CONCENTRATE, THE FEELING THAT YOUR BRAIN IS WRAPPED IN COTTON — AND WALKED OUT WITH A DIAGNOSIS THAT DID NOT QUITE FIT, THIS CHAPTER IS FOR YOU.

Iron deficiency does not just affect your body. It affects your brain. And because the symptoms of low brain iron — anxiety, depression, brain fog, poor focus, irritability — overlap so thoroughly with common mental health diagnoses, millions of women are being treated for conditions they may not have, or whose severity is driven in large part by a mineral deficiency that no one thought to check.

This is not to say that mental health conditions are not real. They are. But iron deficiency can cause mental health symptoms, worsen existing conditions, and make treatment less effective. And unlike many drivers of mental health struggles, iron deficiency is straightforward to test for and treat.

The Iron-Brain Connection

Your brain is the most metabolically active organ in your body. It accounts for only about 2 percent of your body weight but consumes roughly 20 percent of your oxygen and 20 percent of your energy. And iron is the central player in both oxygen transport and energy production.

But iron's role in the brain goes deeper than fuel delivery. Iron is essential for the synthesis of neurotransmitters — the chemical messengers that allow your brain cells to communicate.

Dopamine is the neurotransmitter of motivation, focus, pleasure, and reward. It is what gives you the drive to start a task and the satisfaction of completing it. Dopamine synthesis requires an iron-dependent enzyme called tyrosine hydroxylase. When your brain does not have enough iron, it cannot produce dopamine efficiently. The result looks a lot like the core symptoms of ADHD: difficulty initiating tasks, poor focus, low motivation, a sense that you have to push yourself to do things that should feel automatic.

Serotonin is the neurotransmitter of mood regulation, appetite, sleep, and emotional stability. It is synthesized through a pathway that also depends on iron. Low serotonin is associated with depression, anxiety, irritability, and sleep disturbances. When brain iron is low, serotonin production drops, and your emotional regulation suffers.

Norepinephrine is involved in alertness, attention, and the fight-or-flight response. Its synthesis also requires iron. Low norepinephrine can contribute to fatigue, low energy, and difficulty staying alert.

Your brain also needs iron to build myelin — the insulating sheath around your nerve cells that allows electrical signals to travel quickly. Without enough iron, myelination slows, and your neural communication becomes less efficient. This is a direct contributor to the brain fog that so many iron-deficient women describe: the feeling that your thinking is slow, that words are harder to find, that you are processing information through a layer of resistance.

Why Low Iron Causes Brain Fog

Brain fog is one of the most common and most distressing symptoms of iron deficiency. Women describe it as feeling like their thinking is fuzzy, like they are moving through mental molasses, like they have to work harder to think clearly.

The mechanisms are multiple. Your brain is sensitive to oxygen levels, and low iron means less oxygen reaching your brain tissue. But it is not just about oxygen. The neurotransmitter disruption slows communication between brain regions. The energy deficit in your neurons means they cannot fire as efficiently. The myelin deficit means signals travel more slowly.

This combination — reduced oxygen, less dopamine, less serotonin, slower neural transmission — creates the experience of brain fog. You are not imagining it. You are not lazy or unfocused. Your brain is literally operating at reduced capacity because it does not have the raw materials it needs to function at full speed.

Many women report that brain fog was the symptom that convinced them something was wrong — more than fatigue, more than hair loss. It is the symptom that affects work performance, parenting, relationships, and the sense of being able to trust your own mind. If your brain feels unreliable, everything feels harder.

Anxiety: The Misdiagnosis

Anxiety is a common symptom of iron deficiency, and it is frequently misdiagnosed as generalized anxiety disorder (GAD). The mechanism is biological: your body's stress response system is activated by low oxygen and by dysregulated neurotransmitter production. When your brain is starved for iron, it can trigger a state of physiological arousal that feels exactly like anxiety.

Women with iron deficiency describe feeling on edge, restless, easily startled, and unable to settle their minds. They may experience racing thoughts, a sense of dread, physical symptoms like a racing heart or tight chest — all of which are indistinguishable from anxiety of psychological origin.

The difference is that this type of anxiety may not respond fully to therapy or anti-anxiety medication, because it has a physiological driver that those treatments do not address. One study found that women with iron deficiency were significantly more likely to be diagnosed with anxiety disorders — and that treating the iron deficiency reduced anxiety symptoms in many cases.

This is not to say that your anxiety is "just" iron deficiency. But if you have been treated for anxiety and your symptoms have not fully resolved, or if your anxiety started or worsened around the same time as other iron-related symptoms (fatigue, brain fog, heavy periods, hair loss), it is worth asking whether low iron is part of the picture.

Depression and Treatment-Resistant Depression

The link between iron deficiency and depression is one of the most well-documented — and most overlooked — connections in women's mental health.

Research has consistently found that women with iron deficiency are more likely to meet criteria for depression than women with normal iron stores. One study found that the odds of depression were nearly double in women with iron deficiency compared to iron-replete women. Another study found that women with iron deficiency anemia had significantly higher depression scores than women with normal iron levels.

The striking finding is what happens when the deficiency is treated. Several studies have reported that iron supplementation improves mood in iron-deficient women — and that the improvement can be significant. One study found that iron supplementation improved depressive symptoms in women who had not responded to antidepressant medication. The implication is profound: for some women, depression that appears treatment-resistant may actually be depression that is driven or worsened by a correctable mineral deficiency.

This is not about blaming depression on iron or suggesting that iron supplements can replace antidepressants. But it is about making sure that iron deficiency is on the diagnostic radar before a woman is labeled as treatment-resistant. If your depression has not responded to multiple medication trials, or if it started alongside other iron symptoms, ask for a ferritin test. It is a simple blood draw that could change your treatment trajectory.

The ADHD-Iron Connection

The overlap between iron deficiency and ADHD symptoms is striking enough that several research groups have investigated whether low iron contributes to ADHD severity — and the evidence suggests it does.

Dopamine is the neurotransmitter most closely associated with ADHD. Stimulant medications like methylphenidate and amphetamines work by increasing

dopamine availability in the brain. If iron deficiency impairs dopamine synthesis, it makes sense that low iron would worsen ADHD symptoms — and that treating the deficiency could improve them.

Studies in children have found that low ferritin is associated with more severe ADHD symptoms and that iron supplementation can improve those symptoms. The research in adults is less extensive, but the biological mechanism is the same. Women with ADHD who are also iron deficient may find that their symptoms feel harder to manage, that their medication is less effective, or that the combination of low iron and ADHD creates a cognitive burden that neither condition alone would explain.

If you have ADHD and you have noticed that your symptoms seem worse than they used to be, or that your medication does not work as well as it once did, check your iron. Restoring your ferritin to optimal levels may not resolve your ADHD — but it may make your baseline easier to manage and restore some of the cognitive capacity that deficiency has been stealing.

The Cognitive Improvement Timeline

If you start iron supplementation and your brain symptoms improve, you might wonder how long it takes and what to expect. Here is the general pattern that many women report.

Within the first one to two weeks. Some women notice subtle improvements in energy and mental clarity within days. The brain fog may start to lift. You might find it easier to concentrate, to find words, to follow a train of thought. These early improvements are often small — not a complete transformation, but a sense that something is shifting.

Around four to six weeks. More noticeable changes in mood and focus. Anxiety may feel less acute. Your baseline mood may lift. You may feel more motivated and less overwhelmed. This is also when some women report that their response to ADHD medication improves — the medication seems to work better or last longer.

At two to three months. Significant improvements for many women. The brain fog that felt constant may be mostly gone. Your ability to concentrate, remember, and process information may feel closer to what it was before the deficiency took hold. Mood stability often improves in this window.

Full recovery of brain iron takes longer. The brain is protective of its iron stores and may take longer to replenish than the rest of your body. Even after your ferritin has returned to a healthy level, your brain may need additional time to fully recover. Some women report that cognitive improvements continue for six months or more after starting treatment.

This timeline is not universal. Some women respond faster; some take longer. The important thing is that improvement is possible, and if you are not seeing any

changes after several months of consistent treatment, it is worth revisiting your dose, your form of iron, and whether there are other factors at play.

Advocating for Iron Testing with Mental Health Diagnoses

If you are seeing a therapist, psychiatrist, or other mental health provider, you may need to be the one who brings up iron. Most mental health professionals do not routinely screen for iron deficiency, even though the evidence connecting iron to mental health is robust.

Here is how to have that conversation.

Connect the dots for your provider. "I have been reading about the connection between iron deficiency and mental health. I know that iron is needed to produce dopamine and serotonin, and I am wondering whether low iron could be contributing to my symptoms. Could you help me get a ferritin test?"

Bring your physical symptoms into the conversation. If you have heavy periods, fatigue, hair loss, ice cravings, or any of the physical signs of iron deficiency, mention them. The combination of mental health symptoms and physical symptoms makes a stronger case for testing.

Frame it as ruling something out. "I would like to rule out iron deficiency as a contributing factor to my mental health, so that we can make sure we are treating the right thing." This is a reasonable request, and most providers will agree.

If your provider is dismissive, ask why. "I understand that iron deficiency may not be the main cause of my symptoms, but since it is treatable and easy to test for, I would feel more confident ruling it out. Could we add ferritin to my next blood work?"

Consider an integrated approach. If your mental health provider cannot or will not order iron testing, your primary care doctor can. You can also order your own labs through direct-to-consumer services and bring the results to both your medical and mental health providers.

The Big Picture

Your brain is not separate from your body. The idea that mental health symptoms live entirely in the realm of psychology — divorced from biochemistry, nutrition, and physical health — is outdated and actively harmful to women who are suffering from a condition that has a clear physiological basis.

Iron deficiency can cause or worsen anxiety, depression, brain fog, and ADHD symptoms. It can make mental health treatment less effective. And it is completely treatable.

If you have been struggling with your mental health and wondering why treatment has not worked as well as you hoped, consider whether iron could be part of the picture. It is not the answer for everyone. But it is a possibility that deserves to be explored — and too often, it is not explored at all.

You are not broken. You may be deficient. And deficiency has a fix.

Toolkit Summary

- **Iron is essential for brain function** — it is required for dopamine, serotonin, and norepinephrine synthesis, for oxygen delivery to the brain, for energy production in brain cells, and for myelin formation. Low iron directly impairs cognition and mood.
- **Brain fog is a predictable effect of low brain iron** — reduced oxygen, neurotransmitter disruption, and slower neural transmission combine to create the experience of fuzzy, slow thinking. It is a biological phenomenon, not a character flaw.
- **Anxiety and depression are frequently linked to iron deficiency** — and frequently misdiagnosed when the underlying deficiency is not identified. Iron deficiency can cause anxiety symptoms, worsen depression, and make treatment-resistant depression harder to resolve.

- **ADHD symptoms may be worsened by low iron** — because dopamine synthesis depends on iron. Restoring ferritin to optimal levels may improve focus, motivation, and medication response.
- **Cognitive improvement typically begins within 1-2 weeks** of effective supplementation and continues over several months. Full recovery of brain iron may take longer than recovery of blood iron.
- **Advocate for ferritin testing with your mental health provider** — the evidence connecting iron to mental health is strong, but most providers do not screen for it automatically. If you have mental health symptoms alongside physical signs of iron deficiency, bring up testing. Talk to your doctor about whether iron testing is right for your situation.

ch10

THE SYMPTOMS THAT SHOW UP IN YOUR APPEARANCE MIGHT NOT BE THE MOST DANGEROUS EFFECTS OF IRON DEFICIENCY, BUT THEY ARE OFTEN THE ONES THAT FINALLY GET YOU TO TAKE ACTION. THERE IS SOMETHING DEEPLY UNSETTLING ABOUT WATCHING YOUR HAIR FALL OUT IN THE SHOWER, LOOKING IN THE MIRROR AND SEEING A FACE THAT SEEMS TO HAVE LOST ITS COLOR, OR REALIZING THAT YOUR NAILS HAVE BECOME SO BRITTLE THEY BARELY GROW. THESE ARE THE VISIBLE SIGNS THAT YOUR BODY IS RUNNING LOW ON A RESOURCE IT NEEDS — AND THEY ARE THE SIGNS THAT OTHER PEOPLE CAN SEE TOO.

This chapter covers the full range of visible and physical iron deficiency symptoms: the hair loss, the changes in your skin and nails, the restless legs that keep you up at night, the strange cravings that you might have been too embarrassed to mention to anyone. These symptoms are not random. They are predictable consequences of your body prioritizing the iron it has for essential functions — red blood cell production, brain function, immune response — and cutting back on the "optional" uses of iron like hair growth and nail health.

Telogen Effluvium: Why Hair Thins Diffusely

The medical name for iron-related hair loss is telogen effluvium, and understanding how it works helps you understand why it happens the way it does — and why it takes so long to resolve.

Your hair grows in cycles. The active growth phase (anagen) lasts about two to six years for scalp hair. At any given time, about 85 to 90 percent of your hair is in this phase. The resting phase (telogen) lasts about three months. About 10 to 15 percent of your hair is in this phase at any given time, waiting to shed.

Normally, this is a slow, steady process. You lose about 50 to 100 hairs per day, and new hairs grow to replace them. The ratio of growing to resting hair stays constant.

Iron deficiency disrupts this balance. When your body detects that iron stores are low, it sends a signal to your hair follicles: slow down. Non-essential growth functions get deprioritized. A large number of follicles simultaneously shift from the active growth phase into the resting phase. About three months later — the length of the telogen phase — all of those follicles shed their hair at once.

This is why hair loss from iron deficiency is delayed. You do not lose hair the same week your iron drops. You lose hair about three months after your iron drops, and the shedding can continue for months afterward because

the shift happens gradually. This is also why women often do not connect the hair loss to iron deficiency — the time delay makes the relationship invisible.

The hair loss of telogen effluvium is diffuse. It does not typically cause bald patches, though hair may be noticeably thinner overall. You might notice more hair in your brush, in the shower drain, on your pillow. Your ponytail may feel thinner. Your part may look wider. You may be able to see more of your scalp when your hair is wet or under bright light.

The same process can happen after childbirth (postpartum telogen effluvium), after a major illness or surgery, after significant weight loss, and after severe stress — all of which place metabolic demands on your body that can trigger the same signal to hair follicles.

Why Hair Loss Is So Distressing

There is a reason hair loss hits harder than many other iron deficiency symptoms. It is visible. You cannot hide it the way you can hide fatigue or brain fog. People can see that your hair looks different, and some of them will comment on it. "Your hair looks thinner," someone says, meaning to be helpful, and you feel a wave of shame about something you cannot control.

Hair is culturally tied to femininity, to health, to attractiveness. Losing it can feel like losing a part of your identity. Women report that hair loss affects their

confidence, their willingness to be seen in public, their sense of themselves as healthy and vital. It is a real emotional burden, and it deserves to be acknowledged.

If you are experiencing hair loss from iron deficiency, you are not being vain or shallow to feel upset about it. It is not just hair. It is a visible signal that your body is struggling, and it changes the way you see yourself in the mirror every day.

Nail Changes: Koilonychia and Beyond

Your nails, like your hair, require iron-dependent enzymes to grow properly. When iron is low, your nails are one of the first places to show it.

The earliest sign is often brittleness. Your nails may chip, peel, and split more easily than they used to. They may grow more slowly. You might notice vertical ridges running from your cuticle to the tip — these are common in nail aging, but they can also be a sign of nutritional deficiency.

In more significant iron deficiency, nails can develop a shape called koilonychia — more commonly known as spoon nails. The nail becomes thin and concave, flattening out in the middle and rising at the edges, creating a shallow depression that can hold a drop of water. Spoon nails are a classic sign of long-standing iron deficiency. They are less common today than they were

historically because severe, prolonged deficiency is less common — but they still occur, especially in women whose iron has been low for a long time.

Nail changes are reversible. When your ferritin returns to a healthy level, your nails will grow out normally. But because nail growth is slow — a fingernail takes about four to six months to grow from cuticle to tip — it will take time to see the improvement. The new, healthy nail grows in from the cuticle, and the old, brittle or spoon-shaped nail gradually grows out and is trimmed away.

Pallor: Looking as Tired as You Feel

Pale skin — known medically as pallor — is one of the oldest recognized signs of iron deficiency. It happens because hemoglobin gives red blood cells their color, and when you have fewer red blood cells or less hemoglobin per cell, your blood is less red. Less red blood flowing through the small vessels near your skin's surface means your skin looks paler.

Pallor is most visible in areas with less pigment: the inside of your lower eyelid (the conjunctiva), your nail beds, your palms, and the lining of your mouth. If you pull down your lower eyelid and see pale pink or white instead of a healthy reddish-pink, that is a sign of reduced red blood cells — and a strong indicator that your iron is low.

Pallor can also show up as a general loss of color in your complexion. You may look washed out. People may comment that you look tired or unwell. The pallor of iron deficiency is often described as looking "sallow" or "ashen," and it does not improve with rest or better sleep.

For many women, the pallor of iron deficiency creates a frustrating cycle: you feel exhausted, and you look exhausted, which means people treat you like you are exhausted. They ask if you are okay. They suggest you take a nap. They comment on your appearance in ways that reinforce the sense that something is visibly wrong with you.

Restless Legs Syndrome: The Iron-Dopamine Connection

Restless legs syndrome (RLS) is a neurological condition characterized by an overwhelming, nearly irresistible urge to move your legs. It typically strikes in the evening or at night when you are sitting or lying down, and it can make sleep nearly impossible.

The sensation is hard to describe to someone who has not experienced it. It is not exactly pain. It is more like a deep, crawling, tingling, or electric sensation inside your legs — an internal restlessness that builds until you move, and then settles temporarily, only to return when you stop moving.

The connection between restless legs and iron is well established. Iron is required for dopamine production in the brain, and dopamine dysfunction in the basal ganglia — a region of the brain that controls movement — is a key mechanism in RLS. When iron levels in the brain are low, dopamine signaling is disrupted, and the movement control system becomes overactive, producing the sensation of restlessness.

RLS is common in iron deficiency — some studies suggest that up to 25 percent of iron-deficient women experience it — and treating the deficiency often resolves or significantly improves the symptoms. If you have restless legs, mention it to your doctor. It is a strong clue that your iron may be low, even if your hemoglobin is normal.

Pica and Ice Cravings: The Least Known Symptom

If you have ever found yourself craving ice — not enjoying ice in a drink, but actively wanting to chew it, seeking it out, consuming trays of it — you have experienced one of the most specific and least known symptoms of iron deficiency.

Pica is the medical term for craving non-food substances, and the most common form in iron deficiency is pagophagia — the craving for ice. Women with low iron report an almost compulsive urge to chew ice. They keep a cup of ice on their desk. They fill their water bottle with

crushed ice. They stop at convenience stores for a bag of ice to chew on the way home. They hide their ice consumption because they know it is strange, but they cannot stop.

No one knows exactly why iron deficiency causes ice cravings. The leading theory is that chewing ice increases blood flow to the brain — the cold stimulates vasoconstriction followed by a rebound increase in blood flow, which temporarily increases oxygen delivery to oxygen-starved brain tissue. Another theory is that ice chewing triggers a similar neurological response to pain or stress relief. Whatever the mechanism, it is a remarkably consistent symptom. Studies have found that ice craving is one of the most sensitive indicators of iron deficiency — more reliable than many of the other classic symptoms.

Pica can also involve other non-food substances: clay, dirt, cornstarch, chalk, paper, laundry detergent, and even paint chips. If you have any of these cravings — even if it feels embarrassing or strange — mention it to your doctor. It is a strong, specific signal that your iron is low, and your doctor needs to know.

The Visible Recovery Timeline

One of the hardest parts of treating iron deficiency is the waiting. The visible symptoms — the ones that bother you the most — take the longest to resolve. Here is what you can generally expect.

Energy and brain fog often improve first, within the first two to six weeks of effective supplementation. These are the symptoms that respond most quickly because they are tied to oxygen delivery and cellular energy production.

Restless legs may improve within weeks, as dopamine production begins to normalize. Many women report that their RLS symptoms decrease significantly within the first month of treatment.

Pallor improves gradually over several months as your hemoglobin rises and your red blood cells return to normal size and color. This is not a fast change, but most women notice that their skin looks healthier within two to three months.

Ice cravings can resolve surprisingly quickly. Some women report that the urge to chew ice disappears within days of starting supplementation. For others, it fades over several weeks. The resolution of ice cravings is often one of the first and most dramatic signs that treatment is working.

Hair shedding starts to decrease around three to four months after ferritin reaches a healthy level. The hair that was already in the shedding phase must complete its cycle before the reduction is visible. You may stop losing so much hair before you see new growth. Actual regrowth — thicker hair, a fuller ponytail, a narrower part — typically takes six months to a year.

Nail changes improve over four to six months, as the old, brittle nail grows out and is replaced by healthy nail from the cuticle.

The visible recovery timeline is slower than the internal one, and that can be frustrating. You might feel better — more energy, clearer thinking, less anxiety — while your hair is still falling out and your complexion is still pale. This does not mean the treatment is not working. It means your body is prioritizing the internal recovery first. The visible changes follow.

The Emotional Toll of Looking as Tired as You Feel

There is a specific kind of exhaustion that comes from looking as tired as you feel. It is the experience of watching your appearance change in ways that do not reflect who you are on the inside, and of having people respond to that changed appearance as though it is the real you.

When you look in the mirror and see a pale, drawn face with thinning hair and dark circles, it can be hard to remember that this is a temporary state — that your body is going through something, and that it can recover. The visible symptoms can make you feel like the deficiency has taken over your identity, like this tired, pale, shedding version of you is who you are now.

It is not. These symptoms are a reaction to a biological state, and when that state is corrected, they will reverse. The face in the mirror right now is not your permanent face. It is the face of a body that is low on iron and prioritizing its resources. When your stores are restored, your hair will grow back, your skin will regain its color, your nails will strengthen, and the restlessness in your legs will settle.

The waiting is hard. Be gentle with yourself while you wait. Take progress photos so you can see changes that are too gradual to notice day by day. Remind yourself that the visible symptoms always lag behind the internal recovery, and that lag does not mean failure. You are healing from the inside out, and the outside takes longer.

The Big Picture

The visible signs of iron deficiency are distressing, but they are also useful. They give you — and your doctor — concrete evidence that something is wrong. Hair loss, pale skin, brittle nails, restless legs, ice cravings — these are not signs that you are falling apart. They are signs that your body needs more iron, and they are reversible.

When your iron stores are restored, your body will redirect resources back to hair growth, nail strength, and skin health. The recovery takes time, but it happens. And when you see that first new growth — the baby hairs at

your hairline, the healthy strip of nail at your cuticle, the color returning to your face — it is a reminder that your body knows how to heal when it has what it needs.

Toolkit Summary

- **Iron deficiency causes diffuse hair loss (telogen effluvium)** — hair follicles shift from growth to resting phase when iron is low, and shedding occurs about 3 months later. Recovery takes 3-6 months after iron normalizes.
- **Nail changes include brittleness, ridges, and spoon-shaped nails (koilonychia)** — these are signs of long-standing deficiency and reverse over 4-6 months as healthy nail grows out.
- **Pallor (pale skin) is a classic visible sign** — most noticeable in the lower eyelid, nail beds, and palms. It improves gradually as hemoglobin rises.
- **Restless legs syndrome is linked to brain iron levels and dopamine function** — up to 25 percent of iron-deficient women experience it. Treatment often resolves or significantly improves RLS symptoms.
- **Ice cravings (pagophagia) are one of the most specific symptoms of iron deficiency** — if you crave ice, mention it to your doctor. It is a strong signal that your iron is low, and it often resolves within days of starting supplementation.

- **The visible timeline lags behind internal recovery** — energy improves first, followed by pallor, restless legs, and ice cravings. Hair and nails take the longest. Be patient with your body. Talk to your doctor about treatment and monitoring.

ch11

YOU HAVE DONE THE HARD WORK. YOU GOT TESTED. YOU STARTED TREATMENT. YOU NAVIGATED THE SIDE EFFECTS, ADJUSTED YOUR DIET, AND ADVOCATED FOR YOURSELF WITH DOCTORS WHO MAY OR MAY NOT HAVE BEEN ON BOARD. AND NOW, HOPEFULLY, YOU ARE FEELING BETTER — MORE ENERGY, CLEARER THINKING, STEADIER MOOD, LESS HAIR IN THE SHOWER DRAIN.

This is the moment where many women stop paying attention. They feel better, so they assume the problem is solved. They stop supplementing, stop tracking, stop thinking about iron at all. And then, months or years later, they crash again.

This chapter is about making sure that does not happen to you. Long-term iron management is not complicated, but it requires awareness. Your body's need for iron did not disappear when your ferritin hit the optimal range. It changed — but it did not disappear. And the factors that made you deficient in the first place — heavy periods, pregnancy, diet, absorption issues — may still be in play.

The goal is not to obsess over iron for the rest of your life. The goal is to build habits that keep your stores stable so you can stop thinking about iron and just live.

What Recovery Actually Looks Like

Full iron recovery is not a single number. It is an integrated process that involves your blood, your stores, your symptoms, and your daily function.

Blood recovery happens first. Your hemoglobin returns to normal as your body produces healthy red blood cells. This usually takes four to eight weeks of effective treatment, depending on the severity of your deficiency.

Store recovery takes longer. Ferritin rises more slowly — typically 5 to 10 points per month with consistent oral supplementation, faster with IV iron. Most women need three to six months to reach a ferritin of 70 or above.

Symptom recovery follows its own timeline. Energy often improves first, within two to six weeks. Brain fog and mood lift over one to three months. Hair, nails, and restless legs take longer — three to six months or more for full resolution.

Functional recovery — the point where you forget you were ever iron deficient — comes last. One day you realize you have not thought about your iron in weeks. You have energy. You are sleeping well. Your hair looks like it used to. Your brain feels sharp. That is functional recovery, and it is the goal.

Maintaining Stores After Recovery

Once your ferritin is in the optimal range — generally 70 to 100 or above — the question becomes: how do you keep it there?

The answer depends on why you became deficient in the first place. If your deficiency was caused by a one-time event — a pregnancy, a surgery, a period of heavy bleeding that has since resolved — you may be able to maintain your stores through diet alone once they are fully restored. Your iron needs have returned to baseline, and a well-designed diet may be enough.

If your deficiency is driven by ongoing factors — chronic heavy periods, a condition that impairs absorption, a vegetarian or vegan diet, ongoing endurance training — you will likely need some form of ongoing maintenance. This does not mean high-dose supplements forever. It means a lower, sustainable approach that matches your ongoing losses.

Annual ferritin testing is the single most important maintenance habit. Get your ferritin checked once a year, or more often if you have known risk factors. An annual test catches downward trends before they become symptomatic. If your ferritin drops from 85 to 55 over the course of a year, you know you are losing ground and can adjust before you crash.

A maintenance supplement strategy might mean a lower dose of oral iron — 15 to 30 milligrams of elemental iron every other day, or a few times per week — rather

than the high doses used for initial correction. Some women take a multivitamin with iron as their maintenance approach. Others take a targeted low-dose iron supplement. The right approach depends on your needs and your tolerance.

Dietary awareness continues to matter. You do not need to eat every meal through the lens of iron absorption, but keeping the principles in mind — pair iron-rich foods with vitamin C, separate iron from coffee and tea, include heme iron sources if you eat animal foods — helps maintain your stores without effort.

What to Do If Ferritin Drops Again

Ferritin can drop even when you are doing everything right. A change in your periods, a pregnancy, a new medication, a change in your diet — any of these can shift your iron balance. The key is catching the drop early and responding quickly.

If your annual ferritin test shows a significant drop — say, from 80 to 40 — do not panic. You know how to fix this. You have done it before.

Re-evaluate the cause. Have your periods changed? Have you changed your diet? Did you stop taking a supplement you were relying on? Did you start a new medication? Did you have a health event — surgery, illness, blood donation? Identifying the cause helps you decide how aggressive your response needs to be.

Increase your iron intake. If the drop is mild, adding iron-rich foods and a low-dose supplement may be enough to reverse the trend. If the drop is significant, you may need to return to a higher treatment dose for a few months.

Re-test in two to three months. After adjusting your approach, check your ferritin again to confirm that you are moving in the right direction. If the drop continues despite increased intake, talk to your doctor about whether there is a new underlying issue.

Do not wait until you are symptomatic. The advantage of annual testing is that you catch drops early, when they are easy to correct. The crash that comes from ignoring a slow decline is much harder to climb out of.

The "I Thought I Was Fine and Then I Crashed" Trap

This pattern is so common it deserves its own warning label. A woman corrects her iron deficiency. She feels better. She stops monitoring. Six months or a year later, the old symptoms start creeping back — the fatigue, the brain fog, the hair loss. She thinks she is stressed or overworked. She does not connect it to iron because she "fixed" that already. By the time she gets tested again, her ferritin is back in single digits.

The trap exists because recovery feels like a finish line. It is not. It is a waypoint. Your iron stores need ongoing attention, especially if the factors that caused your deficiency are still present. The goal is to turn that attention into a low-effort habit — an annual test, a maintenance supplement, a general awareness of your body — rather than an ongoing project.

Set a reminder on your calendar for your annual ferritin test. Keep a bottle of low-dose iron in your cabinet in case you need it. And trust your body: if the old symptoms start returning, check your iron before you assume it is something else.

Lifestyle Habits That Protect Iron

Beyond testing and supplementation, certain habits support healthy iron levels over the long term.

Manage heavy periods. If heavy periods are the main driver of your iron deficiency, treating the bleeding itself is the most effective long-term strategy. If you have not yet found a treatment that works for you, keep pushing. Every cycle you lighten is a cycle that costs your iron stores less.

Be mindful of blood loss in all forms. Blood donation is a significant iron drain. If you donate blood regularly, your ferritin will trend downward unless you supplement intentionally. The same applies to any

surgery, injury, or medical procedure that involves blood loss. Plan ahead: optimize your iron before elective procedures and check your ferritin after.

Pay attention to medications. Acid-reducing medications (PPIs, H2 blockers) reduce iron absorption over time. If you take them daily, your iron needs are higher than average. Calcium supplements should be taken at a different time than iron-rich meals and iron supplements. NSAIDs taken regularly can cause gastrointestinal bleeding, which is an invisible source of iron loss.

Time your iron-conscious eating. You do not need to be rigid, but keeping a few principles in mind helps: drink coffee and tea between meals, not with them. Pair iron-rich foods with vitamin C. Take calcium at a different time than iron. These small habits add up.

Stay on top of your health. Conditions that affect iron — thyroid disorders, celiac disease, autoimmune conditions, gastrointestinal issues — should be managed with your healthcare team. If you have a condition that affects iron absorption or loss, your ferritin should be part of your routine monitoring.

Building Your Medical Team

You do not need to navigate this alone. The right medical team makes iron management easier and more effective.

Your primary care doctor is your partner for annual ferritin testing and general health maintenance. If you have a good relationship with your PCP, they can manage most of your iron care.

Your gynecologist is essential if heavy periods are part of your story. They can diagnose and treat the underlying causes of heavy bleeding and help you find a menstrual management strategy that works for your iron.

A hematologist can be helpful for complicated cases — severe deficiency, IV iron management, hereditary conditions, or situations where your iron is not responding to treatment as expected. Do not hesitate to ask for a referral if you are stuck.

A gastroenterologist matters if you have absorption issues from celiac disease, Crohn's, colitis, or other digestive conditions. Optimizing your gut health is essential for long-term iron maintenance.

A registered dietitian can help you design a sustainable eating plan that supports your iron stores without requiring you to overhaul your entire diet.

Not everyone needs all of these specialists. But knowing who is available and when to ask for a referral gives you options.

The Identity Shift

Recovering from iron deficiency is not just a physical process. It is an identity shift.

For months or years, you lived with the assumption that you were just tired, just slow, just anxious, just forgetful. You built your life around that assumption. You stopped expecting as much from yourself. You lowered your standards for your energy, your focus, your mood. You learned to operate at 60 or 70 percent capacity and tell yourself it was fine.

Recovery disrupts that. When your energy returns, you have to reconfigure your expectations of what you are capable of. When your brain clears, you have to get used to trusting your mind again. When your mood stabilizes, you have to unlearn the anxious patterns that you developed to cope with a brain that was not getting enough iron.

This is a good problem to have. But it can be disorienting. Give yourself permission to adjust. You are not the same person you were when you were deficient, and it takes time to figure out who you are at full capacity.

A Final Letter to the Reader

If you are reading this book because you suspect your iron is low, I hope you have found validation here — confirmation that your symptoms are real, that your fatigue is not a character flaw, that the brain fog and hair loss and restless legs and ice cravings are all pointing in the same direction.

If you are reading this book because you have already been diagnosed and you are in treatment, I hope you have found practical tools that make the process clearer and less overwhelming. The road back to feeling like yourself is not always straight, but it is traveled by thousands of women every day. You are not alone on it.

If you are reading this book because a doctor told you your iron is fine and you knew in your gut that it was not, I hope you feel armed with the information you need to advocate for yourself. You were right to trust your body. Keep trusting it.

Iron deficiency is a problem that has a solution. It is not a mystery. It is not something you have to live with. The testing exists. The treatment exists. The recovery is real. The barrier is awareness — yours and your doctor's — and that is a barrier that knowledge can break down.

Here is what I want you to remember above all else:

You are not lazy. You are not making this up. You are not weak or broken or failing at being a functional adult. You may be iron deficient, and iron deficiency is a well-understood, treatable condition.

The path forward is straightforward: test, treat, recover, maintain. You have everything you need to walk that path. And on the other side of it is a version of you who has energy to spare, a clear mind, steady mood, strong hair, warm hands, and a body that feels like it is on your side again.

That version of you is not a fantasy. She is waiting. Go get your ferritin checked. You deserve to meet her.

Toolkit Summary

- **Full recovery takes 3-6 months** — hemoglobin rises first, then ferritin, then symptoms resolve over their own timeline. Functional recovery — forgetting you were ever deficient — comes last.
- **Annual ferritin testing is the cornerstone of maintenance** — catch downward trends before they become symptomatic. Set a yearly reminder.
- **Maintenance supplements may be needed long-term** — especially if the causes of your deficiency (heavy periods, absorption issues, diet) are ongoing. A lower dose is often sufficient for maintenance. Talk to your doctor about what is right for you.
- **Watch for the "I thought I was fine" trap** — it is common to crash again after feeling better. Stay aware, especially if your risk factors have not changed.
- **Build your medical team** — primary care, gynecology, hematology, gastroenterology, and dietetics all have roles to play depending on your situation. You do not have to navigate this alone.

- **Recovery is an identity shift** — when your energy and clarity return, you will need to renegotiate your expectations of yourself. That is a good thing. Give yourself time to adjust.
- **You are not broken. You were deficient. And deficiency has a fix.** The tools are here. The next step is yours.

resources

Glossary of Key Terms

FERRITIN — THE PROTEIN THAT STORES IRON IN YOUR BODY. FERRITIN IS THE BEST MEASURE OF YOUR TOTAL IRON STORES. LOW FERRITIN (<30 NG/ML) INDICATES IRON DEFICIENCY EVEN WHEN HEMOGLOBIN IS NORMAL.

Hemoglobin — The protein in red blood cells that carries oxygen from your lungs to the rest of your body. Low hemoglobin defines anemia.

Hematocrit — The percentage of your blood volume made up of red blood cells. A low hematocrit indicates anemia.

Serum Iron — The amount of iron circulating in your blood at the time of the test. Highly variable day to day.

TIBC (Total Iron-Binding Capacity) — A measure of how much iron your blood can carry. High TIBC indicates low iron stores (your body is making more transferrin to try to catch iron).

Transferrin Saturation — The percentage of iron-binding sites that are actually occupied by iron. Calculated from serum iron $\div$ TIBC. Below 20% suggests deficiency.

RDW (Red Cell Distribution Width) — A measure of variation in red blood cell size. High RDW is an early indicator of iron deficiency (your body starts making smaller, irregular cells as iron runs out).

Hepcidin — The master regulator of iron absorption. High hepcidin (caused by inflammation) blocks iron absorption. Low hepcidin allows more absorption.

Heme Iron — Iron bound to hemoglobin and myoglobin, found only in animal foods (meat, poultry, fish). Absorbed much more efficiently than non-heme iron.

Non-Heme Iron — Iron found in plant foods (spinach, beans, lentils) and fortified foods. Absorbed less efficiently and affected by other dietary components.

Menorrhagia — The medical term for heavy menstrual bleeding: periods lasting more than 7 days, soaking through a pad or tampon every hour, passing clots larger than a quarter.

Telogen Effluvium — A temporary form of hair loss triggered by physiological stress (including iron deficiency). Hair shifts from growing phase to shedding phase, typically 3 months after the trigger.

Pica — The craving of non-food substances. Ice (pagophagia) is the most common form associated with iron deficiency. Other forms include clay, dirt, paper, and starch.

Supplement Comparison Chart

Type	Elemental Iron Per Pill	Absorption	Side Effects	Best For
Ferrous Sulfate	65 mg (per 325 mg pill)	Moderate	High (constipation, nausea)	Budget option; most studied
Ferrous Fumarate	33 mg (per 100 mg pill)	Moderate	Moderate	Lower dose alternative
Ferrous Bisglycinate	25–50 mg	High (gentle on stomach)	Low	Sensitive stomachs; long-term use
Heme Iron Polypeptide	11–12 mg	High (no hepcidin block)	Very low	Best absorption; expensive
Iron Protein Succinylate	40 mg	High	Low	EU standard; harder to find in US

Dosing tip: Every-other-day dosing (e.g., 65 mg elemental iron every other day) is more effective than daily dosing because hepcidin levels return to baseline between doses, allowing better absorption.

Iron-Rich Foods Cheat Sheet

Heme sources (best absorption): - Beef liver — 5 mg per 3 oz (highest) - Oysters — 3-5 mg per 3 oz - Beef (chuck, sirloin) — 2-3 mg per 3 oz - Dark meat chicken/turkey — 1-2 mg per 3 oz - Sardines — 2 mg per 3 oz

Non-heme sources (pair with vitamin C): - Lentils — 3 mg per half cup cooked - Spinach (cooked) — 3 mg per half cup - Chickpeas — 2 mg per half cup - Pumpkin seeds — 2 mg per ounce - Fortified breakfast cereal — 4-18 mg per serving (check label) - Black beans — 2 mg per half cup - Tofu — 2 mg per half cup - Cashews — 2 mg per ounce

Absorption multipliers: Eat non-heme iron with vitamin C (citrus, bell peppers, tomatoes, strawberries). Cooking in a cast-iron pan adds 1-2 mg of iron per serving.

Absorption blockers (separate by 1-2 hours): Coffee, tea (tannins), calcium supplements/dairy, whole grains (phytates), soy.

Lab Tracker Template

Create a simple spreadsheet with columns:

Date	Ferritin	Hemoglobin	Hematocrit	Serum Iron	TIBC	TSAT %	R
------	----------	------------	------------	------------	------	--------	---

Track every 3 months during recovery, then annually for maintenance.

Medication Timing Guide

- **Iron supplements:** Take on an empty stomach (1 hour before or 2 hours after meals) with vitamin C (a glass of orange juice or a 250 mg vitamin C tablet). Separate from coffee, tea, calcium, and high-fiber meals by at least 1-2 hours.
- **Thyroid medication:** Separate from iron by at least 4 hours. Iron binds levothyroxine and prevents absorption.
- **Antibiotics (certain types):** Tetracyclines and fluoroquinolones should be separated from iron by 2-4 hours.
- **PPIs (acid reducers):** Long-term use reduces iron absorption. Talk to your doctor about monitoring your ferritin.

Further Reading

Books: - The Iron Disorders Institute Guide to Anemia — Cheryl Garrison - What to Eat When You Have Low Iron — Karen Kaye - Living With Iron — Dr. Thomas H. Bothwell (classic reference, dense) - The Complete Guide to Iron Deficiency — Dr. Robert T. Means Jr.

Websites and Organizations: - American Society of Hematology (hematology.org) — Patient education resources - National Heart, Lung, and Blood Institute (nhlbi.nih.gov) — Anemia information - Iron Disorders Institute (irondisorders.org) — Dedicated to iron-related conditions - CDC Iron and Iron Deficiency (cdc.gov) — Public health data and guidelines

Online Communities: - r/Anemia on Reddit — Active community for sharing experiences and advice - r/Ferritin on Reddit — Specifically for low ferritin discussion - The Iron Protocol on Facebook — Evidence-based iron deficiency support group

Medical clearance: Before starting any iron supplementation, talk to your physician. Iron overload (hemochromatosis) affects approximately 1 in 200 people and can cause serious organ damage. A simple blood test (ferritin and transferrin saturation) before you start supplements can rule this out. Iron is not a vitamin you excrete when you have too much — it accumulates. Getting the dose right matters, and your doctor is an essential partner in getting it right.

About the Author

Hannah Wells is a health writer who spent years battling unexplained fatigue, brain fog, and hair loss before discovering that her ferritin — a measure of iron storage — sat at 12 ng/mL, a level most labs flag as —normal. Her work focuses on translating medical research into practical guidance for women living with common, overlooked conditions. She is not a doctor, and this book is not medical advice — it is the resource she wishes someone had handed her the day she first learned what ferritin even was.

Also by AUTHOR_NAME

This is her first book.

[This page intentionally left blank]

[This page intentionally left blank]

[This page intentionally left blank]