

THE PCOS RESET

A Woman's Guide to Understanding
Polycystic Ovary Syndrome,
Rebalancing Your Hormones, and
Taking Back Control of Your Health

Hannah Wells

THE PCOS RESET

A Woman's Guide to Understanding Polycystic Ovary Syndrome,
Rebalancing Your Hormones, and Taking Back Control of Your Health

Copyright © 2026 Hannah Wells

All rights reserved.

ISBN: 978-0-000-00000-0

Published by Pink Mango Publishing

For every woman who has been told it's all in her head
—
your body was telling the truth.
This book is for learning to listen,
and for finding your way back to yourself.

Contents

01. Foreword

02. ch01

03. ch02

04. ch03

05. ch04

06. ch05

07. ch06

08. ch07

09. ch08

10. ch09

11. ch10

12. ch11

13. resources

Foreword

I WAS TWENTY-THREE YEARS OLD WHEN MY DOCTOR TOLD ME I HAD POLYCYSTIC OVARY SYNDROME.

Her words were clinical: "Your ultrasound shows multiple follicles on your ovaries. Your periods are irregular because you are not ovulating regularly. Your testosterone is elevated. This is PCOS. It is very common."

Common. The word hung in the air, landing somewhere between a diagnosis and a dismissal. If it was so common, I thought, why had I never heard a real conversation about it? Why did I have to piece together on my own, through late-night internet searches and fragmented Reddit threads, what this condition actually meant for my body, my fertility, my mental health, my entire life?

Over the next fifteen years, I learned what nobody taught me that day. PCOS is not just a fertility problem. It is not just a cosmetic problem — though the acne, the hair growth, the hair loss, the weight gain can feel like a cruelty that no one prepared you for. It is a metabolic condition. It is a hormonal condition. It is an inflammatory condition. It affects your insulin, your blood sugar, your cholesterol, your risk of diabetes, your risk of heart disease, your mood, your energy, your sleep, your relationship with food, your relationship with exercise, your relationship with yourself.

It is also the most common endocrine disorder in women of reproductive age, affecting one in ten women worldwide. And yet the standard medical advice — "just lose weight" — has

been handed down like a magic spell for decades, as if weight loss alone could fix a condition that is fundamentally about how your body processes insulin and produces hormones.

Here is what nobody told me about PCOS: it is treatable. Not curable, but treatable — and manageable in ways that go far beyond weight loss. The right nutrition approach (which is not starvation or extreme restriction), the right movement approach (which is not punishing cardio), the right medical support (which is more than just birth control), the right supplements (which actually have evidence behind them) — all of these can shift your symptoms, improve your metabolic health, and help your body function the way it should.

The last decade has brought a flood of new research on PCOS. We now understand the central role of insulin resistance in driving the condition. We have evidence-supported nutrition and supplement protocols. We know that metformin, inositol, and GLP-1 agonists (yes, the same class of drugs that changed weight-loss treatment) can be powerful tools for PCOS management. We know that lifestyle changes can be as effective as medication for many women. And we know that the old advice — eat less, exercise more, come back when you want to get pregnant — was never sufficient.

This book is what I wish someone had handed me the day I got that ultrasound report. It covers what PCOS actually is, the insulin connection that most doctors skip, every symptom and what it means, how to get the right diagnosis and find a doctor who takes you seriously, how to eat and move and supplement in ways that actually help, how to navigate

fertility and pregnancy, and how the condition affects your mental health — because the emotional toll of PCOS is real and it deserves to be named.

Every chapter ends with a toolkit summary you can use immediately.

I am not a doctor. I am a health writer who has lived with PCOS for fifteen years and spent that time researching and writing about the conditions that affect women differently. This book is a guide, not a prescription. Work with your doctor on your treatment plan — and if your doctor tells you to "just lose weight" without offering you real support, this book will help you find a better conversation.

You are not broken. Your body is not failing you. PCOS is a real, complex condition — and there is a real path through it.

Let us walk it together.

Toolkit Summary

- **PCOS is more common than you think** — affecting 1 in 10 women worldwide, it is the most common endocrine disorder in women of reproductive age.
- **It is not just a fertility problem** — PCOS is a metabolic, hormonal, and inflammatory condition that affects every system in your body.
- **The standard advice — just lose weight — is not enough** — proper management requires nutrition, movement, medication, supplements, and mental health support.

- **Treatment exists beyond birth control** — metformin, inositol, GLP-1 medications, and targeted lifestyle changes can all help.
- **I am not a doctor** — this book is a companion, not a prescription. Work with your doctor on your treatment plan.
- **You are not broken** — your diagnosis is real, your struggles are real, and there is a path forward that does not require you to shrink yourself to be worthy of care.

ch01

IF YOU ARE READING THIS BOOK, THERE IS A GOOD CHANCE YOU HAVE EXPERIENCED THAT MOMENT — THE ONE WHERE YOU ARE SITTING IN A DOCTOR'S OFFICE BEING TOLD YOU HAVE "PCOS" AND WALKING OUT WONDERING WHAT THAT EVEN MEANS. MAYBE YOU WERE HANDED A PAMPHLET. MAYBE YOU WERE TOLD TO LOSE WEIGHT AND COME BACK. MAYBE THE DOCTOR SEEMED TO THINK IT WAS NO BIG DEAL.

But you know it is a big deal. The irregular periods, the unexplained weight gain around your midsection, the acne that came back at twenty-five like a particularly aggressive teenager, the hair growing where you do not want it and falling out where you do. You know something is off with your body, and you have probably been told — by doctors, by family, by the internet — that this is somehow your fault. That you just need to "eat less and move more." That your hormones are fine. That you are exaggerating.

You are not exaggerating.

Polycystic Ovary Syndrome (PCOS) affects an estimated one in ten women of reproductive age worldwide. That makes it the most common endocrine disorder among women. And yet, despite being this prevalent, it remains one of the most misunderstood, underdiagnosed, and poorly managed conditions in all of medicine. The average time to diagnosis is between two and four years, and many women

see multiple specialists — a gynecologist, a dermatologist, an endocrinologist, maybe a fertility specialist — before anyone connects the dots.

This chapter is going to give you the foundation you need to understand what PCOS actually is, what is happening in your body, and why so many women with this condition have been let down by the medical system.

The Diagnostic Criteria: What Actually Counts as PCOS

PCOS is diagnosed using what is called the Rotterdam criteria, established in 2003 by an international consensus of experts. To receive a PCOS diagnosis, you need to meet at least two of the following three criteria:

1. Irregular or absent ovulation (oligomenorrhea or amenorrhea). This means you have fewer than eight menstrual periods per year, or your cycles are consistently longer than 35 days. Some women with PCOS have no periods at all. Others bleed irregularly and unpredictably — sometimes twice a month, sometimes not for four months.

2. Clinical or biochemical signs of hyperandrogenism. "Hyperandrogenism" is a mouthful, but it simply means too many androgens — the group of hormones that includes testosterone. Clinically, this shows up as acne (especially along the jawline and chin), hirsutism (excess hair growth on the face, chest, back, or abdomen),

and androgenic alopecia (male-pattern hair thinning). Biochemically, it means elevated testosterone, DHEA-S, or androstenedione on blood work.

3. Polycystic ovaries on ultrasound. This means the ovaries appear enlarged and contain twelve or more small follicles (2-9 mm in diameter) on each ovary, or the ovarian volume is greater than 10 mL. Polycystic means "many cysts," but these are not harmful cysts — they are immature follicles that failed to develop and release an egg.

Importantly, the Rotterdam criteria require that other conditions be ruled out first. This includes thyroid disorders, elevated prolactin, congenital adrenal hyperplasia (specifically non-classic 21-hydroxylase deficiency), and Cushing's syndrome — all of which can mimic PCOS.

The Four Phenotypes: Why PCOS Looks Different on Everyone

This is one of the most important things to understand about PCOS: it is not one condition — it is four. Researchers have identified four distinct phenotypes based on the Rotterdam criteria:

Phenotype A (the "classic" PCOS): Irregular periods + hyperandrogenism + polycystic ovaries. This is the most severe form and carries the highest risk for metabolic problems like insulin resistance.

Phenotype B (the "non-PCO" PCOS): Irregular periods + hyperandrogenism, but normal ovaries on ultrasound. This group also has high metabolic risk.

Phenotype C (the "ovulatory" PCOS):
Hyperandrogenism + polycystic ovaries, but regular periods. These women may ovulate normally but still have elevated androgens and all the symptoms that come with them — acne, hair loss, hair growth. This is the phenotype most commonly missed by doctors who assume regular periods mean no PCOS.

Phenotype D (the "normoandrogenic" PCOS):
Irregular periods + polycystic ovaries, but normal androgen levels. These women have cycle irregularities and ovarian changes but no visible signs of elevated androgens. This group is the most likely to be told "everything looks fine" when it is not.

Knowing your phenotype matters because it shapes your treatment approach and your long-term health risks. Phenotypes A and B carry higher metabolic risk. Phenotype C is often the most responsive to treatment. Phenotype D is the hardest to diagnose because the blood work looks normal.

The Hormonal Underpinnings: What Is Actually Going Wrong

To understand PCOS, you need to understand the hormonal conversation happening in your body every day. In a normal cycle, the hypothalamus sends gonadotropin-releasing hormone (GnRH) to the pituitary gland, which then releases follicle-stimulating hormone (FSH) and luteinizing hormone (LH). FSH tells the ovaries to mature a follicle. LH triggers ovulation. It is a carefully choreographed sequence of signals.

In PCOS, this signal goes off track. The pulse frequency of GnRH speeds up, which causes the pituitary to release more LH and less FSH. This elevated LH-to-FSH ratio — often 2:1 or higher — is a hallmark of PCOS. With too little FSH, ovarian follicles cannot mature properly. They start developing but stall out, never reaching the point of ovulation. Instead of releasing an egg, each stalled follicle becomes a tiny cyst on the ovary.

Meanwhile, the excess LH stimulates the ovaries to produce more androgens. These androgens — testosterone, androstenedione, DHEA-S — create the symptoms you can see and feel: acne, hirsutism, hair loss, and abdominal weight gain. They also create a vicious cycle, because androgens further disrupt the GnRH pulse, which leads to even more LH, which leads to even more androgen production.

This is not a condition of "bad ovaries." It is a condition of disrupted hormone signaling.

The Insulin Connection: Why PCOS Is Metabolic, Not Just Reproductive

Here is the piece most doctors miss: PCOS is fundamentally a metabolic condition, not just a reproductive one. The reproductive symptoms — the irregular periods, the infertility, the acne — are downstream effects of a metabolic problem.

Approximately 70 percent of women with PCOS have insulin resistance, regardless of their body weight. Insulin resistance means your cells do not respond properly to

insulin, the hormone that moves glucose out of your bloodstream and into your cells for energy. To compensate, your pancreas pumps out more insulin. This is called compensatory hyperinsulinemia — high levels of insulin circulating in your blood.

And here is where it gets relevant to PCOS: insulin stimulates the ovaries to produce androgens. Directly. Theca cells in the ovaries have insulin receptors, and when excess insulin binds to them, they ramp up testosterone production. At the same time, insulin suppresses the production of sex hormone-binding globulin (SHBG) in the liver. SHBG is the protein that normally soaks up free testosterone in your blood. When SHBG levels are low, more testosterone stays active and unbound.

So the metabolic problem (insulin resistance) drives the hormonal problem (hyperandrogenism), which drives the reproductive and cosmetic symptoms. This is the central loop of PCOS, and understanding it changes everything about how you approach treatment.

If you address the insulin resistance — through nutrition, movement, medications, or supplements — you break the loop. Lower insulin means lower androgen production, which means more normal SHBG levels, which means fewer symptoms, which means more regular ovulation.

Why Women with PCOS Get Misdiagnosed and Dismissed

There is an uncomfortable truth about PCOS: women with this condition are routinely dismissed by the medical system. Study after study has found that women with PCOS report feeling "not taken seriously," "blamed for their condition," and "frustrated by the lack of useful information" from healthcare providers.

Part of this is the weight bias that pervades medicine. Because PCOS is strongly associated with weight gain and difficulty losing weight, many doctors default to telling patients to "just lose weight" without addressing the underlying hormonal and metabolic barriers that make weight loss difficult with PCOS. This is not helpful. It is actively harmful.

Part of it is the siloed nature of modern medicine. A dermatologist treating your acne may not check your hormone levels. A gynecologist treating your irregular periods may not screen your insulin. An endocrinologist managing your blood sugar may not ask about your menstrual cycle. No one is connecting the dots because no one is looking at the whole picture.

And part of it is that PCOS simply does not fit the "sick patient" mold. You are not obviously diseased. You are walking around, functioning, showing up to work — just struggling in ways that are invisible. This invisibility makes it easy for doctors to minimize what you are experiencing.

The Reality of Living with a Condition Nobody Explains Well

If you have PCOS, you have probably been told a lot of things that are not true. That you cannot get pregnant (usually false — PCOS is highly treatable). That you are making yourself sick by eating the wrong foods (false — PCOS makes certain foods harder to process). That your only option is birth control pills (false — there are many treatment options). That you should just accept your symptoms (absolutely false — you deserve better than that).

Here is the truth: PCOS is a chronic condition. It will not go away. But it is also one of the most manageable chronic conditions in existence. With the right approach — the kind this book will help you build — most women with PCOS see dramatic improvements in their symptoms, their metabolic health, and their quality of life.

You are not broken. Your metabolism is not broken. Your hormones are not permanently out of whack. There is a specific mechanism driving your symptoms, and once you understand it, you can do something about it.

This book is going to walk you through everything: the insulin connection, the symptom map, the diagnostic journey, the nutrition strategies that actually work, the movement that helps, the medications and supplements with real evidence behind them, the mental health piece that nobody talks about, and the long-term plan that keeps you healthy for decades.

But before all of that, you need to hear this: PCOS is not your fault. You did not cause this by eating too much sugar or not exercising enough or being too stressed. PCOS is a genetic, metabolic, and hormonal condition with deep biological roots. Your job is not to blame yourself. Your job is to learn how to manage it.

And you can. You absolutely can.

Getting Started with Your PCOS Journey

If you have just been diagnosed — or if you have been living with PCOS without understanding it until now — the beginning can feel overwhelming. There is so much information, so many things to track, so many appointments to schedule. Start small. Pick one thing from the next seven chapters and try it for two weeks. Maybe it is tracking your cycles. Maybe it is adding a 15-minute walk after dinner. Maybe it is swapping your breakfast cereal for eggs and vegetables. One change at a time compounds into significant progress.

The women who manage PCOS best are not the ones who do everything perfectly. They are the ones who stay consistent enough, stay curious enough, and forgive themselves when they fall off track. Perfection is not the goal. Progress is.

Toolkit Summary

- PCOS is diagnosed using the Rotterdam criteria (2 of 3): irregular ovulation, hyperandrogenism, or polycystic ovaries on ultrasound
- There are four phenotypes of PCOS (A, B, C, D), each with different symptom profiles and metabolic risks
- PCOS is fundamentally a metabolic disorder driven by insulin resistance in approximately 70 percent of women
- The elevated LH-to-FSH ratio and compensatory hyperinsulinemia drive ovarian androgen production
- Women with PCOS are routinely misdiagnosed and dismissed, with an average 2–4 year delay in diagnosis
- Understanding the insulin-androgen feedback loop is the key to effective management
- PCOS is highly treatable — most women see significant improvement with the right comprehensive approach
- If you suspect you have PCOS or have been diagnosed but feel lost, talk to your doctor about a full workup

ch02

IF YOU HAVE PCOS, YOU HAVE PROBABLY BEEN TOLD — DIRECTLY OR INDIRECTLY — THAT YOUR METABOLISM IS BROKEN. MAYBE YOUR DOCTOR IMPLIED THAT YOU MUST BE EATING TOO MUCH. MAYBE A FAMILY MEMBER SUGGESTED THAT IF YOU JUST TRIED HARDER, THE WEIGHT WOULD COME OFF. MAYBE YOU HAVE INTERNALIZED THE MESSAGE YOURSELF, WONDERING WHY EVERYONE ELSE SEEMS TO BE ABLE TO EAT A NORMAL AMOUNT OF FOOD WITHOUT THEIR BODY REBELLING.

Here is what is actually happening: your metabolism is not broken, but it is stuck. And the mechanism that keeps it stuck is insulin.

This chapter is going to explain insulin resistance — what it is, how it drives PCOS, and why the classic advice of "eat less, move more" completely misses the point. By the end, you will understand the single most important concept in managing PCOS: the insulin-androgen feedback loop.

Insulin 101: What It Does and How It Works

Insulin is a hormone produced by your pancreas, specifically by beta cells in the islets of Langerhans. Every time you eat food that contains carbohydrates, your blood glucose (blood sugar) rises. The pancreas detects this rise and releases insulin into your bloodstream.

Insulin acts like a key. It travels through your blood and binds to insulin receptors on the surface of your cells — primarily muscle cells, fat cells, and liver cells. When insulin binds to these receptors, it unlocks the cell, allowing glucose to move from your bloodstream into the cell, where it can be used for energy.

That is insulin's main job, but it is not its only job. Insulin also signals your liver to store glucose as glycogen for later use, tells your fat cells to store fat, and instructs your liver to stop producing glucose when levels are already high enough.

In a body with normal insulin sensitivity, this system works beautifully. You eat, glucose rises, insulin opens the door, glucose enters your cells, and your blood sugar returns to a healthy baseline. This is the metabolic equivalent of a well-run train station: passengers arrive, check in, board the train, and the platform clears for the next arrival.

What Goes Wrong in PCOS: Insulin Resistance

Now imagine that same train station, but the doors are jammed. Passengers are piling up on the platform (in your bloodstream), and the station manager (your pancreas) keeps calling for more trains (more insulin) because the platform is not clearing.

This is insulin resistance. Your cells have become less responsive to insulin's signal. The "doors" are not opening properly. Glucose stays in your bloodstream longer than it

should. Your pancreas, seeing that blood sugar is not coming down, pumps out more and more insulin to force the doors open.

This is compensatory hyperinsulinemia — high levels of insulin in your blood. It is a compensatory mechanism. Your body is trying to solve a problem. But this high-insulin state creates a whole new set of problems, especially if you have PCOS.

Insulin resistance is extremely common in PCOS. Studies consistently find that somewhere between 50 and 80 percent of women with PCOS have insulin resistance, depending on which diagnostic criteria are used and which PCOS phenotype is present. The rates are highest in women with phenotypes A and B — those with both irregular ovulation and hyperandrogenism.

Critically, insulin resistance in PCOS is not limited to women who are overweight or obese. Lean women with PCOS also have higher rates of insulin resistance compared to women without PCOS. This is known as "lean PCOS," and it is a real and underrecognized phenomenon. If you are thin and wondering why you are being told you have a metabolic condition, this is why.

How Insulin Drives Androgens: The Vicious Cycle

Here is the piece that connects the metabolic to the hormonal. Insulin, at high levels, stimulates the ovaries to produce androgens. This happens through two mechanisms.

First, direct stimulation. The theca cells of the ovaries have insulin receptors and IGF-1 receptors. When excess insulin binds to these receptors, it activates enzyme pathways (specifically 17 α -hydroxylase and 17,20-lyase) that convert cholesterol into androstenedione and testosterone. More insulin means more androgen production.

Second, the SHBG connection. Your liver produces a protein called sex hormone-binding globulin (SHBG). SHBG circulates in your blood and binds to testosterone, estrogen, and other sex hormones, keeping them inactive. When testosterone is bound to SHBG, it is not causing acne, hair loss, or irregular periods — it is just sitting there, harmless.

Insulin suppresses the liver's production of SHBG. So when insulin is high, SHBG is low. When SHBG is low, more of your testosterone is "free" — unbound and active. Even if your total testosterone levels are within the normal range, your free testosterone can be elevated simply because there is not enough SHBG to carry it.

This is why some women with PCOS have "normal" testosterone on blood work but still have all the symptoms of hyperandrogenism. The total number looks fine, but the active fraction is too high.

So here is the feedback loop: insulin resistance → high insulin → high ovarian androgen production + low SHBG → high free androgens → PCOS symptoms. And because androgens themselves can worsen insulin resistance, the cycle feeds on itself.

Metabolic Syndrome and PCOS: The Long-Term Health Risks

Because insulin resistance is so central to PCOS, women with this condition are at significantly higher risk for developing metabolic syndrome — a cluster of conditions that occur together and increase the risk of heart disease, stroke, and type 2 diabetes.

Metabolic syndrome is diagnosed when you have three or more of the following: - Abdominal obesity (waist circumference over 35 inches for women) - Elevated triglycerides (150 mg/dL or higher) - Low HDL cholesterol (under 50 mg/dL for women) - High blood pressure (130/85 mmHg or higher) - Elevated fasting blood glucose (100 mg/dL or higher)

Women with PCOS have a two-to-three times higher risk of developing metabolic syndrome compared to women without PCOS. They also have a significantly higher risk of impaired glucose tolerance and type 2 diabetes — and this risk is independent of body weight. Lean women with PCOS have higher rates of diabetes than lean women without PCOS.

There is also a strong link between PCOS and non-alcoholic fatty liver disease (NAFLD). Insulin resistance drives fat accumulation in the liver, and studies suggest that up to 50 to 70 percent of women with PCOS may have NAFLD. This is something your doctor should be monitoring with liver function tests, even if you are not experiencing any symptoms.

The Inflammation Connection

Insulin resistance is not just about glucose and insulin. It is also about inflammation. Adipose tissue (fat tissue) in an insulin-resistant state releases inflammatory cytokines — chemical messengers that promote a state of low-grade, chronic inflammation throughout the body. This is not the acute inflammation of an injury or infection. It is a slow, smoldering background inflammation that damages cells and tissues over time.

This chronic inflammation worsens insulin resistance, which worsens hyperandrogenism, which worsens PCOS symptoms. It is another feedback loop, layered on top of the insulin-androgen loop.

Women with PCOS tend to have elevated markers of inflammation, including C-reactive protein (CRP), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNF- α). This inflammation contributes to the metabolic risks of PCOS, but it also contributes to the more subjective symptoms — the fatigue, the brain fog, the general feeling of being unwell that many women with PCOS describe.

GLP-1 Agonists and PCOS: The New Frontier

In recent years, a class of medications called GLP-1 receptor agonists has emerged as a promising treatment for PCOS, particularly for women with insulin resistance and metabolic complications. GLP-1 (glucagon-like peptide-1) is a

naturally occurring hormone that stimulates insulin secretion, suppresses glucagon release, slows gastric emptying, and promotes satiety.

Medications like semaglutide (Ozempic, Wegovy) and tirzepatide (Mounjaro, Zepbound) are GLP-1 agonists — they mimic the action of GLP-1. Studies in women with PCOS have found that these medications improve insulin sensitivity, promote significant weight loss, reduce androgen levels, and can even restore regular ovulation in some women.

This is an exciting development, but it comes with important caveats. These are prescription medications that should only be used under a doctor's supervision. They have side effects, primarily gastrointestinal, and they can be expensive. They are not a "quick fix" — most women need to stay on them long-term to maintain benefits, and they work best when combined with nutrition and lifestyle changes.

If you are interested in GLP-1 agonists for PCOS, talk to your doctor. There are specific prescribing criteria, and not everyone is a candidate.

Why "Eat Less" Does Not Solve the Insulin Problem

If there is one message I want you to take away from this chapter, it is this: the "eat less, move more" approach does not work for insulin resistance because it misunderstands the problem.

When your cells are insulin resistant, they are not responding to normal insulin signals. Restricting calories does not fix insulin resistance. It may temporarily lower your weight, but if the underlying metabolic dysfunction is not addressed, the weight will return and the insulin will remain high.

Furthermore, aggressive calorie restriction can actually worsen insulin resistance in the short term by raising cortisol and triggering a stress response that mobilizes glucose. Your body interprets food scarcity as a threat and responds by making more glucose available — exactly what you do not want.

The solution is not to eat less food. It is to eat food that does not spike your insulin, to eat at regular intervals that keep blood sugar stable, and to address the underlying mechanisms through the combination of nutrition, movement, medications, and supplements discussed in the coming chapters.

Your metabolism is not broken. It is stuck. And now you know the mechanism that keeps it from moving. That knowledge is the first and most important step toward unsticking it.

Insulin Resistance Testing: What to Ask Your Doctor For

If you suspect you have insulin resistance, the standard screening tests are a fasting insulin level and a fasting glucose level drawn together. Many doctors will tell you that

a "normal" fasting glucose rules out insulin problems — but this is incorrect. Your glucose can be perfectly normal while your insulin is high, because your pancreas is working overtime to keep your blood sugar in check. This is early-stage insulin resistance, and it is the stage where intervention is most effective.

An oral glucose tolerance test (OGTT) with insulin measurements is more sensitive than a simple fasting test. In this test, you drink a sugar solution and have blood drawn at 0, 60, and 120 minutes to see how your glucose and insulin respond. A prolonged, exaggerated insulin response on the OGTT is a hallmark of insulin resistance. If your doctor is willing to order this test, it provides much more information than fasting values alone.

Some practitioners also use the HOMA-IR (Homeostatic Model Assessment for Insulin Resistance) calculation, which uses your fasting insulin and glucose to estimate insulin resistance. A HOMA-IR score above 2.5 indicates significant insulin resistance. You can calculate this online and discuss the result with your doctor.

Toolkit Summary

- Insulin resistance affects 50–80 percent of women with PCOS and is the primary driver of symptoms
- High insulin stimulates ovarian androgen production and suppresses SHBG, creating the insulin-androgen feedback loop
- PCOS carries elevated risk for metabolic syndrome, type 2 diabetes, NAFLD, and cardiovascular disease

- Chronic inflammation worsens insulin resistance and contributes to fatigue and brain fog
- GLP-1 agonists are an emerging treatment for PCOS with metabolic complications — talk to your doctor if you are interested
- "Eat less, move more" does not fix insulin resistance — targeted metabolic interventions are needed
- Understanding the insulin connection is the foundation of every effective PCOS management plan
- Work with your doctor to assess your insulin resistance through fasting insulin, glucose, and HbA1c testing

ch03

YOUR MENSTRUAL CYCLE IS THE MOST BASIC MEASURE OF REPRODUCTIVE HEALTH. IT IS ALSO, FOR MANY WOMEN WITH PCOS, THE MOST VISIBLE SIGN THAT SOMETHING IS WRONG. MAYBE YOUR PERIODS COME EVERY FEW MONTHS, COMPLETELY UNPREDICTABLY. MAYBE THEY SKIP AN ENTIRE SEASON. MAYBE THEY COME TOO OFTEN — EVERY TWO WEEKS, HEAVY AND PAINFUL. MAYBE THEY HAVE STOPPED ENTIRELY.

Whatever your pattern looks like, it is telling you something about what is happening inside your body. This chapter will help you understand what a healthy cycle looks like, what PCOS does to that cycle, and what your options are if you are hoping to conceive.

What a Normal Cycle Actually Looks Like

Before we talk about what goes wrong, let's establish what "normal" means for a menstrual cycle. A typical cycle ranges from 21 to 35 days, from the first day of one period to the first day of the next. The average is about 28 days, but anything in that 21-to-35 range is considered normal.

The cycle has two main phases:

The follicular phase begins on day one of your period. The pituitary gland releases FSH (follicle-stimulating hormone), which tells the ovaries to recruit and mature a follicle containing an egg. This phase lasts about 10 to 14 days in a normal cycle.

Ovulation occurs when a mature surge of LH (luteinizing hormone) triggers the release of the egg from the ovarian follicle. This typically happens around day 14 in a 28-day cycle.

The luteal phase follows ovulation. The ruptured follicle transforms into the corpus luteum, which produces progesterone to prepare the uterine lining for a potential pregnancy. If no pregnancy occurs, the corpus luteum breaks down, progesterone drops, and your uterine lining sheds as your period. The luteal phase is relatively fixed at 12 to 16 days — it does not vary much from person to person.

This means that if your cycles are consistently longer than 35 days, the delay is almost certainly in the follicular phase. You are taking longer to develop and release an egg. And if you are not ovulating at all, you may still have bleeding — but it is not a true period. It is breakthrough bleeding caused by estrogen building up the uterine lining until it becomes unstable and sheds on its own.

What PCOS Does to Your Cycle: Anovulation

The hallmark of PCOS-related cycle disruption is anovulation — the absence of ovulation. Without ovulation, there is no corpus luteum, no progesterone production, and no predictable cycle.

Instead, the PCOS cycle looks like this: The ovaries try to develop follicles, but the elevated LH-to-FSH ratio and the hyperinsulinemia stall the process. Follicles start growing but never mature enough to ovulate. They become the small, fluid-filled structures that appear as "cysts" on an ultrasound — technically, they are immature follicles.

Without ovulation, the ovaries continue producing estrogen, but not progesterone. This unopposed estrogen builds up the uterine lining (the endometrium) thicker and thicker. Eventually, the lining becomes unstable and sheds irregularly. This is why PCOS periods are often unpredictable, heavy, and sometimes prolonged.

Over time, prolonged anovulation carries a risk: the constant buildup of the endometrium without regular shedding can lead to endometrial hyperplasia, a precancerous condition. This is why it is important for women with PCOS who are not on birth control to have at least four periods per year — either naturally or induced with medication. Talk to your doctor about this if you are going more than three months without a period.

How to Tell If You Are Ovulating

You do not have to guess whether you are ovulating. There are reliable signs you can track at home:

Cervical mucus changes. As ovulation approaches, estrogen rises and cervical mucus becomes clear, slippery, and stretchy — like raw egg white. This is fertile-quality

mucus. After ovulation, progesterone thickens the mucus, making it sticky and opaque. If you never see egg-white mucus, you may not be ovulating.

Basal body temperature (BBT). Your body temperature rises by about 0.3 to 0.5 degrees Fahrenheit after ovulation, thanks to progesterone's thermogenic effect. If you take your temperature every morning before getting out of bed (same time, after at least three hours of sleep), you will see a sustained temperature shift after ovulation. If BBT stays flat across your cycle, you are likely not ovulating.

Ovulation predictor kits (OPKs). These detect the LH surge that precedes ovulation. However, women with PCOS often have chronically elevated LH levels, which can cause false positives. OPKs may show "positive" multiple times in a cycle or never show a clear surge. They are not always reliable for PCOS.

Progesterone testing. Your doctor can measure serum progesterone on day 21 of a 28-day cycle (or day 7 before expected period). A level above 3 ng/mL confirms ovulation. For women with irregular cycles, the timing is trickier — your doctor may ask you to come in for weekly blood draws until ovulation is confirmed.

Fertility and PCOS: The Most Treatable Cause of Infertility

Here is the most encouraging thing I can tell you about PCOS and fertility: PCOS is the most common cause of ovulatory infertility, and it is also the most treatable. The vast

majority of women with PCOS — somewhere between 70 and 80 percent — can achieve pregnancy with the right treatment.

The primary barrier to fertility in PCOS is anovulation. If you are not releasing eggs, you cannot conceive. But the eggs themselves are typically healthy. The problem is not your eggs; it is the hormonal environment that prevents them from being released. When you address that environment — through ovulation induction medications, insulin-sensitizing treatments, or even lifestyle changes — ovulation resumes in many women.

This is an important reframe. PCOS-related infertility is not the same as primary ovarian insufficiency or diminished ovarian reserve. Your ovarian reserve is often normal or even higher than average, because the follicles accumulate over cycles without being released. The problem is mechanical-hormonal, not about egg quantity or quality — though insulin resistance and inflammation can affect egg quality over time.

Ovulation Induction: Letrozole vs. Clomiphene

When a woman with PCOS is ready to conceive and not ovulating on her own, the first-line treatment is ovulation induction with medication. Two drugs are commonly used.

Letrozole (Femara) is now considered the first-line ovulation induction agent for PCOS, based on a landmark 2014 study (the AMIGOS trial) that found letrozole resulted in higher live birth rates and lower rates of multiple

pregnancy compared to clomiphene. Letrozole is an aromatase inhibitor — it temporarily reduces estrogen production, which tricks the pituitary into releasing more FSH, which stimulates follicle growth. It is taken orally on cycle days 3 through 7.

Clomiphene citrate (Clomid) was the traditional first-line drug. It is a selective estrogen receptor modulator that blocks estrogen receptors in the hypothalamus, also increasing FSH release. Clomiphene is effective but has a higher risk of multiple pregnancy (10 percent twin rate vs. about 3 percent with letrozole) and can thin the uterine lining, which may reduce implantation rates in some women.

Letrozole is preferred for most women with PCOS, but clomiphene remains a viable option, particularly if cost or availability is a concern. Both are oral medications — no injections, no daily clinic visits. Your doctor should monitor with ultrasound and blood work to track follicle development and time intercourse appropriately.

The Fertility Preservation Conversation

For women with PCOS who are not ready to conceive but worry about their future fertility, here is some reassurance: PCOS does not typically cause early ovarian aging. Your ovarian reserve is usually not diminished. However, age still matters. Egg quality declines with age for everyone, including women with PCOS.

If you are in your twenties or early thirties with PCOS and do not plan to have children for several years, you do not usually need to rush into egg freezing. Your fertility

prognosis with treatment is generally excellent. But if you are in your late thirties or have additional risk factors (endometriosis, family history of early menopause), a conversation with a reproductive endocrinologist about your options may be helpful.

IVF and PCOS: Higher Success but Also Higher Risk

For women with PCOS who need in vitro fertilization (IVF) — either because ovulation induction has not worked, or because there are additional fertility factors at play — the prognosis is generally good. Women with PCOS tend to produce more eggs per retrieval cycle compared to women with normal ovaries, thanks to their higher antral follicle count. More eggs often mean more embryos, which improves the chances of pregnancy.

However, there is a significant risk that comes with this egg abundance: ovarian hyperstimulation syndrome (OHSS). OHSS is a potentially serious complication in which the ovaries become swollen and painful after medication stimulation, and fluid leaks into the abdominal cavity. Women with PCOS are at higher risk for OHSS because their ovaries respond more vigorously to stimulation medications.

Your reproductive endocrinologist can minimize this risk by using lower medication doses, using a GnRH agonist trigger shot instead of hCG, and considering a "freeze-all" approach where embryos are frozen for later transfer rather than transferred fresh.

The Emotional Weight of Fertility Challenges

It is important to acknowledge the emotional toll of fertility challenges with PCOS. Fertility is rarely mentioned in the same sentence as the word "sadness" in medical texts, but it should be. The month-after-month disappointment of negative pregnancy tests, the invasive nature of fertility treatments, the strain it puts on relationships, and the grief of feeling like your body is not doing what it is supposed to do — this is real and it is heavy.

If you are on a fertility journey with PCOS, you deserve support. That may mean a therapist who specializes in reproductive health, a support group (online or in-person), or simply a partner or friend who can sit with you in the hard moments without trying to fix everything.

And remember: the statistics are on your side. Most women with PCOS who want to conceive eventually do. It just may take longer, and it may require more intervention, than it does for women without PCOS.

When to See a Reproductive Specialist

You do not necessarily need a reproductive endocrinologist right away. Many women with PCOS ovulate with simple lifestyle changes or with medication from their regular gynecologist. However, there are situations where seeing a specialist earlier rather than later can save you time, money, and emotional energy.

Consider a referral to a reproductive endocrinologist if: you have been trying to conceive for six months with regular ovulation on medication but have not become pregnant, you are over 35 and trying to conceive for six months with ovulation induction, you have additional fertility factors (male factor, tubal issues, endometriosis), or you have tried three to six ovulatory cycles on oral medications without success. Reproductive endocrinologists have access to a broader range of treatments, including injectable medications, intrauterine insemination (IUI), and IVF.

If you are under 35 and ovulating on medication without other known fertility factors, many fertility specialists recommend trying for twelve ovulatory cycles before moving to more advanced treatment. The monthly chance of conception in a healthy couple is only about 20 to 25 percent, so patience is built into the process even under ideal circumstances.

Toolkit Summary

- A normal cycle is 21-35 days with a predictable follicular phase, ovulation, and luteal phase
- PCOS causes anovulation — irregular or absent ovulation — due to elevated LH and hyperinsulinemia
- Track ovulation with cervical mucus, BBT, OPKs (with caution), or progesterone blood testing
- PCOS is the most common cause of ovulatory infertility and also the most treatable — 70-80 percent of women achieve pregnancy with treatment

- Letrozole is first-line for ovulation induction; clomiphene is an alternative
- IVF success rates are generally good for PCOS, but OHSS risk is higher — talk to your doctor about prevention strategies
- If you are going more than three months without a period, talk to your doctor about protecting your endometrial health
- Fertility challenges are emotionally heavy — seek support and remember that most women with PCOS do eventually conceive

ch04

PCOS DOES NOT PRESENT THE SAME WAY IN ANY TWO WOMEN, AND THAT IS ONE OF THE THINGS THAT MAKES IT SO FRUSTRATING TO DIAGNOSE AND SO CONFUSING TO LIVE WITH. YOUR SISTER WITH PCOS MAY HAVE COMPLETELY DIFFERENT SYMPTOMS THAN YOU DO. YOUR FRIEND FROM THE SUPPORT GROUP MAY BE DEALING WITH THINGS YOU HAVE NEVER EXPERIENCED.

But despite this variability, there are common symptom patterns — clusters of signs and symptoms that point to the underlying hormonal and metabolic drivers. This chapter maps the full symptom landscape of PCOS so you can recognize what is PCOS, what is something else entirely, and how your specific pattern should guide your treatment approach.

The Skin Symptoms: Acne and Hyperandrogenism

Acne in PCOS has a signature pattern. It is not the forehead-and-nose acne of adolescence. PCOS acne is jawline and chin acne — the "beard distribution." It is deep, cystic, and painful. It tends to flare in the week before bleeding (if you are having periods), and it often does not respond well to over-the-counter treatments.

The cause is elevated androgens. Androgens stimulate sebaceous glands to produce more sebum — the oily substance that clogs pores. They also promote the growth of *Cutibacterium acnes*, the bacteria that contribute to inflamed acne. When you add in the chronic inflammation that comes with insulin resistance, you have a perfect recipe for persistent, treatment-resistant breakouts.

Standard acne treatments may help but often fall short for PCOS-related acne unless the underlying androgen excess is addressed. That is why topical retinoids and benzoyl peroxide may not be enough, and why targeting the root cause — elevated androgens and insulin — makes a bigger difference. For some women, spironolactone (a medication that blocks androgen receptors) combined with a good skincare routine can dramatically improve PCOS acne. Talk to your doctor about whether this is right for you.

Hirsutism: When Hair Grows Where It Should Not

Hirsutism is the medical term for excess hair growth in a male pattern — on the face (upper lip, chin, sideburns), chest, back, shoulders, abdomen (from the pubic area up toward the navel), and inner thighs. It affects approximately 60 to 80 percent of women with PCOS who have elevated androgens.

The mechanism is straightforward: androgens convert vellus hairs (the fine, barely visible hairs that cover most of your body) into terminal hairs (thick, dark, visible hairs). This is the same process that happens in puberty when hair

appears in the armpits and pubic area — but in PCOS, it happens in places where women typically do not grow thick hair.

Hirsutism is one of the most distressing PCOS symptoms for many women, and it is also one of the slowest to respond to treatment. Even when your androgen levels come down, existing terminal hairs do not simply disappear. They may become finer and grow more slowly, but they remain. That is why a combination approach works best: medical treatment to lower androgens (spironolactone, birth control, insulin-sensitizing medications) combined with direct hair removal methods (electrolysis, laser, threading, shaving). Talk to your doctor about the medical options; a licensed electrologist or laser technician can help with the cosmetic side.

Androgenic Alopecia: Hair Loss in PCOS

Androgenic alopecia is the medical name for male-pattern hair thinning, and it affects many women with PCOS. Unlike the diffuse hair loss that can happen with thyroid conditions, iron deficiency, or stress, androgenic alopecia follows a distinct pattern: thinning at the crown of the head and widening of the center part. The front hairline typically remains intact.

This happens because androgens (specifically dihydrotestosterone, or DHT) bind to androgen receptors on hair follicles, causing them to miniaturize. Each successive hair growth cycle produces a shorter, finer hair until the

follicle eventually stops producing visible hair. The process is gradual — it happens over years, not weeks — which is why many women do not notice it until the thinning is significant.

Androgenic alopecia in PCOS can be treated, but it requires patience. Topical minoxidil (Rogaine) is the most widely studied treatment and works well for many women. Spironolactone can also help by blocking androgen receptors. Low-level laser therapy and platelet-rich plasma (PRP) injections have some evidence as well. Talk to your doctor about which options are appropriate for you.

One very important note: hair loss can also be caused by iron deficiency, thyroid disorders, and other conditions that overlap with PCOS. Before assuming all your hair loss is androgenic, have your doctor check ferritin (iron stores), TSH, and vitamin D. You may be dealing with multiple contributing factors.

Acanthosis Nigricans: The Skin Marker of Insulin Resistance

Acanthosis nigricans (AN) is a skin condition that appears as dark, velvety patches on the back of the neck, in the armpits, under the breasts, and in skin folds. It can look like dirt that will not wash off, though it is not dirt at all. It is a sign of significant insulin resistance.

AN is caused by high insulin levels stimulating the growth of skin cells. When insulin binds to receptors on keratinocytes (skin cells), it triggers proliferation, leading to

the thickened, darkened appearance. The presence of AN is a strong indicator that your insulin resistance is significant and needs to be addressed.

The good news is that AN is reversible. When insulin levels come down — through nutrition, movement, medications, or supplements — the skin patches often fade. Improvement in AN is one of the visible signs that your metabolic health is getting better.

Weight Gain and Body Composition: The PCOS Pattern

Weight gain in PCOS follows a characteristic pattern: it is centered in the abdomen, creating an apple-shaped body rather than a pear-shaped one. This is visceral fat — fat stored deep in the abdomen around the organs — and it is metabolically active, meaning it produces inflammatory compounds that worsen insulin resistance and hormone imbalance.

The reasons for this pattern are complex. Insulin resistance directs the body to store fat rather than burn it. High insulin also inhibits lipolysis — the breakdown of fat for energy. And the elevated androgens themselves promote abdominal fat storage. It is a triple threat.

Women with PCOS also tend to have higher rates of metabolic adaptation — their resting metabolic rate may be lower than expected for their body size. This means they burn fewer calories at rest compared to women without PCOS of the same age and weight. Add to this the fatigue

and exercise intolerance that many women with PCOS experience, and you have a condition that creates multiple biological barriers to weight management.

This is not your fault. These are real biological mechanisms, not character flaws. And they can be addressed — not by "eating less," but by taking the specific metabolic approaches covered in this book.

Sleep Apnea and Fatigue

Sleep apnea is significantly more common in women with PCOS, with studies finding rates of 8 to 15 percent — higher than the general female population even after controlling for body weight. The combination of insulin resistance, inflammation, and androgens affects the central nervous system's regulation of breathing during sleep. Additionally, abdominal obesity can physically compress the diaphragm and airways.

PCOS fatigue has multiple sources. There is the sleep disruption from apnea. There is the chronic inflammation, which produces fatigue through cytokine activity. There is the metabolic inefficiency — when your cells are insulin resistant, they do not take up glucose efficiently, and your muscles and brain may not be getting the energy they need. And there is the emotional burden of managing a chronic condition, which is exhausting in its own right.

If you are experiencing significant fatigue, talk to your doctor about sleep apnea screening and also check your vitamin D, iron, ferritin, and thyroid function — all of which are commonly low in PCOS and contribute to fatigue.

Mood Changes and the PCOS-Brain Connection

Mood changes in PCOS are real and have physiological roots. Androgens affect the brain's neurotransmitter systems, including serotonin and GABA. Insulin resistance affects glucose availability to the brain, which can cause brain fog and difficulty concentrating (sometimes called "PCOS brain"). And chronic inflammation has been linked to depression through the cytokine effects on brain function.

Chapter 9 covers the mental health dimensions of PCOS in depth. For now, know that the mood symptoms you may be experiencing — irritability, anxiety, depression, brain fog — are not just "in your head."

Body Image and PCOS

PCOS challenges your relationship with your body in ways that go beyond "I wish I looked different." It is a condition that makes your body feel like it is betraying you. The acne makes you feel like a teenager again. The hair growth feels unfeminine. The hair loss feels like you are aging too fast. The weight gain makes you feel like you have no control.

These feelings are normal. They are not shallow. Your appearance is tied to your identity and your sense of self, and when a medical condition hijacks your appearance, it is legitimate to grieve that.

But here is the counterpoint: as you learn to manage your PCOS — and you will learn — your relationship with your body can heal. Not because your symptoms will disappear

completely (they may not), but because you will understand what is happening and why. The sense of betrayal comes from not knowing. The sense of control comes from understanding.

Tracking Your Symptom Patterns

One of the most useful things you can do as you start managing PCOS is to track your symptoms consistently. Not obsessively, but consistently enough to see patterns. A simple tracker can include: - Date - Bleeding (if any) - Acne (location and severity on a 1-5 scale) - Hair growth (self-assessment) - Hair shedding - Weight - Energy level (1-5) - Mood (1-5) - Notable stress events - Food and exercise notes

When you bring this tracker to your doctor, it is worth a thousand words. It turns your subjective experience into objective data that can guide diagnosis and treatment.

When Symptoms Overlap: The Ferritin Connection

There is an important overlap between PCOS symptoms and iron deficiency that deserves its own section. Low ferritin (low iron stores) can cause fatigue, hair loss, brain fog, pale skin, and even irregular periods — all symptoms that look like PCOS or worsen PCOS symptoms. Many women with PCOS have heavy periods, which deplete iron. Others avoid red meat, have restrictive eating patterns, or have poor iron absorption due to the gut inflammation associated with insulin resistance.

Before attributing all your symptoms to PCOS, have your doctor check your ferritin level. Ideally, ferritin should be above 30 ng/mL for sufficient iron stores. Some experts recommend keeping it above 50 ng/mL for optimal hair growth. If your ferritin is low, iron supplementation (under your doctor's guidance, with retesting to avoid overload) can make a meaningful difference in your energy, hair health, and cognitive function.

Similarly, vitamin B12 deficiency, which is common in women taking metformin long-term, can cause fatigue, brain fog, and mood changes. And thyroid disorders, which are more common in women with PCOS, can cause many of the same symptoms. Always check the full picture before assuming a symptom is "just PCOS."

Toolkit Summary

- PCOS acne follows a jawline/chin pattern driven by androgen excess and inflammation
- Hirsutism affects 60–80 percent of women with PCOS and requires both medical and cosmetic management — talk to your doctor about spironolactone
- Androgenic alopecia causes crown thinning and part widening; minoxidil and spironolactone are first-line treatments
- Acanthosis nigricans is a visible marker of insulin resistance — it improves as insulin levels decrease
- PCOS weight gain is abdominal/visceral, driven by insulin resistance and androgen excess

- Sleep apnea is more common in PCOS — talk to your doctor if you experience significant fatigue
- Mood symptoms and brain fog are physiologically linked to PCOS
- Track your symptoms consistently to identify patterns and guide treatment

ch05

GETTING A PCOS DIAGNOSIS SHOULD BE STRAIGHTFORWARD. YOU GO TO A DOCTOR, YOU DESCRIBE YOUR SYMPTOMS, THEY RUN SOME TESTS, AND YOU GET AN ANSWER. SIMPLE, RIGHT?

For too many women, the reality looks very different. The doctor says your blood work looks "basically normal." They tell you to lose ten pounds. They hand you a prescription for birth control and send you on your way. You leave with the same questions you came in with, plus the feeling that you have been dismissed.

This chapter is here to change that. It walks you through exactly what tests you need, what the results mean, and how to find a doctor who will take you seriously. Consider it your diagnostic roadmap.

The Rotterdam Criteria: Your Diagnostic Foundation

PCOS is diagnosed using the Rotterdam criteria: you need at least two of the following three:

1. **Oligomenorrhea or anovulation** — fewer than eight periods per year, cycles consistently longer than 35 days, or confirmed absence of ovulation
2. **Hyperandrogenism** — either clinical signs (acne, hirsutism, alopecia) or elevated androgens on blood work

3. **Polycystic ovaries on ultrasound** — twelve or more follicles per ovary, or ovarian volume greater than 10 mL

Before applying these criteria, your doctor must rule out other conditions that can mimic PCOS. The standard differential diagnosis includes:

- **Thyroid disorders** — both hypothyroidism and hyperthyroidism can cause irregular periods. A TSH test rules this out.
- **Hyperprolactinemia** — elevated prolactin can cause anovulation and irregular periods. A prolactin blood test rules this out.
- **Non-classic congenital adrenal hyperplasia (NCAH)** — a genetic condition that causes elevated 17-hydroxyprogesterone (17-OHP) and mimics PCOS. A morning 17-OHP blood test rules this out. This is important because NCAH is more common in certain ethnic groups and requires different treatment.
- **Cushing's syndrome** — a rare condition of excess cortisol that can cause weight gain, irregular periods, and insulin resistance. If you have classic Cushing's features (moon face, buffalo hump, purple stretch marks), your doctor may test for this.

The Complete PCOS Lab Workup

Here is the blood work you should have if you are being evaluated for PCOS. You can request this as a "PCOS panel" — many labs now offer one. Your insurance may require specific codes, so talk to your doctor about what is covered.

Hormone Panel: - Total testosterone - Free testosterone (calculated from total and SHBG, or measured directly by equilibrium dialysis) - DHEA-S (dehydroepiandrosterone sulfate — an adrenal androgen) - Androstenedione (a precursor to testosterone) - Sex hormone-binding globulin (SHBG) — this is critical but often skipped - LH (luteinizing hormone) - FSH (follicle-stimulating hormone) - Prolactin - 17-hydroxyprogesterone (17-OHP) — to rule out NCAH

Metabolic Panel: - Fasting glucose - Fasting insulin — this is the most commonly skipped test and the most important one for PCOS - HbA1c (measures average blood sugar over the past 2-3 months) - Lipid panel (total cholesterol, LDL, HDL, triglycerides) - Liver function tests (ALT, AST) — PCOS increases NAFLD risk

Other: - Thyroid panel (TSH, free T4, free T3) - Vitamin D — deficiency is extremely common in PCOS - Ferritin — iron deficiency mimics many PCOS symptoms

Understanding Your Results

Total testosterone: Normal ranges vary by lab, but generally 15-70 ng/dL. In PCOS, it may be elevated or in the upper half of "normal."

Free testosterone: This is more useful than total testosterone. Even if total is normal, free testosterone can be elevated if SHBG is low. Many women with PCOS have normal total but elevated free testosterone.

SHBG: This is one of the most telling numbers. Normal SHBG is roughly 40–120 nmol/L. Low SHBG (below 40) is a hallmark of PCOS with insulin resistance. It tells you that your liver is being suppressed by high insulin.

LH and LH/FSH ratio: A ratio of 2:1 or higher suggests PCOS. However, this ratio can be normal in PCOS (particularly phenotype D), and it fluctuates through your cycle, so it is not diagnostic alone.

DHEA-S: This reflects adrenal androgen production. It is elevated in about 60 percent of women with PCOS. Very high DHEA-S (>600 mcg/dL) raises suspicion for adrenal tumors or NCAH.

Fasting insulin: This is the test most doctors skip, and it is the one that tells you whether insulin resistance is driving your PCOS. Fasting insulin over 8–10 mIU/L is considered elevated. Over 15 mIU/L indicates significant insulin resistance. Talk to your doctor about what your specific lab's reference range means.

HbA1c: Under 5.7 percent is normal. 5.7–6.4 percent indicates prediabetes. Over 6.5 percent indicates diabetes. Many women with PCOS have HbA1c in the 5.2–5.6 range but have fasting insulin over 10 — meaning they are insulin resistant but keeping their blood sugar in check by producing extra insulin. This is often missed.

The Ultrasound: What to Expect

A transvaginal ultrasound is the standard for assessing PCOS. It is slightly uncomfortable but not painful. The technician measures the size of each ovary and counts the number of follicles (the small fluid-filled structures that contain immature eggs).

The ultrasound criteria for PCOS are: - Twelve or more follicles (2-9 mm in diameter) in one or both ovaries - Ovarian volume greater than 10 mL

Important caveats: If you are on birth control pills, your ovaries may appear "normal" because the pill suppresses follicle development. The ultrasound should ideally be done when you are off hormonal contraception for at least three months. Also, polycystic ovaries are sometimes seen in women without PCOS (about 20-30 percent of women with regular cycles have polycystic-appearing ovaries), and some women with PCOS have "normal" ovaries on ultrasound — particularly those with phenotype B.

The ultrasound is one piece of the puzzle, not the whole picture.

The Caveats: When PCOS Does Not Look Like PCOS

PCOS can be tricky. Here are the most common situations where women get told they do not have PCOS when they actually do:

Case 1: You are thin, your blood work is "normal," but you have irregular periods. Some women with PCOS have normal androgens and normal insulin but still have irregular cycles. They fall into phenotype D. The diagnosis may hinge on the ultrasound or on more subtle markers like low SHBG.

Case 2: You have all the symptoms but normal ovaries on ultrasound. This is phenotype B. You have irregular periods, elevated androgens, and all the symptoms, but your ovaries look fine. Some doctors will dismiss you because "your ovaries are normal."

Case 3: You have regular periods but terrible symptoms. This is phenotype C — ovulatory PCOS. You ovulate regularly, so your cycles look normal, but you have high androgens and polycystic ovaries. Many women with this phenotype struggle with acne, hair loss, and hirsutism for years without anyone connecting it to PCOS.

Case 4: You are on birth control, and everything looks "fine." Birth control suppresses ovarian function and lowers androgen levels. Your blood work will look normal and your ovaries will look quiet on ultrasound. But that does not mean you do not have PCOS — it means the hormonal contraception is masking it. To get an accurate diagnosis, you need to be off hormonal birth control for at least three months.

How to Find a Doctor Who Actually Listens

Finding the right doctor may be the most important step in your PCOS journey. Here is what to look for:

An endocrinologist is often the best specialist for PCOS because PCOS is fundamentally an endocrine (hormonal and metabolic) condition. Reproductive endocrinologists are particularly good if you are interested in fertility. Many gynecologists are PCOS-literate, but some focus primarily on reproductive aspects and miss the metabolic piece.

Questions to ask before your appointment: - "Do you treat many patients with PCOS?" - "What tests do you order for a PCOS workup?" - "Do you check fasting insulin and SHBG routinely?" - "What treatment options do you offer beyond birth control pills?"

What to bring: - Your symptom tracker (dates, symptoms, patterns) - A list of which tests you want — reference this chapter - A family history (PCOS, diabetes, thyroid conditions, or infertility in close relatives) - A brief written summary of your history: when symptoms started, what you have tried, and what your goals are (better cycles, clearer skin, fertility, weight management, etc.)

If a doctor dismisses you, fire them. This is your health. You deserve a doctor who takes your symptoms seriously, who orders the right tests, and who partners with you in finding solutions. You can ask for a second opinion. You can switch providers. You can research PCOS-specialized clinics, which are becoming more common in major cities and through telehealth platforms.

Telehealth and PCOS: A Viable Option

If you live in an area without PCOS-literate providers, or if you have had difficulty finding a doctor who listens, telehealth PCOS clinics are becoming an increasingly good option. Several telemedicine platforms now specialize in PCOS and women's hormonal health, offering the full diagnostic workup, medication management (including metformin, spironolactone, and GLP-1 agonists for eligible patients), and ongoing monitoring — all from your home.

Telehealth is not appropriate for every situation (physical exams and ultrasounds still require in-person visits), but it can be an excellent option for the initial consultation, lab review, and ongoing medication management. Many of these services accept insurance, and the quality of care is often higher than what you would get from a generalist who sees one or two PCOS patients per year. If you have been struggling to find good care locally, telehealth is worth exploring.

The Emotional Toll of the Diagnostic Journey

It is also worth acknowledging that the diagnostic process itself can be emotionally exhausting. The months or years of not knowing. The gaslighting. The dismissive doctors. The friends and family who tell you "it is probably nothing." Getting diagnosed is a relief — but it also brings a complicated set of emotions: grief for the time you lost,

anger at the system that failed you, and fear about what the diagnosis means for your future. All of these are normal. Give yourself permission to feel them.

Advocating for Your Diagnosis When Tests Are "Normal"

One of the most common frustrations in the diagnostic journey is being told "your tests are normal" when you clearly have symptoms. If your lab results fall within the reference range but your symptoms tell a different story, remember that the reference range is not a diagnostic cutoff — it is a statistical distribution. A testosterone level at the 95th percentile of the "normal" range may still cause symptoms in a woman whose body is sensitive to androgens.

Push back gently but firmly. Ask your doctor: "My results may be in the normal range, but they may be high-normal for me. Can we look at my free testosterone, my SHBG, and my fasting insulin to get a more complete picture?" If your doctor is unwilling to look beyond the reference range, that is a strong signal that you need a second opinion. There are providers who understand that "normal" does not always mean "healthy" in the context of PCOS.

Toolkit Summary

- PCOS diagnosis requires at least 2 of 3 Rotterdam criteria after ruling out other conditions

- The essential lab workup includes testosterone (total and free), SHBG, LH, FSH, DHEA-S, fasting insulin, HbA1c, lipids, prolactin, 17-OHP, TSH, vitamin D, and ferritin
- Fasting insulin is the most commonly skipped but most important metabolic test for PCOS
- An ultrasound must be interpreted in context — women on birth control or with phenotype B may have "normal" ovaries
- PCOS can present atypically in women who are lean, on birth control, or have regular periods
- An endocrinologist is often the best specialist for comprehensive PCOS care
- Bring a symptom tracker and test list to your appointment — and do not hesitate to seek a second opinion if you are dismissed

ch06

IF YOU HAVE PCOS, YOU HAVE PROBABLY RECEIVED MORE UNSOLICITED NUTRITION ADVICE THAN YOU EVER WANTED. GO KETO. GO VEGAN. CUT ALL SUGAR. EAT SIX SMALL MEALS A DAY. INTERMITTENT FAST. NEVER EAT CARBS AFTER 4 P.M. THE SHEER VOLUME OF CONTRADICTORY ADVICE CAN MAKE YOU WANT TO GIVE UP AND JUST EAT WHATEVER IS IN THE FRIDGE.

Here is the truth: there is no single "PCOS diet" that works for everyone. But there are evidence-based nutrition principles that consistently help women with PCOS lower their insulin, reduce inflammation, improve their symptoms, and feel better — without requiring you to live on chicken and broccoli or give up all the foods you love.

This chapter lays out those principles and gives you a practical framework you can adapt to your life, your preferences, and your budget.

The Nutritional Goals of PCOS Management

Before we talk about specific foods, let's clarify what we are trying to accomplish nutritionally for PCOS. There are three primary goals:

Goal 1: Stabilize blood sugar. When your blood sugar spikes, your pancreas releases a surge of insulin to bring it down. For women with PCOS and insulin resistance, each blood sugar spike triggers an exaggerated insulin response.

Over time, these spikes worsen insulin resistance. The goal is to eat in a way that keeps blood sugar steady throughout the day — no dramatic spikes, no dramatic crashes.

Goal 2: Improve insulin sensitivity. Some foods and eating patterns can actually make your cells more responsive to insulin over time. These are the foods you want to build your diet around.

Goal 3: Reduce inflammation. PCOS is a state of chronic low-grade inflammation driven by insulin resistance and adipose tissue dysfunction. Anti-inflammatory foods can help calm this background inflammation, which in turn improves insulin sensitivity and reduces symptoms.

The Mediterranean Diet: The Most Evidence-Based Pattern for PCOS

When researchers look at which dietary pattern has the strongest evidence for PCOS, the Mediterranean diet consistently comes out on top. It is not a "diet" in the weight-loss sense — it is a way of eating that emphasizes whole foods, healthy fats, and plenty of vegetables.

What the evidence shows: Multiple studies have found that women with PCOS who follow a Mediterranean-style eating pattern show improvements in insulin sensitivity, reductions in inflammatory markers, decreases in androgen levels, and improvements in body composition. These benefits appear whether weight loss occurs or not — the metabolic improvements are independent of weight change.

The Mediterranean pattern looks like this: - Vegetables of all colors, especially leafy greens, cruciferous vegetables (broccoli, cauliflower, Brussels sprouts), and red/orange vegetables - Fruits, with an emphasis on berries and lower-glycemic options - Whole grains — quinoa, farro, barley, oats, whole wheat, brown rice - Legumes — lentils, chickpeas, beans of all types - Fish and seafood, especially fatty fish like salmon, sardines, and mackerel (rich in omega-3s) - Poultry, eggs, and dairy in moderate amounts - Red meat rarely (a few times per month) - Olive oil as the primary fat source - Nuts, seeds, and avocados - Herbs and spices for flavor instead of salt-heavy seasonings

The Mediterranean diet is not low-carb. It is moderate-carb — about 40 to 50 percent of calories from carbohydrates, but from high-quality, fiber-rich sources that digest slowly and do not spike blood sugar.

The Low-Carb vs. Moderate-Carb Debate

There is a persistent debate in the PCOS community about whether low-carb or ketogenic diets are necessary. Here is what the evidence says:

Very low-carb (keto) diets can work for some women with PCOS. They dramatically lower insulin levels because they minimize the glucose that triggers insulin release. Several small studies have found that keto diets improve PCOS symptoms, including regularizing periods and lowering androgens.

However, keto is not necessary for most women with PCOS. The improvement seen in keto studies is largely driven by the reduction in insulin, and you can achieve meaningful insulin reduction with a moderate-carb Mediterranean approach — without the extreme dietary restriction, the potential for nutrient deficiencies, or the difficulty of sustaining keto long-term.

The middle ground: Research suggests that for most women with PCOS, consuming about 40 percent of calories from carbohydrates — from high-fiber, low-glycemic sources — is sufficient to improve insulin sensitivity and symptoms. That works out to about 150 to 200 grams of carbohydrates per day for a 1,800-calorie diet. That is not low-carb by any standard, but it is a significant reduction from the standard American diet, which often exceeds 250 grams of carbohydrates per day, largely from refined sources.

The exception: Some women with PCOS have severe insulin resistance and find that they genuinely feel and function better at 30 percent carbohydrate or lower. This is worth trying if moderate-carb has not helped within three months. Talk to your doctor and consider working with a registered dietitian who is PCOS-informed to develop a plan that works for you.

The Glycemic Load Concept

Not all carbohydrates are created equal. A baked potato and a bowl of lentils have similar amounts of total carbohydrates, but they have very different effects on your blood sugar. This is where glycemic load comes in.

Glycemic load is a measure of how much a serving of food will raise your blood sugar. It accounts for both the glycemic index (how fast a food raises blood sugar) and the amount of carbohydrate in a serving. High-glycemic-load foods cause rapid spikes in blood sugar; low-glycemic-load foods cause gradual, modest rises.

High-glycemic-load foods to limit: - White bread, white rice, white pasta - Sugary beverages (soda, juice, sweetened coffee drinks) - Sugary snacks and desserts - Breakfast cereals (most, even the "healthy" ones) - Potatoes and French fries - White flour tortillas, pastries, bagels

Low-glycemic-load foods to emphasize: - Non-starchy vegetables (broccoli, spinach, peppers, zucchini, tomatoes, cauliflower) - Legumes (lentils, chickpeas, black beans) - Whole grains (quinoa, barley, steel-cut oats) - Berries and stone fruits - Nuts and seeds - Avocados and olives

A simple rule: if a carbohydrate source has fiber (at least 3 grams per serving) and/or protein, it will have a lower glycemic load than a refined, fiberless version of the same food.

The Anti-Inflammatory Eating Pattern

PCOS-related inflammation responds well to dietary changes. The key anti-inflammatory strategies are:

Increase omega-3 fatty acids. Omega-3s (found in fatty fish, flaxseeds, chia seeds, and walnuts) have direct anti-inflammatory effects. Aim for two servings of fatty fish per week, or consider a fish oil supplement (talk to your doctor about dosing).

Reduce omega-6 to omega-3 ratio. The standard Western diet is heavily weighted toward omega-6 fatty acids (found in vegetable oils, processed foods, and grain-fed meat), which promote inflammation. Reducing processed foods and incorporating more omega-3-rich foods helps rebalance this ratio.

Eat a rainbow of plant foods. Different colored plant foods contain different phytochemicals — quercetin in onions, lycopene in tomatoes, anthocyanins in berries — each with unique anti-inflammatory properties. Eating a wide variety ensures you get the full spectrum.

Limit ultra-processed foods. Foods made in a factory with long ingredient lists (including additives, preservatives, artificial flavors, and refined oils) tend to promote inflammation. Foods that come from nature tend to reduce it.

Meal Timing and Insulin

When you eat matters almost as much as what you eat. The key insulin-friendly meal timing principles are:

Eat regular meals. Skipping meals leads to blood sugar crashes followed by compensatory overeating, which causes spikes. Three moderate meals per day, or three meals and a small snack, tends to work best for blood sugar stability.

Avoid late-night eating. Eating large meals late at night can worsen next-morning fasting insulin levels. Try to finish your last meal at least three hours before bed.

Consider the "protein and produce" first strategy. When you eat a meal, eating protein and vegetables first, and carbohydrates last, can reduce the post-meal blood sugar spike by up to 30 percent due to the way food order affects glucose absorption.

What to Eat: A Practical Framework

If all of the above feels overwhelming, here is a simplified framework you can start with tomorrow:

Build every meal around: 1) a protein source (eggs, chicken, fish, tofu, beans, Greek yogurt), 2) non-starchy vegetables (fill half your plate), and 3) a high-quality fat source (olive oil, avocado, nuts, seeds). Add a small portion of low-glycemic carbohydrates (quinoa, sweet potato, lentils, berries) if you are hungry and active.

What to keep in your kitchen: - Extra virgin olive oil - Canned salmon or sardines - Frozen vegetables (broccoli, spinach, cauliflower) - Lentils and chickpeas (canned or dried) - Quinoa and oats - Eggs - Nuts and seeds (almonds, walnuts, chia seeds, flaxseeds) - Berries (fresh or frozen) - Plain Greek yogurt - Avocados

What to minimize in your kitchen: - Sugary beverages of any kind - White bread, white pasta, white rice - Sugary breakfast cereals and granola - Crackers, chips, and

processed snack foods - Sweetened yogurt and flavored oatmeals - Most salad dressings (make your own with olive oil and vinegar)

The Intuitive Eating Adaptation for PCOS

There is a tension between "PCOS nutrition" and the anti-diet movement, and it deserves acknowledgment. Many women with PCOS have a history of disordered eating — the frustration of unexplained weight gain leads to dieting, which leads to restriction, which leads to bingeing, which leads to guilt, which leads to more dieting.

The way forward is not more restriction. It is not punishing your body for having PCOS. The way forward is learning which foods make your body feel good and which ones do not, and making choices from that knowledge rather than from fear.

This is intuitive eating adapted for PCOS. You do not have to measure your food. You do not have to count grams of carbohydrates for the rest of your life. You need to learn the general principles — more vegetables, better carbohydrates, regular meals, adequate protein and fat — and then find the version of those principles that fits your actual life. Food is not punishment. Food is medicine, yes, but it is also pleasure, culture, connection, and joy. You get to have all of that.

Practical Meal-Planning Strategies

One of the biggest challenges with PCOS nutrition is not knowing what to eat — it is having the energy and time to prepare food when you are already tired from managing a chronic condition. Meal planning can help break that cycle. Start by identifying three dinners you can cook in under 30 minutes that follow the PCOS-friendly framework (protein, vegetables, healthy fat, optional low-GI carb). Keep the ingredients for those three meals on hand at all times.

Batch cooking is another practical strategy. On a Sunday afternoon, cook a large batch of quinoa or lentils, roast a tray of vegetables, and make a protein source (grilled chicken, hard-boiled eggs, or baked tofu). Having these components ready means you can assemble a PCOS-friendly meal in five minutes on a busy weekday. The barrier to eating well is almost never about knowing what to eat — it is about having the energy to prepare it when you are tired and hungry. Reduce that barrier by preparing ahead.

The Hydration and PCOS Connection

Hydration is an underappreciated factor in blood sugar management. Even mild dehydration can elevate cortisol and raise blood glucose levels. Women with PCOS should aim for at least eight to ten cups of fluid per day, primarily from water, herbal tea, and other unsweetened beverages. Staying well-hydrated supports kidney function (important if you take metformin or spironolactone), helps with fatigue, and may

reduce sugar cravings. If you struggle to drink enough water, try keeping a water bottle on your desk and setting a reminder to refill it twice during the day.

Toolkit Summary

- The three nutritional goals for PCOS are blood sugar stability, improved insulin sensitivity, and reduced inflammation
- The Mediterranean diet has the strongest evidence for improving PCOS symptoms and metabolic health
- Most women with PCOS do not need keto — 40 percent of calories from high-quality carbohydrates works for most
- Glycemic load matters more than glycemic index — choose fiber-rich carbohydrate sources
- Anti-inflammatory eating means more omega-3s, fewer processed foods, and a wide variety of plant foods
- Meal timing matters: regular meals, avoid late-night eating, and consider food order (protein and vegetables first)
- Build meals around protein, vegetables, and healthy fats with small portions of low-glycemic carbohydrates
- You do not need to restrict or count forever — learn the principles and find your own sustainable version

- Talk to your doctor or a registered dietitian before making significant dietary changes, especially if you are on medication

ch07

IF YOU HAVE BEEN TOLD TO "JUST EXERCISE MORE" FOR YOUR PCOS, YOU ALREADY KNOW HOW UNHELPFUL THAT ADVICE IS. IT IGNORES EVERYTHING THAT MAKES EXERCISE CHALLENGING WITH THIS CONDITION — THE FATIGUE, THE INFLAMMATION, THE EXERCISE INTOLERANCE, THE WAY YOUR BODY SEEMS TO HOLD ONTO WEIGHT NO MATTER WHAT YOU DO.

But here is what that advice gets right: exercise is one of the most effective tools for managing PCOS. The key is doing the right kind, at the right intensity, for the right reasons. This chapter breaks down exactly which types of movement improve insulin sensitivity, which types lower stress, and how to build an exercise routine that works with your hormones rather than against them.

Why Exercise Helps PCOS Specifically

Exercise works on PCOS through several interconnected mechanisms:

Insulin sensitivity improvement. Muscle contractions increase glucose uptake into muscle cells through an insulin-independent pathway. Put simply: when you move your muscles, they pull glucose out of your bloodstream whether or not insulin is working properly. This lowers your blood sugar and reduces the demand on your pancreas to produce

extra insulin. Over time, regular exercise improves your insulin sensitivity more broadly, meaning your cells become more responsive to insulin signals.

Stress reduction. Exercise lowers cortisol, the primary stress hormone. High cortisol worsens insulin resistance and directly stimulates the adrenal glands to produce more androgens. By lowering cortisol through appropriate exercise, you break this feedback loop.

Inflammation reduction. Regular exercise reduces levels of inflammatory markers, including C-reactive protein (CRP) and interleukin-6. Lower inflammation means less insulin resistance and less of the background inflammatory drive that worsens PCOS symptoms.

Body composition improvement. Exercise — especially resistance training — builds muscle. More muscle mass means a higher resting metabolic rate and more glucose disposal capacity. Exercise also promotes favorable changes in fat distribution, particularly reducing the visceral abdominal fat that drives metabolic dysfunction in PCOS.

Mood and energy improvement. The endorphin release and stress reduction from exercise directly counteract the higher rates of depression and fatigue seen in PCOS. This is not just a mental benefit — it is a physiological one that supports adherence to all your other PCOS management strategies.

What Type of Exercise Works Best: The Evidence

For decades, exercise advice for PCOS was generic: "do cardio." We now know that the best approach is more nuanced. The research points to a combination of resistance training and steady-state cardio as having the strongest evidence for PCOS.

Resistance training (strength training) improves insulin sensitivity by increasing muscle mass and by upregulating glucose transporter proteins (GLUT4) in muscle cells. It is also the most effective exercise modality for improving body composition — building lean mass while reducing fat mass. The evidence suggests that women with PCOS who do resistance training two to three times per week see significant improvements in insulin sensitivity, even without weight loss.

Steady-state cardio (brisk walking, jogging, cycling, swimming) improves cardiovascular fitness and contributes to insulin sensitivity improvements. The key phrase is "steady-state" — moderate, sustained effort that you can maintain for 30 to 45 minutes. This type of exercise is well-tolerated by most women with PCOS and does not typically over-elevate cortisol.

The combination appears to be more effective than either type alone. A 12-week program combining resistance training and steady-state cardio leads to greater improvements in insulin sensitivity, body composition, and symptom scores than either exercise type alone.

HIIT: Helpful or Harmful for PCOS?

High-Intensity Interval Training (HIIT) — short bursts of very intense exercise followed by brief recovery periods — has become popular in recent years. For PCOS, the evidence is mixed.

The upside of HIIT: It is time-efficient. A 20-minute HIIT session produces similar or better cardiovascular benefits than 40 minutes of steady-state cardio. It also improves insulin sensitivity, sometimes more rapidly than moderate-intensity exercise.

The downside of HIIT for PCOS: HIIT elevates cortisol. The intense, stressful nature of maximal-effort intervals triggers a significant cortisol response. For women with PCOS who already have elevated cortisol or who are sensitive to stress hormones, HIIT may worsen symptoms — particularly fatigue, mood instability, and sleep disruption.

The practical takeaway: HIIT is not off-limits for PCOS, but it is not a good choice as your primary exercise modality, especially if you are already under significant life stress. If you enjoy HIIT, limit it to one to two sessions per week, keep the "all-out" intervals to 20 to 30 seconds, and prioritize recovery afterward. Pay attention to how you feel — if HIIT leaves you wired and exhausted rather than energized, it may not be right for your body right now.

The Goldilocks Principle for PCOS Exercise

There is a concept I call the "Goldilocks principle" for PCOS exercise: not too little, not too much. Just the right amount for your body.

Too little exercise (zero or one session per week) does not provide enough stimulus to improve insulin sensitivity, reduce inflammation, or lower cortisol.

Too much exercise (daily high-intensity sessions, extended endurance training, training while under-recovered) can raise cortisol, increase inflammation, and worsen fatigue. This is particularly true for women with PCOS because their stress response systems are already dysregulated.

The right amount for most women with PCOS is three to five sessions per week, combining resistance and cardio, at moderate intensity. Sessions should last 30 to 50 minutes. This gives you enough stimulus to create metabolic improvements without overtaxing your stress systems.

Exercise Timing Relative to Meals

When you exercise in relation to meals can affect how much your blood sugar improves.

Exercise after meals (particularly after dinner) is especially effective for lowering post-meal blood sugar spikes. A 15-minute walk after a meal can reduce the glycemic response by 20 to 30 percent.

Exercise in a fasted state (first thing in the morning before eating) may improve insulin sensitivity more than exercise after eating, because the body is forced to use stored glucose and fat for fuel. However, some women with PCOS feel weak or lightheaded with fasted exercise, especially if their blood sugar regulation is already compromised. If you try fasted exercise and it does not feel good, eat something small beforehand — a protein-rich snack or a small piece of fruit — and exercise after.

The practical guideline: The best time to exercise is the time you can consistently do it. The metabolic benefits of exercise accumulate over weeks and months, not in a single session. Consistency matters far more than optimizing the timing.

Building Exercise Tolerance When You Are Starting from Zero

If you are not currently exercising, the most important thing is to start slowly enough that you do not get injured, overwhelmed, or discouraged. Here is a reasonable progression:

Week 1-2: Walk for 15 to 20 minutes per day, five days per week. That is it. Walking at a comfortable pace has measurable benefits for insulin sensitivity and stress reduction.

Week 3-4: Add one resistance training session per week. This can be bodyweight exercises (squats, lunges, push-ups against a wall, glute bridges) or machines at a gym. Two sets of 10 to 12 repetitions per exercise is plenty.

Week 5-6: Increase walking to 30 minutes, five days per week. Add a second resistance training session per week.

Week 7-8: You now have a routine: walking most days plus two resistance sessions per week. This is a solid foundation for PCOS management. You can continue at this level indefinitely or gradually increase intensity as your fitness improves.

PCOS and Cortisol: Why Stress Management Matters as Much as Exercise

One of the most overlooked aspects of PCOS management is the interaction between exercise and cortisol. Women with PCOS tend to have higher baseline cortisol levels and a more reactive stress response system. This means that exercise that is too intense, too frequent, or undertaken during high-stress life periods can actually worsen symptoms rather than improve them.

The signs that your exercise routine is raising your cortisol too much include: - You feel exhausted rather than energized after exercise - Your sleep quality worsens - You feel anxious or irritable - Your PCOS symptoms flare — more breakouts, more hair shedding, worse periods - You get sick more often - Your muscles are chronically sore

If you notice these signs, scale back. Reduce intensity, reduce frequency, or switch to a purely restorative practice (walking, gentle yoga, stretching) for a couple of weeks and see how you feel. The goal is to exercise in a way that makes you feel better, not worse.

Putting It All Together: A Sample Weekly Template

Here is a sample weekly exercise plan following the principles in this chapter:

Monday: Resistance training (full body, 30–40 minutes) **Tuesday:** Steady-state cardio (brisk walk or bike ride, 30 minutes) **Wednesday:** Resistance training (full body, 30–40 minutes) **Thursday:** Steady-state cardio (brisk walk or bike ride, 30 minutes) **Friday:** Active recovery (gentle yoga, stretching, or easy walk, 20 minutes) **Saturday:** Longer cardio (45–60 minute walk, hike, or swim) or HIIT (if you tolerate it, 15–20 minutes) **Sunday:** Rest or gentle movement (stretching, leisure walk)

This template is a guide, not a prescription. Listen to your body, adjust as needed, and celebrate consistency over perfection.

Exercise and Your Menstrual Cycle

For women with PCOS who do have some degree of cycle regularity, aligning exercise intensity with your cycle phase can be helpful. During the follicular phase (the week or two

leading up to ovulation), estrogen is rising and your body tends to handle higher-intensity exercise well. This is a good time for heavier resistance training or slightly more challenging workouts. During the luteal phase (after ovulation), progesterone rises and your body temperature is higher, which can make intense exercise feel harder. This phase is better suited to moderate steady-state cardio, lighter resistance work, and more restorative movement.

For women with PCOS who are not ovulating regularly, this cycling approach does not apply in the same way — but the principle of listening to your body is even more important. Your energy levels may fluctuate unpredictably. Let them guide you. Some weeks you will feel strong and capable of pushing harder. Other weeks, a gentle walk and some stretching is the right choice. The goal is not to hit a specific intensity target every session. The goal is to move consistently and in a way that makes you feel better, not worse.

Overcoming Common Exercise Barriers

Many women with PCOS face real barriers to consistent exercise: fatigue is the most common, followed by time constraints, lack of energy after work, and the discouragement of not seeing visible results quickly. If fatigue is your primary barrier, focus on the minimum effective dose: ten minutes of movement on the days you have the least energy. Ten minutes of walking, stretching, or

gentle bodyweight exercises still stimulates glucose uptake and lowers cortisol. It counts. Do not let perfect be the enemy of done.

If lack of visible results is discouraging you, shift your focus from appearance outcomes to process outcomes. Track how you feel after exercise, not what the scale says. Track how your energy levels change over the course of a week. Track whether your sleep improves on days you exercise. These are meaningful, measurable benefits that happen long before your body composition changes. The metabolic improvements happen first — the visible changes follow.

Toolkit Summary

- Exercise improves PCOS through insulin sensitivity increase, cortisol reduction, inflammation reduction, and body composition improvement
- The best evidence supports combining resistance training (2–3 times per week) with steady-state cardio
- HIIT has mixed effects — limit to 1–2 sessions per week and pay attention to how you feel afterward
- The Goldilocks principle: 3–5 sessions per week, 30–50 minutes, moderate intensity
- Exercise after meals reduces post-meal blood sugar spikes — a 15-minute walk after eating helps
- Build tolerance gradually — start with daily walking, then add resistance training

- If exercise leaves you exhausted, scale back — more is not always better for PCOS
- Stress management is essential — you may need different exercise approaches during high-stress periods

ch08

NAVIGATING THE WORLD OF PCOS TREATMENTS CAN FEEL OVERWHELMING. BETWEEN PRESCRIPTION MEDICATIONS WITH INTIMIDATING SIDE EFFECT PROFILES AND SUPPLEMENTS SOLD WITH DRAMATIC PROMISES BUT LITTLE EVIDENCE, IT IS HARD TO KNOW WHAT IS WORTH TAKING AND WHAT IS NOT.

This chapter cuts through the noise. It covers the most common medications and supplements used for PCOS, what the evidence actually says, who they tend to help most, and the important cautions to discuss with your doctor.

A critical note before we begin: every medication and supplement discussed in this chapter works differently for different people. What helps your friend with PCOS may not help you. Finding the right combination often requires patience and experimentation under medical supervision. Never start or stop medications or supplements without talking to your doctor first.

Metformin: The Insulin-Sensitizing Workhorse

Metformin is the most studied insulin-sensitizing medication for PCOS. It works primarily by reducing glucose production in the liver and improving glucose uptake in muscle tissue, which lowers circulating insulin levels.

How it helps PCOS: By lowering insulin, metformin reduces ovarian androgen production, improves SHBG levels, and can help restore ovulation in some women. Studies find

that metformin increases ovulation rates by about 30 to 50 percent in women with PCOS and improves pregnancy rates when used alongside ovulation induction medications. It also modestly improves weight loss, particularly when combined with nutrition and lifestyle changes.

Who benefits most: Women with PCOS who have insulin resistance — which, as we have covered, is about 70 percent of women with PCOS. Metformin is less effective for women with PCOS who do not have insulin resistance. Women with higher BMI and those with phenotype A or B (the metabolic phenotypes) tend to have the best response.

Side effects and how to manage them: The most common side effects are gastrointestinal — nausea, diarrhea, stomach upset, and loss of appetite. These affect approximately 20 to 30 percent of women starting metformin. The extended-release version (metformin ER) is significantly better tolerated than the immediate-release version. Starting with a low dose (500 mg once daily) and gradually increasing over 4 to 6 weeks reduces GI side effects. Taking metformin with food also helps. If GI side effects are severe, talk to your doctor about the extended-release formulation or about discontinuing — there are other options.

Metformin and vitamin B12: Long-term metformin use (more than 2 years) can deplete vitamin B12. If you take metformin for more than a year, your doctor should check your B12 levels annually. Supplementing with B12 is reasonable.

Inositol: The Supplement with Real Evidence

Inositol is a naturally occurring sugar alcohol that acts as a second messenger in insulin signaling pathways. It is not a vitamin or mineral, but it is often grouped with B vitamins. There are two forms relevant to PCOS.

Myo-inositol (MI) is the most abundant form in the body. It is found in fruits, beans, grains, and nuts, but PCOS treatment requires supplemental doses.

D-chiro-inositol (DCI) is a derivative that acts through a different pathway. In women with PCOS, the conversion of MI to DCI may be impaired, leading to a DCI deficiency in certain tissues.

The 40:1 ratio: Research suggests that the optimal MI-to-DCI ratio for PCOS is approximately 40:1 — closely matching the physiological ratio found in the body. Many PCOS supplements now offer this ratio in a single product. High doses of DCI alone (without MI) may actually worsen egg quality and ovarian response, so combination products are preferred.

What the evidence shows: Multiple meta-analyses have found that inositol supplementation (MI at 2–4 grams per day plus DCI at 50–100 mg per day, or combined in the 40:1 ratio) improves insulin sensitivity, reduces triglycerides, lowers blood pressure, reduces androgen levels, and improves ovulation rates — in some studies, comparable to metformin. The evidence is strongest for ovulation and metabolic outcomes. Inositol is generally very well-tolerated with minimal side effects (occasional mild GI upset).

Dosing: The typical therapeutic dose is 2 to 4 grams of myo-inositol plus 50 to 100 mg of D-chiro-inositol per day, taken in divided doses (morning and evening). Higher doses are not necessarily better and may cause GI discomfort.

Birth Control Pills: The Pros and the Caveats

Combined oral contraceptives (the birth control pill) are the most commonly prescribed treatment for PCOS. They work by suppressing ovarian function, which stops the production of ovarian androgens and creates a regular, predictable withdrawal bleed.

The benefits: - Regular, predictable bleeding (reduces endometrial cancer risk from prolonged anovulation) - Dramatic improvement in acne (estrogen increases SHBG, which lowers free testosterone) - Reduction in hirsutism (slows new hair growth, though existing hair needs cosmetic management) - Reduction in hair loss - Reliable contraception

The downsides: - Birth control masks the underlying metabolic problem — insulin resistance continues progressing even if symptoms are controlled - It can worsen insulin sensitivity in some women (especially pills with higher progestin doses or more androgenic progestins) - It is not a treatment — it is symptom management - When you stop the pill, symptoms may return more severely than before ("post-pill PCOS flare") - It increases blood clot risk, particularly in women with PCOS who already have higher cardiovascular risk - It can affect mood — some women experience depression or mood changes on hormonal contraception

The bottom line: Birth control is a reasonable option for short-to-medium-term symptom management, especially in women who are not trying to conceive and do not have significant metabolic concerns. But it should not be the only treatment. If you are on birth control for PCOS, you should also be addressing insulin resistance and metabolic health through the nutrition, movement, and stress management approaches discussed in this book. Talk to your doctor about whether birth control is right for you and which formulation is best.

Spirolactone: The Anti-Androgen

Spirolactone is a medication that blocks androgen receptors and reduces androgen production. It is not FDA-approved specifically for PCOS, but it is widely used off-label for the androgenic symptoms — acne, hirsutism, and androgenic alopecia.

How it works: Spirolactone binds to androgen receptors and prevents testosterone from binding, and it also inhibits enzymes involved in testosterone production. The result is less androgen activity at the skin and hair follicle level.

What it helps: Cystic acne (most women see significant improvement within 2 to 3 months), hirsutism (new hair grows more slowly, finer, and lighter — takes 4 to 6 months to see results), and androgenic alopecia (some women see reduced shedding and improved hair density).

The pregnancy warning: Spironolactone is teratogenic — it can cause birth defects in a male fetus. Anyone taking spironolactone who could become pregnant must use effective contraception. This is typically why doctors prescribe it alongside birth control.

Side effects: The most common is frequent urination (spironolactone is a diuretic). Others include breast tenderness, fatigue, and dizziness. It can also cause elevated potassium, so your doctor should check your potassium levels after starting the medication.

Dosing and timeline: Typical doses are 50 to 200 mg per day. Start at a low dose and increase gradually. Results take time — 2 to 3 months for acne, 4 to 6 months for hair growth, and 6 to 12 months for hair loss improvement. Do not expect overnight results.

GLP-1 Agonists: The Emerging PCOS Treatment

GLP-1 receptor agonists (semaglutide, tirzepatide) are emerging as effective treatments for PCOS, particularly for women with significant insulin resistance, metabolic syndrome, or obesity.

The evidence in PCOS: Studies have found that GLP-1 agonists improve insulin sensitivity, promote significant weight loss, reduce androgen levels, and can restore ovulation. Some women with PCOS who have not responded to metformin find GLP-1 agonists much more effective.

Who is a candidate: These are prescription medications with specific prescribing criteria. Insurance coverage varies and can be a barrier. Talk to your doctor about whether you meet the criteria.

Side effects: Primarily gastrointestinal — nausea, vomiting, diarrhea, constipation. These are most pronounced when starting and when increasing the dose. Taking them with food and starting at the lowest dose helps.

Beyond Metformin: Other Anti-Diabetic Medications

Other medications used off-label for PCOS include:

Pioglitazone is a thiazolidinedione that improves insulin sensitivity by activating PPAR-gamma receptors. It is effective for PCOS but has significant side effects (weight gain, fluid retention, potential bone loss) and safety concerns that make it a third-line option at best.

Acarbose blocks carbohydrate absorption in the intestine, reducing post-meal blood sugar spikes. It is less commonly used due to GI side effects.

Supplement Evidence Review

Beyond inositol, these are the supplements with the strongest evidence for PCOS:

Vitamin D: Deficiency rates in PCOS are extremely high — 60 to 80 percent of women have insufficient levels. Vitamin D deficiency is linked to worse insulin resistance,

higher androgen levels, and worse mood. Supplementing to reach sufficient levels (30–50 ng/mL) improves metabolic and hormonal outcomes. A typical dose is 1,000 to 4,000 IU per day, but your doctor should test your levels and guide dosing.

Omega-3 fatty acids: Fish oil reduces triglycerides, lowers inflammation, and modestly improves insulin sensitivity in PCOS. Two grams per day of combined EPA and DHA is a typical dose.

Magnesium: Magnesium deficiency is common in PCOS (partly because insulin resistance increases urinary magnesium excretion). Supplementing improves insulin sensitivity and may improve sleep and mood. Magnesium glycinate is well-absorbed and gentle on the stomach.

Zinc: Zinc plays a role in insulin signaling, inflammation, and androgen metabolism. Studies in PCOS have found that zinc supplementation reduces insulin resistance, lowers triglycerides, and may reduce hirsutism. Typical dose is 20 to 30 mg per day.

Chromium: Chromium picolinate (200–1,000 mcg per day) may improve glucose tolerance and reduce insulin resistance. The evidence is modest but consistent.

N-acetylcysteine (NAC): NAC is an antioxidant that improves insulin sensitivity and has been studied for ovulation induction in PCOS. At 1,200 to 1,800 mg per day, some studies find ovulation rates similar to metformin. NAC is generally well-tolerated.

Berberine: Berberine is a plant alkaloid that improves insulin sensitivity through AMPK activation — the same pathway as metformin. A 2016 meta-analysis found that

berberine improved insulin sensitivity, reduced androgens, and improved waist-to-hip ratio in PCOS. Dose is typically 500 mg two to three times per day with meals.

Spearmint tea: There is small but promising research that spearmint tea (two cups per day) may reduce free testosterone levels in women with PCOS. The mechanism is not fully understood. It is a mild intervention but very safe to try.

Supplement Safety and Quality

A few important notes on supplements: They are not regulated by the FDA. Quality varies dramatically between brands. Look for third-party tested products (USP, NSF, or ConsumerLab seals). Buy from reputable companies. And remember: supplements are adjuncts, not substitutes. They work best alongside the nutrition, movement, and medical strategies discussed throughout this book.

The Cost-Benefit Framework for PCOS Treatments

Not every treatment is worth pursuing for every woman. A useful framework for deciding whether a medication or supplement is right for you involves weighing three factors: evidence strength, side effect burden, and cost. High-evidence, low-side-effect, low-cost interventions (like inositol, vitamin D, or consistent eating patterns) are almost always

worth trying first. Lower-evidence, higher-cost, higher-side-effect interventions should be reserved for situations where first-line approaches have not been sufficient.

This is where your symptom tracker and annual lab review become essential tools. If your fasting insulin is declining and your cycles are becoming more regular on a basic regimen of nutrition changes, inositol, and metformin, there may be no need to pursue more advanced treatments. But if your insulin remains high and your symptoms are not improving after six months of consistent effort, that is the time to have a more aggressive conversation with your doctor about GLP-1 agonists, higher medication doses, or additional supplements. The right treatment is the minimum effective dose — the simplest regimen that achieves your goals with the fewest side effects and the lowest burden.

Toolkit Summary

- Metformin improves insulin sensitivity, ovulation, and modest weight loss in PCOS — start low and go slow to manage GI side effects
- Inositol (myo-inositol plus D-chiro-inositol in a 40:1 ratio) has strong evidence for improving insulin sensitivity and ovulation
- Birth control pills manage symptoms but do not treat the underlying metabolic condition — they should not be your only treatment
- Spironolactone works well for acne and hirsutism but requires birth control due to pregnancy risks

- GLP-1 agonists are an emerging treatment with strong metabolic benefits — talk to your doctor if you have insulin resistance or weight concerns
- Vitamin D, omega-3s, magnesium, zinc, chromium, NAC, and berberine all have supportive evidence for PCOS
- Never start or stop any medication or supplement without talking to your doctor first
- Choose third-party tested supplement brands for quality and purity

ch09

IF THERE IS ONE ASPECT OF PCOS THAT IS MOST CONSISTENTLY OVERLOOKED BY THE MEDICAL SYSTEM, IT IS THE MENTAL HEALTH DIMENSION. DOCTORS WILL PRESCRIBE BIRTH CONTROL FOR YOUR ACNE, METFORMIN FOR YOUR INSULIN, AND SPIRONOLACTONE FOR YOUR HAIR GROWTH — BUT WHO PRESCRIBES THE SUPPORT FOR THE DEPRESSION, THE ANXIETY, THE BODY GRIEF, AND THE EXHAUSTION OF MANAGING A CHRONIC CONDITION THAT NOBODY SEEMS TO UNDERSTAND?

This chapter is for that part of the PCOS experience. The part that no blood test measures and no ultrasound shows. The part that is just as real as the hormonal imbalances and just as deserving of attention.

The PCOS–Mental Health Connection: What the Numbers Say

The statistics are striking. Women with PCOS have significantly higher rates of depression compared to women without PCOS. Meta-analyses estimate that approximately 40 percent of women with PCOS experience clinically significant depressive symptoms, compared to roughly 20 percent in the general female population. Anxiety rates are similarly elevated.

But the mental health burden extends beyond depression and anxiety. Women with PCOS have higher rates of: - Eating disorders (binge eating disorder in particular, which is linked to insulin dysregulation) - Body dysmorphic disorder (related to the visible symptoms — acne, hirsutism, hair loss, weight changes) - Sexual dysfunction (lower libido, reduced satisfaction) - Reduced quality of life (across physical, emotional, and social domains) - Suicidal ideation (risk is elevated, particularly in women with significant hirsutism or infertility)

These are not small increases. These are substantial, clinically meaningful differences from the general population. And they suggest that the mental health impact of PCOS is driven by something deeper than just "feeling bad about having a health condition."

The Hormonal Contribution to Mental Health

The connection between PCOS and mental health is not purely psychosocial, though that dimension is real too. There are biological mechanisms at work.

Androgens and the brain. Testosterone and other androgens affect brain function in complex ways. Androgen receptors are present throughout the brain, including in regions that regulate mood, anxiety, and reward. Elevated androgens in PCOS may directly affect neurotransmitter systems, particularly serotonin and GABA. Low serotonin is associated with depression; low GABA is associated with anxiety. The elevated androgens of PCOS may dysregulate both systems.

Insulin and brain function. The brain is a glucose-hungry organ — it consumes about 20 percent of the body's glucose. When insulin resistance is present, the brain's ability to take up and use glucose efficiently may be compromised. This can cause brain fog, difficulty concentrating, and memory issues (sometimes reported by women with PCOS as "PCOS brain"). Low brain glucose availability also affects mood regulation.

Inflammation and depression. The chronic inflammation that accompanies PCOS has been directly linked to depression through the "cytokine hypothesis" of depression. Inflammatory cytokines (including interleukin-6 and TNF-alpha) can cross the blood-brain barrier and affect neurotransmitter metabolism, particularly by diverting tryptophan away from serotonin production. This is a plausible biological pathway linking PCOS-related inflammation to depression.

Cortisol dysregulation. Many women with PCOS have altered cortisol rhythms — higher baseline cortisol, a blunted morning cortisol peak, or a slower decline throughout the day. Cortisol dysregulation is strongly associated with depression, anxiety, fatigue, and sleep disruption.

The Psychosocial Burden: Living with an Invisible Condition

The biological factors are real, but they do not tell the whole story. There is also the lived experience of PCOS — the burden of managing a chronic condition that is poorly understood, visibly impacts your appearance, and affects some of the most intimate aspects of your life.

The frustration with symptoms. Acne that will not clear despite expensive skincare. Hair that grows where you do not want it and falls out where you do. Weight that seems to accumulate despite your best efforts. The feeling of being at war with your own body.

The body image struggle. PCOS attacks the physical markers of femininity in a culture that places enormous value on female appearance. The acne, the hair growth, the hair loss, the abdominal weight — each symptom chips away at how you see yourself and how you believe others see you. This is not vanity. This is a real and painful assault on your identity.

The fertility grief. For women with PCOS who want children, the fertility component can be devastating. Each negative pregnancy test, each failed ovulation induction cycle, each month of waiting — the grief accumulates. And it is often invisible grief, because infertility is not something most people discuss openly.

The weight of invisibility. PCOS is an invisible condition. You look "fine" to the outside world. No one sees the irregular periods, the internal hormonal chaos, the three

years of trying to get diagnosed. This invisibility means you are often suffering without the support network that visible illness sometimes brings.

Therapy Approaches That Help

If you are struggling with the mental health dimensions of PCOS, professional support can be transformative. Here are the therapy approaches with the strongest evidence:

Cognitive Behavioral Therapy (CBT) focuses on identifying and changing unhelpful thought patterns and behaviors. For PCOS, CBT can help address body image distortions, reduce catastrophizing about symptoms, and develop behavioral strategies for managing stress and mood.

Acceptance and Commitment Therapy (ACT) focuses on accepting difficult thoughts and feelings rather than fighting them, and committing to values-based action. For PCOS, ACT can help you live a meaningful life alongside your symptoms, rather than waiting until your symptoms are "fixed" to start living.

PCOS-specific counseling. Some therapists now specialize in PCOS or reproductive health. If you can find one, a specialist who understands the hormonal dimension and the fertility grief can offer targeted support that a general therapist might not.

Support groups. Online and in-person support groups for PCOS provide validation, practical advice, and a sense of community. Organizations like PCOS Challenge and the PCOS Awareness Association offer support groups and resources.

Medical Treatment That Helps Mental Health

One of the most hopeful findings in PCOS research is that treating the underlying PCOS often improves mental health. When you address the insulin resistance, the androgen excess, and the inflammation, mood and anxiety often improve alongside the physical symptoms.

Several studies have found that: - Metformin therapy is associated with reductions in depressive symptoms - Inositol supplementation improves anxiety and depression scores in some women - Weight loss (through any approach) is associated with improved quality of life and mood - Regular ovulation and predictable periods reduce the anxiety of reproductive uncertainty - Improvement in visible symptoms (acne, hirsutism) improves body image and self-esteem

This does not mean you should skip therapy and just take supplements. But it does mean that your physical treatment and your mental health treatment are not separate tracks — they are connected. When one gets better, the other tends to follow.

What Helps at the Individual Level

Beyond professional treatment, there are strategies you can start using today to support your mental health with PCOS:

Separate the condition from your identity. PCOS is something you manage. It is not who you are. Practice saying (out loud if it helps): "I have PCOS, but PCOS does not have me." This reframe matters more than you think.

Name the grief. It is okay to be sad that you have PCOS. It is okay to be angry about the time you lost before getting diagnosed. It is okay to grieve the body you thought you would have. Let yourself feel these things without judgment.

Find your people. Whether it is a partner, a friend, a therapist, or an online community, find someone who gets it. PCOS is isolating, but it does not have to be endured alone.

Practice self-compassion. When you look in the mirror and see symptoms you do not want, try saying something kind to yourself. "This is hard. I am doing my best. My body is not my enemy, even when it feels that way." The research on self-compassion shows it reduces depression and anxiety and improves health behaviors.

Set boundaries with unsolicited advice. When someone tells you to "just try keto" or "just relax and you will get pregnant," you are allowed to say: "I appreciate that you care, but my PCOS is something I am managing with my doctor." You do not owe anyone an explanation.

A Word on Suicidal Ideation

If you are having thoughts of harming yourself, please know that you are not alone and that help is available. The National Suicide Prevention Lifeline (988 in the United States) is available 24/7. You can also text HOME to 741741. These resources are free and confidential.

PCOS can make life feel unbearable at times — there is no shame in that. There is only the need to reach out for support, and you deserve that support.

Building Emotional Resilience Over Time

Living with PCOS is a marathon, not a sprint. The mental health challenges will not be resolved in a week or a month. But over time, as you build a management plan that works and develop the skills to navigate the hard days, emotional resilience grows. You learn that bad days end. You learn that a symptom flare does not mean you are back at square one. You learn that you have survived every hard day so far, and you will survive the next one too.

Resilience is not about being positive all the time. It is about being honest about the difficulty while also trusting your capacity to handle it. It is about knowing when to push through and when to rest. It is about having a toolkit of strategies — the right therapist, the right community, the right self-care practices — that you can draw on when the weight of PCOS feels heavy.

Some days, the best you can do is show up for your one non-negotiable health habit (maybe it is taking your inositol, or going for your walk, or eating a decent breakfast) and let the rest slide. That is not failure. That is wise resource management. You are living with a chronic condition, and you are doing it with grace.

When to Seek Professional Mental Health Support

If you are experiencing persistent depression (low mood, loss of interest, changes in appetite or sleep for more than two weeks), significant anxiety (racing thoughts, avoidance behaviors, panic attacks), disordered eating patterns (restriction, binge eating, purging, or obsessive thoughts about food), or body dysmorphic symptoms (spending hours focused on your appearance, avoiding social situations because of how you look), it is important to seek professional help. These are not character flaws — they are medical conditions that respond to treatment.

A good therapist who understands PCOS or chronic illness can make a profound difference. If therapy feels inaccessible due to cost or location, consider online therapy platforms that accept insurance or offer sliding-scale fees. Many have therapists who specialize in reproductive health. Your doctor can also help by screening for depression and anxiety at your annual visit and, if needed, prescribing medications that are compatible with PCOS management. You do not have to carry this alone.

Toolkit Summary

- Depression and anxiety rates in PCOS are approximately double the general population
- Biological drivers include androgen effects on brain chemistry, insulin-related brain glucose dysregulation, inflammation-driven neurotransmitter disruption, and cortisol dysregulation
- Psychosocial burdens include body image struggles, fertility grief, and the invisibility of PCOS
- CBT and ACT are effective therapy approaches for PCOS-related mental health struggles
- Treating the underlying PCOS — insulin resistance, androgens, inflammation — often improves mental health
- Self-compassion, naming your grief, finding community, and setting boundaries all support mental health
- The National Suicide Prevention Lifeline (988) is available 24/7 if you need immediate support
- If you are experiencing depression or anxiety, talk to your doctor about a comprehensive approach that addresses both your PCOS and your mental health

ch10

IF YOU HAVE PCOS AND ARE THINKING ABOUT PREGNANCY, YOU PROBABLY HAVE A MIX OF HOPE AND ANXIETY. HOPE BECAUSE PCOS-RELATED INFERTILITY IS THE MOST TREATABLE FORM OF INFERTILITY — AND MOST WOMEN WITH PCOS WHO WANT TO CONCEIVE EVENTUALLY DO. ANXIETY BECAUSE YOU HAVE HEARD THAT PCOS MAKES PREGNANCY HARDER, THAT MISCARRIAGE IS MORE COMMON, THAT THE RISKS ARE ELEVATED.

Both the hope and the anxiety are grounded in real data. This chapter gives you the full picture — what the research says about fertility treatment, miscarriage risk, pregnancy complications, and postpartum considerations — so you can go into pregnancy planning informed and prepared.

Fertility Treatment for PCOS: Where to Start

For women with PCOS who are not ovulating regularly, fertility treatment begins with ovulation induction. The standard approach involves oral medications that stimulate follicle development.

Letrozole (Femara) is first-line. Multiple studies, including the landmark AMIGOS trial, have established letrozole as the superior medication for ovulation induction in PCOS. In that study, women taking letrozole had a 27.5 percent live birth rate, compared to 19.1 percent with

clomiphene — a significant difference. Letrozole also carries a lower risk of multiple pregnancy and does not thin the uterine lining the way clomiphene can.

Clomiphene (Clomid) remains a reasonable alternative, particularly if cost or insurance coverage is a concern. It works in a different way — blocking estrogen receptors rather than lowering estrogen — but the overall principle is the same: it triggers FSH release, which stimulates follicle growth.

Metformin as an adjunct. Adding metformin to ovulation induction may improve response rates, particularly in women who are insulin resistant. Some studies show that metformin alone can restore ovulation in about 30 percent of women with PCOS. The addition of metformin to letrozole or clomiphene may improve pregnancy rates, though the evidence is mixed. Talk to your doctor about whether adding metformin is right for you.

Inositol for egg quality. Inositol (myo-inositol plus D-chiro-inositol in the 40:1 ratio) has been studied for improving egg quality in women with PCOS undergoing fertility treatment. Some research suggests that inositol supplementation before and during treatment improves oocyte maturity, embryo quality, and pregnancy rates. If you are planning fertility treatment, starting inositol two to three months beforehand is reasonable.

When ovulation induction fails. If you have not conceived after three to six ovulatory cycles on oral medications, the next step is typically injectable

gonadotropins (FSH injections) with close monitoring, or proceeding to IVF. Your reproductive endocrinologist will guide this decision based on your specific situation.

Miscarriage Risk in PCOS

This is a heavy topic, but it deserves direct discussion. Women with PCOS have a higher risk of miscarriage. Studies estimate the risk at 30 to 50 percent in women with PCOS, compared to 15 to 20 percent in the general population.

The reasons are likely multifactorial: - **Insulin resistance** is associated with early pregnancy loss, possibly because it affects endometrial receptivity — the ability of the uterine lining to support an implanted embryo. - **Inflammation** may also impair implantation and early pregnancy development. - **Elevated LH levels** may contribute to higher rates of miscarriage by affecting egg quality at the time of ovulation. - **Progesterone insufficiency** is thought to occur in some women with PCOS due to the underlying hormonal dysregulation.

What reduces miscarriage risk: The most important intervention is improving metabolic health before conception. Multiple studies have found that women with PCOS who take metformin before and during early pregnancy have lower miscarriage rates compared to those who do not. Inositol supplementation before pregnancy also shows promise for reducing miscarriage risk. Optimizing blood sugar control, achieving a healthy weight (if weight is a factor), and addressing inflammation are all worth discussing with your doctor before trying to conceive.

Progesterone support after ovulation. Some reproductive endocrinologists prescribe progesterone supplementation during the luteal phase and early pregnancy for women with PCOS, reasoning that progesterone insufficiency may contribute to miscarriage. The evidence is not definitive, but it is a reasonable conversation to have with your doctor.

Pregnancy Complications with PCOS

Women with PCOS have higher rates of several pregnancy complications. Knowing these risks ahead of time allows you to work with your doctor on prevention and monitoring.

Gestational diabetes (GDM): This is the most significant pregnancy complication for women with PCOS. The risk of GDM is approximately two to four times higher in women with PCOS compared to women without PCOS. This is directly related to the underlying insulin resistance. Screening with a glucose tolerance test is standard in all pregnancies, but women with PCOS may benefit from earlier or more frequent screening. Talk to your doctor about when your screening should occur.

Preeclampsia: The risk of preeclampsia (a dangerous condition of high blood pressure during pregnancy) is increased in PCOS. The mechanism likely involves the combination of insulin resistance, inflammation, and vascular dysfunction. Management includes regular blood pressure monitoring and, if needed, low-dose aspirin starting in the second trimester for prevention.

Preterm birth: Some studies find a modestly increased risk of preterm birth in women with PCOS. The risk appears to be highest in women with significant insulin resistance or metabolic syndrome. Close monitoring of cervical length and fetal growth is standard.

Cesarean delivery: Women with PCOS have higher rates of C-section. This is partly due to higher rates of pregnancy complications that necessitate surgical delivery, and partly due to higher rates of large-for-gestational-age babies (driven by insulin resistance and GDM).

Managing PCOS During Pregnancy

Blood sugar monitoring: If you have PCOS, especially with known insulin resistance, your doctor may recommend earlier or more frequent glucose screening. Some women with PCOS who use continuous glucose monitoring during pregnancy find it helpful for keeping blood sugar in the optimal range.

Metformin during pregnancy: Metformin is considered safe to continue during pregnancy. There is good evidence that continuing metformin through the first trimester reduces the risk of pregnancy loss in women with PCOS. Some women continue metformin throughout pregnancy to reduce GDM risk and manage weight gain. Talk to your doctor about whether you should continue metformin when you conceive.

Inositol during pregnancy: Inositol is generally considered safe during pregnancy, and some studies suggest it may reduce the risk of GDM in women with PCOS. The

typical approach is to continue the same 40:1 ratio supplement at the same dose. Always run any supplement by your obstetrician.

Weight gain guidelines: If you start pregnancy with a higher body weight, your weight gain targets may differ from standard recommendations. Your doctor can help you determine the appropriate weight gain range for your situation.

Postpartum Considerations

After pregnancy, PCOS does not simply disappear. In fact, the postpartum period can bring significant changes — both positive and challenging.

PCOS symptoms after pregnancy: Some women find that their PCOS symptoms improve after pregnancy and breastfeeding. This may be due to hormonal changes, weight loss from breastfeeding, or simply the passage of time. Other women find that symptoms return or worsen. There is no way to predict which category you will fall into.

Breastfeeding and PCOS: Women with PCOS may face additional challenges with breastfeeding. The hormonal imbalances can affect milk supply — both the elevated androgens and the insulin resistance may impair lactation. However, most women with PCOS are able to breastfeed successfully, especially with support from a lactation consultant. Breastfeeding also has metabolic benefits — it burns extra calories, improves insulin sensitivity, and may reduce the risk of type 2 diabetes later in life.

Metabolic health after pregnancy: Pregnancy itself is a metabolic stress test. Women with PCOS have a higher risk of developing type 2 diabetes after a pregnancy complicated by GDM. If you had GDM, you should have a glucose tolerance test 6 to 12 weeks after delivery, and then annually thereafter. Maintaining the nutrition and exercise habits you built during pregnancy is important for long-term metabolic health.

Fertility after pregnancy: If you want more children, your fertility after a first pregnancy may be different from your fertility before. Some women ovulate more regularly after a full-term pregnancy; others return to their pre-pregnancy anovulatory pattern. If you are breastfeeding, ovulation may be suppressed, but that is not reliable for contraception or for fertility planning.

The Emotional Side of PCOS and Pregnancy

Pregnancy after PCOS can bring complicated emotions. There is often joy mixed with anxiety — the joy of finally achieving pregnancy, and the anxiety of "what if it goes wrong" after years of struggling with infertility. This is normal. Allow yourself to feel both.

If you have experienced pregnancy loss, you may carry that grief into a subsequent pregnancy. The fear of another loss can be overwhelming. Seek support — a therapist, a support group, or a compassionate obstetrician who understands pregnancy after loss.

And if you are reading this chapter and not yet pregnant, the waiting period — the months or years between deciding to start trying and actually conceiving — is some of the hardest terrain. Be gentle with yourself. You are not alone.

The Partner's Role in PCOS Pregnancy Planning

If you have a partner, it can help to include them in the PCOS pregnancy planning process. Fertility treatment can feel isolating — the blood draws, the ultrasounds, the timed intercourse, the negative pregnancy tests — and partners often do not know how to help. A few ways a partner can be genuinely supportive include: attending doctor's appointments, tracking medication schedules and appointments, being emotionally present without trying to "fix" the sadness of negative results, offering practical support during the emotional lows, and educating themselves about PCOS so they understand what you are going through.

If you are navigating fertility alone (single by choice or circumstance), it can feel even harder. Building a support network of friends, family, or an online PCOS community is especially important. You do not have to do this alone. Many women navigate PCOS fertility treatment independently and build wonderful families. The road may be different, but the destination is the same.

The PCOS-Friendly Pregnancy Checklist

Once you conceive, here is a checklist to review with your obstetrician: continue metformin if you were taking it (it reduces miscarriage risk and GDM risk), continue inositol (safe and may reduce GDM risk), schedule early glucose screening (before the standard 24–28 week test), monitor blood pressure closely (preeclampsia risk is elevated), ask about low-dose aspirin for preeclampsia prevention if you have additional risk factors, and have a postpartum glucose tolerance test scheduled for 6–12 weeks after delivery. Print this list and bring it to your first prenatal appointment.

Postpartum Hormonal Transitions

After you give birth, your hormones undergo a dramatic shift. The placenta is gone, which removes a major source of hormones, and if you breastfeed, prolactin suppresses ovulation. Some women with PCOS find that their symptoms improve significantly after pregnancy — their insulin sensitivity resets, their cycles become more regular, and their androgen levels drop. Others find that PCOS returns or even worsens, particularly if significant weight was gained during pregnancy or if sleep deprivation and stress cause metabolic disruption.

The postpartum period is a high-risk time for mental health, and women with PCOS are at elevated risk for postpartum depression. The combination of hormonal shifts, sleep deprivation, the physical recovery from childbirth, and the demands of caring for a newborn can be overwhelming. If

you have a history of depression or anxiety, talk to your doctor before delivery about a postpartum mental health plan. Early intervention is far more effective than waiting until you are in crisis.

Toolkit Summary

- Letrozole is first-line ovulation induction for PCOS; clomiphene is an alternative
- Metformin and inositol improve egg quality and may reduce miscarriage risk
- Women with PCOS have higher rates of GDM (2-4x), preeclampsia, and preterm birth — talk to your doctor about early screening
- Metformin is safe to continue during pregnancy and may reduce pregnancy loss and GDM risk
- Postpartum PCOS is unpredictable — some women improve, others return to their pre-pregnancy pattern
- Breastfeeding challenges are real but manageable with support — most women with PCOS can breastfeed successfully
- If you had GDM, get a postpartum glucose tolerance test and annual diabetes screening
- The emotional journey of PCOS and pregnancy is real — seek support for the hard parts and allow yourself to feel the joy

ch11

BY NOW, YOU HAVE LEARNED A LOT ABOUT PCOS. YOU UNDERSTAND THE HORMONAL MECHANISMS, THE INSULIN CONNECTION, THE DIAGNOSTIC PROCESS, THE NUTRITION AND MOVEMENT STRATEGIES, THE MEDICATIONS AND SUPPLEMENTS, AND THE MENTAL HEALTH DIMENSIONS. YOU HAVE THE KNOWLEDGE.

This chapter is about putting it all together into a long-term management plan that actually fits your life. Because knowing what to do is different from doing it consistently, and doing it consistently is different from doing it for years or decades.

Living full with PCOS — not despite it, but with it as a manageable part of your life — requires a system. Not perfection. A system.

The Comprehensive PCOS Plan: The Five Pillars

Your PCOS management plan rests on five pillars. These are not optional extras — they are the foundation of long-term metabolic and hormonal health. Every woman with PCOS needs all five, though the specifics of each pillar will be different for every person.

Pillar 1: Nutrition Foundation. The goal is not a restrictive diet. It is a consistent pattern of eating that stabilizes blood sugar, supports insulin sensitivity, and reduces inflammation. For most women, this means a Mediterranean-style diet with plenty of vegetables, protein, healthy fats, and moderate amounts of high-fiber carbohydrates. Not keto. Not fasting. Not deprivation. Consistent, satisfying, nutrient-dense eating that works with your metabolism, not against it.

Pillar 2: Movement. The goal is not extreme exercise. It is consistent, moderate movement that includes resistance training, steady-state cardio, and stress-reducing practices. Three to five sessions per week, most of them at moderate intensity, is the target. This is not negotiable for long-term health — but it also does not have to be punishing.

Pillar 3: Medications and Supplements. Some women with PCOS need medication to manage insulin resistance, lower androgens, or address specific symptoms. Others manage well with supplements alone. The right regimen for you depends on your phenotype, your symptom burden, your goals, and your preferences. Work with your doctor to find the combination that works. Review it annually.

Pillar 4: Stress Management. Stress is not a soft concern — it is a biological driver of PCOS. Cortisol worsens insulin resistance, increases androgen production, and disrupts ovulation. You need consistent stress management practices: adequate sleep, boundaries, leisure time, mindfulness, therapy if needed, and enough rest to avoid burnout. This pillar is as important as the nutrition pillar, even though no doctor will prescribe it.

Pillar 5: Symptom Tracking and Monitoring. You cannot manage what you do not measure. Tracking your menstrual cycle, your symptoms, your labs, and your treatment response gives you the data you need to make informed decisions and recognize when something needs to change.

The Annual Health Review

Once you have a stable PCOS management plan, you should review it annually — at minimum. Your annual review should include:

Blood work: - Fasting insulin - Fasting glucose - HbA1c - Lipid panel (total cholesterol, LDL, HDL, triglycerides) - Liver function (ALT, AST) — PCOS increases NAFLD risk - Vitamin D - Thyroid panel (TSH) - SHBG, testosterone (if symptoms are active)

Vital signs: - Blood pressure - Weight - Waist circumference

Symptom review: - Cycle regularity (how many periods in the past year?) - Acne, hirsutism, hair loss (worse, same, or better?) - Energy and mood - Sleep quality

Other screening: - Blood sugar screening — if you had GDM, annual glucose tolerance test - Endometrial assessment — if you are having fewer than four periods per year, ultrasound or biopsy to check for endometrial hyperplasia

Your annual review is also the time to reassess your medication and supplement regimen. What worked last year may not be optimal this year. Your body changes. Your life changes. Your PCOS management should change with them.

Monitoring for Long-Term Risks

PCOS is a lifetime condition, and it carries long-term health risks that need monitoring.

Type 2 diabetes: Women with PCOS have a 5-to-10 times higher risk of developing type 2 diabetes compared to women without PCOS. The risk is highest in women with phenotypes A and B. Fasting insulin, HbA1c, and glucose tolerance testing should be part of your annual screening.

Cardiovascular disease: The insulin resistance, dyslipidemia (low HDL, high triglycerides), and inflammation of PCOS contribute to accelerated cardiovascular risk. Women with PCOS have higher rates of hypertension and cardiovascular events. Blood pressure and lipid monitoring should be annual.

Endometrial cancer: Women with PCOS who have fewer than four periods per year and do not take any medication that induces regular bleeding (birth control or cyclic progesterone) have an elevated risk of endometrial hyperplasia and, over time, endometrial cancer. If you are not on birth control and have infrequent periods, you need an ultrasound and potentially an endometrial biopsy every year or two. Taking cyclic progesterone (10 to 14 days per month) to induce a period is an option if you prefer not to take birth control.

NAFLD: Non-alcoholic fatty liver disease affects up to 50 to 70 percent of women with PCOS. Annual liver function tests (ALT, AST) are essential. If you have elevated liver enzymes, a liver ultrasound can assess fatty liver severity. Treating insulin resistance is the primary intervention for NAFLD — the same nutrition and exercise strategies that help PCOS help the liver.

The Flare Protocol: What to Do When Symptoms Worsen

PCOS flares — periods when your symptoms worsen — are common. They are often triggered by: - Significant life stress (job change, relationship stress, grief, moving) - Weight gain (even a few pounds can worsen insulin resistance) - Disrupted routines (travel, illness, schedule changes) - Coming off birth control - Perimenopause transition - Significant changes in sleep or exercise patterns

When you recognize a flare, do not panic. That only raises cortisol and makes things worse. Instead, follow this protocol:

Step 1: Review your fundamentals. Have your sleep, nutrition, movement, or stress management slipped? If so, the first line of defense is returning to your baseline habits.

Step 2: Check your labs. If your symptoms have worsened for more than two months, run your annual lab panel to see if something has changed metabolically or hormonally.

Step 3: Talk to your doctor. If labs show changes, or if your symptoms are severe, your doctor may recommend adjusting medications, adding a medication you are not currently taking, or considering a new approach.

Step 4: Be patient. After you make a change, it takes at least two to three months to see improvement in most PCOS symptoms. Do not expect one week of clean eating to reverse symptoms that took years to develop.

The Identity Shift PCOS Requires

Living with PCOS requires a shift in how you think about health. It moves from "fix it and forget it" to "manage it forever." That can feel discouraging at first, but it is actually liberating.

You are not waiting to be cured. You are not waiting for your body to get back to "normal." There is no "before PCOS" that you are trying to return to. There is only now — managing a chronic condition well, living a full life alongside it, and accepting that the work of management is ongoing.

When you stop waiting for a cure, you stop postponing your life. You start living now, with PCOS, because you recognize that your life is not on hold until your symptoms disappear. Your life is happening right now, and it can be full and rich and meaningful even with acne and insulin resistance and irregular periods.

A Final Letter to the Reader

If I could tell you one thing, it would be this: PCOS is not your identity. It is a condition you manage, and you are capable of managing it well.

You are not "a PCOS patient." You are a person who happens to have a hormonal condition. You are not defined by your acne, your weight, your fertility challenges, or your insulin resistance. You are defined by what you do — how you show up for yourself, how you manage your health, how you live your life.

PCOS has given you something valuable, even if it does not feel that way. It has forced you to understand your body in a way most people never do. You know more about insulin, androgens, and metabolic health than most doctors knew ten years ago. You have developed discipline and self-awareness and the ability to advocate for yourself in medical settings. These are not small things.

The road ahead will have ups and downs. There will be good months and bad months. There will be periods when your symptoms are well-controlled and times when they flare. That is the nature of a chronic condition. It does not mean you are failing. It means you are human.

Keep the plan flexible. Keep your self-compassion strong. Lean on your people. Keep showing up for your annual labs. Keep asking questions. Keep advocating for yourself. And when you feel discouraged, remember that you are not alone. One in ten women has PCOS. You are part of a community, whether you have found it yet or not.

You can do this. You are doing it right now, by reading this book, by learning, by refusing to accept that "this is just how it is." That takes courage. And that courage will carry you through.

Communication Tips for Your Support System

One practical tool that can make living with PCOS easier is learning how to communicate about it with the people in your life. Many women with PCOS feel like they have to explain themselves constantly — why they eat differently, why they cannot make late social plans, why they are tired, why they are not pregnant yet. You do not have to explain everything to everyone. But having a short, clear explanation for the people who matter can reduce the mental load.

Try something like: "I have a hormonal condition called PCOS that affects my metabolism and hormones. It means I need to eat regular meals that keep my blood sugar stable, I need to prioritize sleep, and some things that seem simple (like losing weight) are biologically harder for me. I am managing it well with my doctor, but some days are harder than others. I really appreciate your support." A concise explanation that does not invite debate is powerful. You do not have to defend your condition or your choices. You just need the people around you to understand enough to be supportive.

Celebrating Your Wins, Big and Small

Living with PCOS requires consistent effort, and that effort deserves acknowledgment. When you go a month without a new cystic acne breakout — celebrate it. When you complete your annual lab work — celebrate it. When you have a period that comes on time — yes, celebrate it. When you do the right things consistently, even if you do not see immediate changes — recognize that consistency is itself the win.

Create a simple way to track your wins. It could be a note on your phone, a journal entry each week, or a checkmark on a calendar. The act of noticing and recording progress — even small progress — rewires your brain to see your efforts as effective. This is not about toxic positivity. It is about training yourself to recognize the real, measurable improvements that happen when you manage a chronic condition well. You are doing hard work, and it is working.

Toolkit Summary

- Your PCOS plan rests on five pillars: nutrition, movement, medications/supplements, stress management, and symptom tracking
- Schedule an annual health review with fasting insulin, HbA1c, lipids, liver function, vitamin D, blood pressure, and symptom assessment
- Monitor long-term risks: type 2 diabetes (annual screening), cardiovascular disease (annual lipids and BP), endometrial health (if fewer than 4 periods/year), and NAFLD (annual liver function)

- When symptoms flare, review fundamentals, check labs, talk to your doctor, and be patient — changes take months
- PCOS is a lifetime condition, not a waiting period — do not postpone your life until symptoms disappear
- PCOS is not your identity — it is a condition you manage, and you are capable of managing it well
- Talk to your doctor before making any significant changes to your management plan

resources

THIS SECTION PROVIDES PRACTICAL TOOLS AND REFERENCES TO SUPPORT YOUR PCOS JOURNEY. THE GLOSSARY, TRACKERS, AND COMPARISON CHARTS ARE DESIGNED TO HELP YOU COMMUNICATE MORE EFFECTIVELY WITH YOUR DOCTOR, TRACK YOUR PROGRESS, AND MAKE INFORMED DECISIONS.

Glossary

17-OHP (17-hydroxyprogesterone): A hormone precursor that is elevated in non-classic congenital adrenal hyperplasia (NCAH), a condition that mimics PCOS.

Acanthosis nigricans: Dark, velvety patches of skin (typically on the back of the neck, armpits, and skin folds) that indicate significant insulin resistance.

Amenorrhea: The absence of menstrual periods for three or more months.

Androstenedione: An androgen hormone produced by the ovaries and adrenal glands; a precursor to testosterone.

Anovulation: The absence of ovulation — when the ovary does not release an egg during a menstrual cycle.

DHEA-S (Dehydroepiandrosterone sulfate): An androgen hormone produced primarily by the adrenal glands; often elevated in PCOS.

FSH (Follicle-stimulating hormone): A pituitary hormone that stimulates ovarian follicle development.

GLP-1 (Glucagon-like peptide-1): A naturally occurring hormone that stimulates insulin secretion and promotes satiety; medications that mimic it (GLP-1 agonists) are emerging treatments for PCOS.

HbA1c (Hemoglobin A1c): A blood test that measures average blood sugar levels over the past two to three months.

Hirsutism: Excess hair growth in a male pattern (face, chest, back, abdomen) in women, caused by elevated androgens.

Hyperandrogenism: The presence of elevated androgen levels, either on blood work (biochemical) or in physical symptoms (clinical).

Insulin resistance: A condition in which cells do not respond properly to insulin, requiring the pancreas to produce more insulin to maintain normal blood sugar.

LH (Luteinizing hormone): A pituitary hormone that triggers ovulation; typically elevated in PCOS relative to FSH.

Letrozole: An aromatase inhibitor used as first-line ovulation induction medication for women with PCOS.

Metabolic syndrome: A cluster of conditions (abdominal obesity, high triglycerides, low HDL, high blood pressure, elevated blood sugar) that increase the risk of heart disease and diabetes.

Metformin: An insulin-sensitizing medication commonly used to treat insulin resistance in PCOS.

NAFLD (Non-alcoholic fatty liver disease): Fat accumulation in the liver not caused by alcohol; strongly associated with insulin resistance and PCOS.

OHSS (Ovarian hyperstimulation syndrome): A potentially serious complication of fertility treatment in which the ovaries become swollen and painful; PCOS increases the risk.

Oligomenorrhea: Infrequent menstrual periods — cycles longer than 35 days.

PCOS (Polycystic Ovary Syndrome): A common endocrine disorder characterized by irregular periods, elevated androgens, and polycystic ovaries, driven largely by insulin resistance.

Rotterdam criteria: The internationally accepted diagnostic criteria for PCOS, requiring 2 of 3: irregular ovulation, hyperandrogenism, and/or polycystic ovaries on ultrasound.

SHBG (Sex hormone-binding globulin): A protein produced by the liver that binds to sex hormones, keeping them inactive; low SHBG is common in PCOS and results in more free, active testosterone.

Spirolactone: An anti-androgen medication used to treat acne, hirsutism, and androgenic alopecia in PCOS; requires concurrent birth control due to teratogenicity.

Lab Tracker Template

Use this template to track your lab results over time. Bring it to your annual health review.

Test	Date:	Date:	Date:	Target Range
Fasting insulin				<8-10 mIU/L
Fasting glucose				<100 mg/dL
HbA1c				<5.7%
Total testosterone				15-70 ng/dL
Free testosterone				Lab-specific
SHBG				40-120 nmol/L
DHEA-S				35-430 mcg/dL
LH				Lab-specific
FSH				Lab-specific
LH/FSH ratio				<2:1
Total cholesterol				<200 mg/dL
LDL cholesterol				<100 mg/dL
HDL cholesterol				>50 mg/dL
Triglycerides				<150 mg/dL
ALT				<35 U/L
AST				<35 U/L
Vitamin D				30-50 ng/mL
TSH				0.5-4.5 mIU/L
Ferritin				30-150 ng/mL

Supplement Cheat Sheet

Supplement	Typical Dose	Evidence Level	Notes
Myo-inositol	2–4 g/day	Strong	Combine with DCI in 40:1 ratio
D-chiro-inositol	50–100 mg/day	Strong (with MI)	Not effective alone at high doses
Vitamin D	1,000–4,000 IU/day	Strong	Test levels first; most women need this
Omega-3 (fish oil)	2 g EPA+DHA/day	Moderate	Improves triglycerides and inflammation
Magnesium glycinate	200–400 mg/day	Moderate	Improves insulin sensitivity and sleep
Zinc	20–30 mg/day	Moderate	Take with food; can cause nausea
Chromium picolinate	200–1,000 mcg/day	Modest	May improve glucose tolerance
NAC	1,200–1,800 mg/day	Modest	Comparable to metformin for ovulation
Berberine	500 mg 2–3x/day	Moderate	AMPK activator; take with meals
Spearmint tea	2 cups/day	Preliminary	May lower free testosterone

Medication Comparison Chart

Medication	Primary Use	How It Works	Typical Timeline to Benefit	Key Cautions
Metformin	Insulin resistance	Reduces liver glucose output, improves muscle glucose uptake	4-12 weeks	GI side effects; check B12 annually
Letrozole	Ovulation induction	Lowers estrogen, increases FSH	Within one cycle	Requires monitoring; OHSS risk
Clomiphene	Ovulation induction	Blocks estrogen receptors, increases FSH	Within one cycle	Higher twin rate; uterine lining thinning
Spirolonactone	Acne, hirsutism, hair loss	Blocks androgen receptors	2-6 months	Teratogenic — requires birth control; check potassium
Semaglutide/ Tirzepatide			4-12 weeks	

Medication	Primary Use	How It Works	Typical Timeline to Benefit	Key Cautions
	Insulin resistance, weight	GLP-1 agonist (and GIP for tirzepatide)		GI side effects; cost; long-term commitment
Birth control pill	Symptom management	Suppresses ovarian function	1-3 months	Masks underlying condition; clot risk; may worsen insulin

Symptom Tracking Template

Date	Bleeding (days)	Acne (1-5)	Hair growth (1-5)	Hair shedding (1-5)	Energy (1-5)	Mood (1-5)	Sleep (hrs)	Stre (Y/N)

Scale: 1 = none/very good, 5 = severe/very poor

Sample Meal Plans (Mediterranean-Based, Low-GI)

Breakfast Options: - Greek yogurt (plain) with berries, flaxseed, and a sprinkle of cinnamon - Two scrambled eggs with spinach, mushrooms, and a slice of whole-grain toast - Overnight oats (steel-cut) with chia seeds, almond milk, and berries - Smoothie with protein powder, spinach, frozen berries, unsweetened almond milk, and flaxseed

Lunch Options: - Large salad with leafy greens, grilled chicken, chickpeas, cucumber, tomato, avocado, and olive-oil vinaigrette - Lentil soup with a side of roasted vegetables - Quinoa bowl with roasted broccoli, salmon, tahini dressing, and pickled onion - Turkey and avocado wrap in a whole-grain tortilla with mixed greens

Dinner Options: - Baked salmon with roasted Brussels sprouts and a small sweet potato - Stir-fried tofu or chicken with broccoli, bell peppers, snap peas, and cauliflower rice - Grass-fed beef patty (no bun) on a bed of arugula with roasted asparagus and sweet potato wedges - Mediterranean baked cod with tomatoes, olives, capers, and a side of sautéed spinach

Snack Options: - Apple slices with almond butter - Handful of walnuts and a small piece of dark chocolate - Hard-boiled egg - Celery with hummus - Edamame - Small bowl of berries with a tablespoon of pumpkin seeds

When to Use These Resources

This resources section is designed to be a living document — something you return to as your needs change. When you are first diagnosed, start with the glossary to understand the terminology your doctor uses, then use the lab tracker template at your next appointment. When you are trying to decide whether to try a supplement, consult the supplement cheat sheet and discuss it with your doctor. When you are tracking symptoms, print out the tracking template and keep it by your bed. When you are struggling with what to cook, the meal plans provide a starting point.

Your body and your PCOS will change over time. What works at twenty-four may need adjustment at thirty-four. The resources here are designed to be flexible enough to adapt. Bookmark this section, update your lab tracker each year, and revisit the cheat sheets whenever you are considering a new treatment approach. Managing PCOS is an ongoing process, and the right tools make it manageable.

Common Medications — Expanded Notes

Metformin ER (extended release): If you experience significant gastrointestinal side effects with standard metformin, ask your doctor about the extended-release formulation. It is often much better tolerated and can be taken with the largest meal of the day. Starting at 500 mg once daily for a week, then increasing to 500 mg twice daily,

and finally to 1,000 mg twice daily (the typical therapeutic dose) can minimize side effects. Some women do well at lower doses — 1,000 mg total per day may be sufficient.

Letrozole protocol: Ovulation induction with letrozole typically starts at 2.5 mg taken on cycle days 3 through 7. If ovulation does not occur, the dose can be increased to 5 mg or 7.5 mg in subsequent cycles. Monitoring with ultrasound and blood work is important to track follicle development and time intercourse. Most women ovulate on one of these doses within two cycles.

Spirolactone titration: Starting at 25 mg once daily and increasing by 25 mg every two weeks up to a maintenance dose of 100-200 mg per day reduces side effects. It takes about six to twelve months to see full effects on hair loss and hirsutism. Spirolactone is also available in topical form for some insurance plans, which avoids systemic side effects.

Inositol dosing schedule: Most women take 2 grams of myo-inositol plus 50 mg of D-chiro-inositol twice daily — once in the morning and once in the evening. This keeps levels steady throughout the day. Powder forms can be mixed into water or unsweetened beverages; capsules are also available. Taking inositol on an empty stomach may cause mild nausea in some women — taking it with food can help.

Body Weight and PCOS — Additional Context

Weight is one of the most charged topics in PCOS care, and it deserves additional context. Many women with PCOS carry excess body weight, and weight loss often improves

PCOS symptoms — but the relationship is bidirectional. PCOS makes weight loss harder, and excess weight worsens PCOS. Breaking this cycle requires addressing the metabolic dysfunction, not just the weight.

Research suggests that a weight loss of 5 to 10 percent of starting body weight can significantly improve PCOS symptoms, including ovulation rates, insulin sensitivity, and androgen levels. This is a modest amount — for a 200-pound woman, that is 10 to 20 pounds. You do not need to reach an "ideal" body weight to see benefits. Even small reductions in weight improve metabolic health.

However, weight should never be the only measure of success. Many women with PCOS improve their insulin sensitivity, restore ovulation, reduce their androgen levels, and improve their quality of life without significant weight loss. These metabolic improvements are just as valuable as weight loss, and they often happen first. If you are doing the right things for your PCOS — eating well, moving regularly, managing stress, taking appropriate medications — and your body weight is not changing dramatically, you are still getting healthier. Your labs, your cycles, and your symptoms are the true measures of success.

Further Reading

Books: - The PCOS Plan: Prevent and Reverse Polycystic Ovary Syndrome through Diet and Fasting by Jason Fung and Nadia Brito Pateguana - The Period Repair Manual by Lara Briden - The PCOS Workbook: Your Guide to Complete Physical and Emotional Health by Angela Grassi - The Fifth

Vital Sign: Master Your Cycles and Optimize Your Fertility by Lisa Hendrickson-Jack - PCOS SOS: A Gynecologist's Lifeline to Naturally Restore Your Rhythms, Hormones, and Happiness by Felice Gersh

Websites: - PCOS Awareness Association (pcosaa.org) - PCOS Challenge (pcoschallenge.org) - The PCOS Nutrition Center (pcosnutrition.com) - The Hormone Network — Endocrine Society (hormone.org)

Online Communities: - r/PCOS on Reddit — an active, supportive community of over 200,000 members - PCOS Dietitian Support Group on Facebook - PCOS Warrior (pcOSwarrior.com) — blog and community - The PCOS Protocol on Instagram and YouTube

Organizations: - PCOS Challenge: The National Polycystic Ovary Syndrome Association - The Polycystic Ovary Syndrome Association (PCOSAA) - Endocrine Society — Patient resources on PCOS - American Society for Reproductive Medicine (ASRM) — Fertility and PCOS resources

About the Author

AUTHOR BIO GOES HERE

Also by AUTHOR_NAME

This is her first book.

