

Tools Fill & Sign Comment

a. Are more numerous in antigravity muscles.

c. The striated portion is innervated by afferent motor fibres.

2. In Wallerian degeneration,

b. Chromatolysis occurs.

c. Myelin sheath shows widening of nodes of Ranvier.

d. Schwann cells decrease in number.

3. Interfascicular oligodendrocytes

- a. Are phagocytic cells.
- b. Cytoplasm is fibrillary.
- c. Share in the formation of blood brain barrier.
- d. Are ectodermal in origin.

4. Sympathetic ganglia are characterized by one of the following:

- a. Cells are stellate multipolar.
- b. Many satellite cells.
- c. No synapse between cells.
- d. Cells are separated by myelinated nerve fibres.



Type here to search



ENG

11:50 AM
10/2/2025



5. The largest neuroglia cells:

- a. Are also called mesoglia cells.
- b. Share in the formation of the blood brain barrier.
- c. Are concerned with myelin production.
- d. Are mesodermal in origin.

6. Regarding the peripheral nerve:

- a. The nerve is covered by dense connective tissue called perineurium.
- b. Epineurium is formed of epithelium like cells joined by gap junctions.
- c. The endoneurium forms a barrier to protect the nerve cells.
- d. Sheath of Henle consists of reticular fibers formed by Schwann cells.

7. Nonmyelinated nerve fiber with sheath of schwann cells is found in:

- a. Post ganglionic sympathetic nerve.
- b. Grey matter.
- c. White matter.
- d. Peripheral nerve.

8. Which of the following is true regarding oligodendrocytes:

- a. Satellite oligodendrocytes support nerve cells

8. Which of the following is true regarding oligodendrocytes:

- a. Satellite oligodendrocytes support nerve cells.
- b. They have no centrioles.
- c. Endodermal in origin.
- d. Interfascicular oligodendrocytes are present in grey matter.

9. Regarding the nerve endings, which of the following statements is true?

- a. Free nerve endings in epithelium are receptors for vibration and pressure.
- b. In tendon spindle, sensory nerve ends around the collagen bundles.
- c. Muscle spindles are present in smooth muscle.
- d. Ruffini corpuscles are non-capsulated rounded bodies.

- a. Less common type than electrical synapse.
- b. Axon terminal is rich in mitochondria & synaptic vesicles of transmitter.
- c. Synaptic cleft is 2-3 nm in width, through which ions must cross.
- d. Contains gap junctions between cells for conduction of neurotransmitters.