Historical Britannica: extracted source

Encyclopaedia Britannica; or, A dictionary of arts, sciences, and miscellaneous literature - Fourth edition, Volume 12 Part 2, MAH-Materia medica - EB.9

Source row 328. Parser: nvidia/nemotron-3.5-parse.

This is model-extracted text, not a corrected transcription. OCR errors, reading order and region alignment require human review.

Region region-0; provider order 0.

Pimasts. low B. Short-tailed, yellow, freched with black, with naked black face and bands, hairy on the upper farface–11. _S. Guerrao_, Guerous B. Short-tailed s cinceroy, with the crown jotted with yellow brown fare and pale bear–12. _Duncania_ is dead-ranked **B.** Short-tailed, alb brown, with bluish face, and ve- yr large deer teeth–13. _Va Feya_ Brown B. Short- ith tulled j brown, with coulition behind, a width face, and avry head tofore–15. _A Nugefrica_ Fig-tailed B. Olive brown with short-tailed tail–15. _Grifera_ Crefled B. Short talled , black, with very long hair or the crown and checks i, which break the bare face and hands–16. _S. Andria_ Little B. Short tailed y yellowish, without california behind, with chosen fineing side to the finger, and fromdella with rounded tails; the fingers with narrow ones–17. _Hamwyara_, Dog-East B. Trial ges, with calories behind Harghi draws, and the hairs on each side of the head very long–18. _S. Ferra_, Lion-tailed B. Tailley black, with very large whistling freeding -teard–19. _Cayafora_ Pale brown, bearded, with calificite behind, and with lough-filled-colored faces a whitelit land accord for received, and a loough sharp pointed tail–20. _S. Regeta_, Winkled B. Short, taled, yellowish browns, within beneath, with flesh-coloured eyes, and large blue-red wrinkles callicties behind:

Region region-1; provider order 1.

37 Monkeys.

Region region-2; provider order 2.

Abonkeys.

Region region-3; provider order 3.

21. _L Louane_, Leonardo Monkey. Black, with callotion behind, very large whistle heard, and very long tufed tail–23. _S. Gymalofar_, Harv-tipped B. Long, tailed, bearded, with colulates behind, rifup blind notifir, and schectail all–23. _Yeter_, Purple-faced M. Long-tailed, witch, with black bears–24. _Rolony_ Polkovny or Ralston. Born-tatalized, like iih; white beneath, with triangular face, furrounded by a white divided and cross–25. _S. Chiana_ Pinus and Spotted M. Long-tailed, blackibh, frechedk with white; the bar of the forehead and lizard growing, a potato form with a tunated band scrut the fore-head–26. _S. Naivia_, Long-nodel M. Long-tailed, blackibh's ply-p ath-coloured bereats, with long naked fish-colored faces. _S. Peregione_, Yellow ib M. Long-tailed, bearded, cineensely yellow, with black face and onto–28. _S. Salute_ Green Tue. Long, tailed, yellowish gray, with black face, and callions belined–29. _Zhlaya_, Mangbay on White-eyal M. Long-tailed, bearded, with black face, white eyelids, white frontal band, and the hair on the fore-head upple–20. _G Gehofer_ Multheck M. Long, tailed, blackibh rally, whithb breath, with bearded check and yellowblow crowns red eyelid and whittle marine–31. _S. Niltonen_ Willemed M. Long-tailed, beardife, thick, frechedk with white; the than of the hauled and teard growing, a potato form with a tunated band scrut the fore-head–26. _S. Naivia_, Long-nodel M. Long-tailed, olive coloured, bearded checks, and black ears, node, and foles–33. _S. Mason_ Negro M. Long-tailed, blackibh, free flyerfly fish-coloured face and breath and blackikh beard–23. _S. Aydade_ Eger M. Long-tailed, bear-de fetly grey, with sitting individual but on the crown–35. _S. Rubra_ Red M. Long-tailed, red pale alh-coloured beneath, with bearded cheeks, and a black or white band acre the forehead–36. _S. Sinica_ Chi

Region region-4; provider order 4.

nee M. Beardife, pale brown, with the hair of the Ribs of the forehead preceding round horizontally–37.

S. Freangri the Spanish ra, Vaulting M. Olive black, white berast, with a trimming color blue foot on the one–28. _S. Picenia_ Browned M. Ruth brown, whichily yellow beneath, with black limbs, and the hair in the last rings similarly upward–39. _S. More, Variet M. Olive rally_ white becets, with the cheeks bearded, and a hunted whilha band beside the forehead–40. _S. Naivia_ Pro-fusion R. Long-tailed, bearded, chatin colour, with pale limb and tail, and a very long one–41. _N. Av._ many, Cochin Chin Mm. Long-tailed, with bedral cheeks and white tail–42. _S. Fariva_, Dawyn M. Sub-ferraginous, with the bone part of the back orange, white bearing, with fed-chcolored face and earrs–43. _S. Hierina_ Gaost M. Long-tailed, brown, with blue furrowsers, and the long-pointed bend–43. _Come_. Pull-bottom to B. Long-tailed, black, with very long freezing whitil him on the back, and white tail–45. _S. Aenegaria_, Bay M. Long-tailed, ruby, with black limbs and tail–46. _S. Amalata_ Anmuhalted M. Reilly brown, whichbened the hair amabulated, tail, horses than the body–47. _S. Pichea_ Fox-tailed M. Blackikh brown, with the tips of the hair which, and very gray tail–48. _S. Lucenia_ Siratrician M. Long, tailed, with fireaginary ears; crooked hairy tail and the leaves, thin on the bone being rounded–49. _S. Angela_, Milky M. Long-tailed, nifky, with the black limbs and tail–46. _S. Amalata_ Ammuhalted M. Early brown, whichbened the hair amabulated, tail, horses than the body–49. _S. Pichea_ Fox-tailed M. Blackikh brown, with the tips of the hair which, and very gray tail–48. _S. Lucenia_ Siratrician M. Long, tailed, with fireaginary dairy ears; crooked hairy tail and the leaves, thin on the bone being rounded–49. _S. George_. Kulto, called M. Long red-tailed, bearded left, with the hair of the head breaking downwards, and thus pastall–20. _S. Naiphi_, silky M. Long-tailed, riby, with, only, reddish green, orange-colored feet, and point- ed tail–21. _S. Aprquarie_, Fair M. Long-tailed, bearded, white, with red face and brown tail–32. _S. Berladeki_, Precher M. Bearded, black, the feet and off the oil brown, tail prebehand–44. _So- Anisia_, Royal M. Long-tailed, bearded red jail prebeilk–47. _S. Kaniya_ Your-fingerend M. Long-tailed, black, with Rainangered Feet, tail prebeñike–56. _S. Fatualis_, Horned M. Long-tailed, bearded, with worma like rats on the head- tail prebeñike–47. _S. Pirpida_ Pearl M. Long-tailed, everdefts, with upright hair on the head, and bluish, tail prebeñike–56. _S. Adege_ Weger M. Long-tailed, bearded's brown body, black feet, and without calides, jail prebeñike–50. _S. Capu-cina_, Copacidin M. Long-tailed, beautiful, without calides, with black crown and limbs, and infloria pre-hedlin tail–60. _S. Sherner_, Squirit M. Long-tailed, yellowignity, beardedits, with orange-colored bands and feet, four of the clean, and the that feet, pointed out. _S. Alaguqigh_ Antique M. Blackfish brown, white bereach, with black limbs and face, bearded cheeks, and brown petite tail the–62. _S. Mora_, Nalda-tailed M. Long-tailed, bearded, brown, with dukly auxzie, and adda feel feyl tail.

Region region-5; provider order 5.

Spiceer i. _S. Augusta_ Om Oman, Wild Man of Satrea, the Woods. Chapranzes, Jacko. Taillets Appe–It Ona Orga is generally believed by antarially, that the animals which have been deficient under the mean given above, are only varieties of the fame facies, differing from each other in jazz, colors, fox, and fame other nothing fludes of discriminating. Four remarkable femmes

Region region-6; provider order 6.

Source record

bri-0328-4588acdac3cf

Image SHA-256: 4588acdac3cf6445b62cc57af5bf3158

9799caacfc5eebab02fdbedab2a528ad

Dataset revision: 309f765b1ecc83e0a7f7762e44b4a927a7555443

National Library of Scotland, Encyclopaedia Britannica Illustrated (CC0).

https://huggingface.co/datasets/NationalLibraryOfScotland/encyclopaedia_britannica_illustrated/viewer/default/train?row=328