Historical Britannica: extracted source

Encyclopaedia Britannica; or, A dictionary of arts, sciences, and miscellaneous literature - Fourth edition, Volume 11 Part 1, HYD-JUN - EB.9

Source row 2286. Parser: nvidia/nemotron-3.5-parse.

This is model-extracted text, not a corrected transcription. OCR errors, reading order and region alignment require human review.

Chap. IV.

Region region-0; provider order 0.

Abdominal Ploes

Region region-1; provider order 1.

Gen. 3. AANELFS.

Region region-2; provider order 2.

Anabope.

Region region-3; provider order 3.

Head (mewhat depreffed) month terminal) teeth (mail, and placed on the jaws) eyes protuberant, with double pupils; gil-mountaine ficayed.

Region region-4; provider order 4.

Forr good analysis. Colois smalloy of Limnus–

Region region-5; provider order 5.

Ibulmus: Yellowiti-gray, with longitudinal black lines on each fide. Length from fice to eight or for inches. general appearance like that of a lache, but its eyes differ from their of every known fifth, each being ap- premlyered into two differed eye, upon la a com- mon receptacle; on difection, however, this observation is found to apply only one has another half of the organ. This fish is a mustor of South America, principal feu- quenting the rivers of Surinam, near the fea-coafs.

Region region-6; provider order 6.

Anals.

Region region-7; provider order 7.

Heal bony, naked, rough, and furnished with futures: teeth acute, and clude in the jaw and palates; two beards at the nofs; gill membrane twelve-royed; body faly.

Region region-8; provider order 8.

culva. _Carolina mix—With a black foot at the tail) famil, of a roundish form, and delon enten. Inhalata Carolina.

Region region-9; provider order 9.

Sirama.

Region region-10; provider order 10.

Gen. 4. SILURUS.

Region region-11; provider order 11.

Hend maked, jays, broad, and compreffed; mouth fur silhed with beard; pape and throat wide; lips thick; jaws dentated; tongue thick, (mouch, and very thort; eye final); gil-membrane furnished hair from four to

factors rays; body elongated, compreffed, facile, covered with vidl lines; lateral line near the back; it be fifty of the pectoral fins, or of the der- fal fin, finy, and dentated backwax.

Region region-12; provider order 12.

glanis. _Siq, or European filursas—One folk dorfal fin, fix bearer. Growns are long length of right, or even on term feet, and to the weight of the hundred pounds; but its ordinary fine is from two to three our four feet. It is flagging, and and deposited in the soft bottom of the rivers which is frequent, with his mouth half-open, moving about the breedy, which she finally fiber stalking for women, by hold of, and an extraped. It inhabits the larger rivers of Europe, as well as these parts of Asia and Africa, but is ino high etitimism as a food.

Region region-13; provider order 13.

eflireus. _Fidelity filursas—With one altifoof dorfal fin, and fix trends. About trees include lung, vessels in the fore-parts, depreffed, and of a cinerous colour, with from blocking the towards the salt. It is found in some rivers in Africa, and when truck, gives a gallons thock, though not fo forage as that from the torpedo and gymnasts. It is due to food.

Region region-14; provider order 14.

eficia. _Afriei filursas—Brown, all-coloured beneath, with bearded longer than the body; farket left, and wherers rays in the anal fan. The young of this species are excluded in the form of large over, the integments of which they from bark, but others to the parent till the yolk is confirmed. Native of India.

Region region-15; provider order 15.

estus. _City filursas—one kinder and full of friends; recently away in the anal fan, and eight beards. Inhabit the far and rivers of North America, preying on all kinds of finalize follies, and not fearing even those of its own Vel. X1. East I.

Region region-16; provider order 16.

kind. Talks like an eel, and is much relished by the Abdominal Americans.

Region region-17; provider order 17.

The other facies are denominated militaris, larges, bericharge, terarum, gelatin, moldyta, himmatinas, flyt cinna, clarias, grandecimanilang, erythogerea, borra- chus, fufilris, vulatus, abbirindes, aficus, myusia, angulilaris, undatomasia, various, felis, cours, carrinus, stones, ciliatings, and blood.

Region region-18; provider order 18.

Gen. 5. PLATSYACUS.

Region region-19; provider order 19.

Halbit of filurus & mouth beneath, bearded with circhi; body facallets, depreffed; tail long, compreffled.

Region region-20; provider order 20.

Artshialural plungtures. Silura of globo de Limnus cusylpleo- Grows on the length of a four or more, has a very un-unsur, couth appearance, and is remarkable for the many imall acrobain incredible, or locksers, with which the body is ledes. Nature of the Indian fans and rivers.

Region region-21; provider order 21.

_Flaredal gingences—Brown, marked above by loss-cerraco- giutuinai warted lines, with front anal fin. Smaller,fu- and just elongated than the two preceding. Native of the Indian fees.

Region region-22; provider order 22.

Erd/baped playstaceus—Brown, with longitudinal argaille- white stripes, and the femoral dorsal, and emudral ris, fun united. Length weive up fifteen inches. Native of the Indian fees.

Region region-23; provider order 23.

Gen. 6. LORICATA.

Region region-24; provider order 24.

Head smooth, depreffed & mouth without teeth, and re- tradles; gil-membrane fix-ryed; body covered with a hard craft.

Region region-25; provider order 25.

_Ribbed laricaria. Silura colatia of Limnus—Vel-collate, landh brown, sealed with a single row of linhaile each fide; laid forked. This is a species of great strength and bolodos, which inhibits very painful and dangerous would with the future. It is native of the Indian and American fees.

Region region-26; provider order 26.

_Loyal berriesites—One dorlat fin, two beards] length case- about ten o twelve inches. Much alles to the pec-phroda, ceding, but has a rounded tail. Native of the Ameri- can fees.

Region region-27; provider order 27.

_Soldier laricaria. Siluras callicklay of Limnus—callicb- brown will dependful rounded head, and low offgs., fales on each fide, and roundtail tail. Native of South Ammer, where it is its considerable offers as a food. Spekkel laricaria—Yellow with bismuth buck pancillats, double row of fales on each fide ; fins freckilled with blacks; and darked tail, why fire out of inches long, but very elegant. Native of the river of Siberian.

Region region-28; provider order 28.

Tesulted laricaria—Longhemed y yellowish brown; denials, with roasted, bearable mouth, and highly pointed front. Native of the Indian fees.

Region region-29; provider order 29.

Teliber laricaria Juricaria prepfories of Limnus-flavors. They have, treated with borrowers to doralis first and tail marked by transferife bands. Native of the Indian fees.

Region region-30; provider order 30.

Gen. S. Salno.

Region region-31; provider order 31.

Head mouth and compreffed; mouth larges lips finally; tonque white, cardlinghamous, and movable; eyes midlife-fixed; lateral teeth in the jaws and tongue; gilb-membran furnished with four to ten rays; gill- N cover

Region region-32; provider order 32.

Source record

bri-2286-bf2427da4447

Image SHA-256: `bf2427da444769a49359f48f9473754c`

`b83b2796c186caccae883280aea98d72`

Dataset revision: `309f765b1ecc83e0a7f7762e44b4a927a7555443`

National Library of Scotland, Encyclopaedia Britannica Illustrated (CC0).

https://huggingface.co/datasets/NationalLibraryOfScotland/encyclopaedia_britannica_illustrated/viewer/default/train?row=2286