

This section introduces a few rules that allow us to differentiate a great variety of functions. By proving these rules here, we can differentiate functions without having to apply the definition of the derivative each time.

Powers, Multiples, Sums, and Differences

The first rule of differentiation is that the derivative of every constant function is zero.

RULE 1 Derivative of a Constant Function

If f has the constant value $f(x) = c$, then

$$\frac{df}{dx} = \frac{d}{dx}(c) = 0.$$

EXAMPLE 1

If f has the constant value $f(x) = 8$, then

$$\frac{df}{dx} = \frac{d}{dx}(8) = 0.$$

Similarly,

$$\frac{d}{dx}\left(-\frac{\pi}{2}\right) = 0 \quad \text{and} \quad \frac{d}{dx}(\sqrt{3}) = 0. \quad \blacksquare$$

Proof of Rule 1 We apply the definition of derivative to $f(x) = c$, the function whose outputs have the constant value c (Figure 1.8). At every value of x , we find that

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{c - c}{h} = \lim_{h \rightarrow 0} 0 = 0. \quad \blacksquare$$

The second rule tells how to differentiate x^n if n is a positive integer.

RULE 2 Power Rule for Positive Integers

If n is a positive integer, then

$$\frac{d}{dx} x^n = nx^{n-1}.$$

The Power Rule is actually valid for all nonzero real numbers n , which we prove in Section 1.7 (and in stages for various powers along the way). To apply the Power Rule, we subtract 1 from the original exponent (n) and multiply the result by n .

EXAMPLE 2 Interpreting Rule 2

f	x	x^2	x^3	x^4	\dots
f'	1	$2x$	$3x^2$	$4x^3$	\dots

RULE 3 Constant Multiple Rule

If u is a differentiable function of x , and c is a constant, then

$$\frac{d}{dx}(cu) = c \frac{du}{dx}.$$

In particular, if n is a positive integer, then

$$\frac{d}{dx}(cx^n) = cnx^{n-1}.$$

EXAMPLE 3

(a) The derivative formula

$$\frac{d}{dx}(3x^2) = 3 \cdot 2x = 6x$$

(b) A useful special case

The derivative of the negative of a differentiable function u is the negative of the function's derivative. Rule 3 with $c = -1$ gives

$$\frac{d}{dx}(-u) = \frac{d}{dx}(-1 \cdot u) = -1 \cdot \frac{d}{dx}(u) = -\frac{du}{dx}.$$



The next rule says that the derivative of the sum of two differentiable functions is the sum of their derivatives.

RULE 4 Derivative Sum Rule

If u and v are differentiable functions of x , then their sum $u + v$ is differentiable at every point where u and v are both differentiable. At such points,

$$\frac{d}{dx}(u + v) = \frac{du}{dx} + \frac{dv}{dx}.$$

EXAMPLE 4 Derivative of a Sum

$$\begin{aligned} y &= x^4 + 12x \\ \frac{dy}{dx} &= \frac{d}{dx}(x^4) + \frac{d}{dx}(12x) \\ &= 4x^3 + 12 \end{aligned}$$

The Sum Rule also extends to sums of more than two functions, as long as there are only finitely many functions in the sum. If u_1, u_2, \dots, u_n are differentiable at x , then so is $u_1 + u_2 + \dots + u_n$, and

$$\frac{d}{dx}(u_1 + u_2 + \dots + u_n) = \frac{du_1}{dx} + \frac{du_2}{dx} + \dots + \frac{du_n}{dx}.$$

EXAMPLE 5 Derivative of a Polynomial

$$\begin{aligned}y &= x^3 + \frac{4}{3}x^2 - 5x + 1 \\ \frac{dy}{dx} &= \frac{d}{dx}x^3 + \frac{d}{dx}\left(\frac{4}{3}x^2\right) - \frac{d}{dx}(5x) + \frac{d}{dx}(1) \\ &= 3x^2 + \frac{4}{3} \cdot 2x - 5 + 0 \\ &= 3x^2 + \frac{8}{3}x - 5\end{aligned}$$

Notice that we can differentiate any polynomial term by term, the way we differentiated the polynomial in Example 5. All polynomials are differentiable everywhere. ■

EXAMPLE 6 Finding Horizontal Tangents

Does the curve $y = x^4 - 2x^2 + 2$ have any horizontal tangents? If so, where?

Solution The horizontal tangents, if any, occur where the slope dy/dx is zero. We have,

$$\frac{dy}{dx} = \frac{d}{dx}(x^4 - 2x^2 + 2) = 4x^3 - 4x.$$

Now solve the equation $\frac{dy}{dx} = 0$ for x :

$$\begin{aligned}4x^3 - 4x &= 0 \\ 4x(x^2 - 1) &= 0 \\ x &= 0, 1, -1.\end{aligned}$$

The curve $y = x^4 - 2x^2 + 2$ has horizontal tangents at $x = 0, 1$, and -1 . The corresponding points on the curve are $(0, 2)$, $(1, 1)$ and $(-1, 1)$.

Derivatives of Exponential Functions

Derivative of the Natural Exponential Function

$$\frac{d}{dx}(e^x) = e^x$$

EXAMPLE 7 Finding a Tangent Line

Find an equation for a line that is tangent to the graph of $y = e^x$ and goes through the origin.

Solution Since the line passes through the origin, its equation is of the form $y = mx$, where m is the slope. If it is tangent to the graph at the point (a, e^a) , the slope is $m = (e^a - 0)/(a - 0)$. The slope of the natural exponential at $x = a$ is e^a . Because these slopes are the same, we then have that $e^a = e^a/a$. It follows that $a = 1$ and $m = e$, so the equation of the tangent line is $y = ex$. See (Figure 1.5) ■




Figure 1.5: the tangent line to the curve $y = e^x$

The derivative of the natural exponential function has the form:

$$\frac{de^{ax}}{dx} = a \cdot e^{ax}, \text{ } a \text{ is a constant}$$

The derivative of the general exponential function has the form:

$$\frac{da^x}{dx} = a^x \cdot \ln(a), \text{ } a \text{ is a constant}$$

Example 8:

Find the derivative of the functions $f(x) = 2 \cdot e^{-3x}$ and $g(x) = 5^x$

Solution:

$$f'(x) = 2 \cdot (-3) \cdot e^{-3x} = -6 \cdot e^{-3x}$$

$$g'(x) = 5^x \cdot \ln(5)$$

Products and Quotients

RULE 5 Derivative Product Rule

If u and v are differentiable at x , then so is their product uv , and

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}.$$

The derivative of the product uv is u times the derivative of v plus v times the derivative of u . In *prime notation*, $(uv)' = uv' + vu'$. In *function notation*,

$$\frac{d}{dx}[f(x)g(x)] = f(x)g'(x) + g(x)f'(x).$$

EXAMPLE 8 Using the Product Rule

Find the derivative of

$$y = \frac{1}{x}(x^2 + e^x).$$

Solution We apply the Product Rule with $u = 1/x$ and $v = x^2 + e^x$:

$$\begin{aligned} \frac{d}{dx} \left[\frac{1}{x}(x^2 + e^x) \right] &= \frac{1}{x}(2x + e^x) + (x^2 + e^x) \left(-\frac{1}{x^2} \right) \\ &= 2 + \frac{e^x}{x} - 1 - \frac{e^x}{x^2} \end{aligned}$$

$$\begin{aligned} \frac{d}{dx}(uv) &= u \frac{dv}{dx} + v \frac{du}{dx}, \text{ and} \\ \frac{d}{dx} \left(\frac{1}{x} \right) &= -\frac{1}{x^2} \text{ by} \\ &\text{Example 3, Section 1.7} \end{aligned}$$

EXAMPLE 10 Differentiating a Product in Two Ways

Find the derivative of $y = (x^2 + 1)(x^3 + 3)$.

Solution

(a) From the Product Rule with $u = x^2 + 1$ and $v = x^3 + 3$, we find

$$\begin{aligned}\frac{d}{dx}[(x^2 + 1)(x^3 + 3)] &= (x^2 + 1)(3x^2) + (x^3 + 3)(2x) \\ &= 3x^4 + 3x^2 + 2x^4 + 6x \\ &= 5x^4 + 3x^2 + 6x.\end{aligned}$$

(b) This particular product can be differentiated as well (perhaps better) by multiplying out the original expression for y and differentiating the resulting polynomial:

$$\begin{aligned}y &= (x^2 + 1)(x^3 + 3) = x^5 + x^3 + 3x^2 + 3 \\ \frac{dy}{dx} &= 5x^4 + 3x^2 + 6x.\end{aligned}$$

This is in agreement with our first calculation. ■

RULE 6 Derivative Quotient Rule

If u and v are differentiable at x and if $v(x) \neq 0$, then the quotient u/v is differentiable at x , and

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}.$$

In function notation,

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g(x)f'(x) - f(x)g'(x)}{g^2(x)}.$$

EXAMPLE 11 Using the Quotient Rule

Find the derivative of

$$(a) \quad y = \frac{t^2 - 1}{t^2 + 1}, \quad (b) \quad y = e^{-x}.$$

Solution(a) We apply the Quotient Rule with $u = t^2 - 1$ and $v = t^2 + 1$:

$$\begin{aligned} \frac{dy}{dt} &= \frac{(t^2 + 1) \cdot 2t - (t^2 - 1) \cdot 2t}{(t^2 + 1)^2} & \frac{d}{dt} \left(\frac{u}{v} \right) &= \frac{v(du/dt) - u(dv/dt)}{v^2} \\ &= \frac{2t^3 + 2t - 2t^3 + 2t}{(t^2 + 1)^2} \\ &= \frac{4t}{(t^2 + 1)^2}. \end{aligned}$$

$$(b) \quad \frac{d}{dx}(e^{-x}) = \frac{d}{dx} \left(\frac{1}{e^x} \right) = \frac{e^x \cdot 0 - 1 \cdot e^x}{(e^x)^2} = \frac{-1}{e^x} = -e^{-x}$$

RULE 7 Power Rule for Negative IntegersIf n is a negative integer and $x \neq 0$, then

$$\frac{d}{dx}(x^n) = nx^{n-1}.$$

EXAMPLE 12

$$(a) \quad \frac{d}{dx} \left(\frac{1}{x} \right) = \frac{d}{dx}(x^{-1}) = (-1)x^{-2} = -\frac{1}{x^2}$$

Agrees with Example 3, Section 1.7

$$(b) \quad \frac{d}{dx} \left(\frac{4}{x^3} \right) = 4 \frac{d}{dx}(x^{-3}) = 4(-3)x^{-4} = -\frac{12}{x^4}$$

EXAMPLE 13 Choosing Which Rule to Use

Rather than using the Quotient Rule to find the derivative of

$$y = \frac{(x-1)(x^2-2x)}{x^4},$$

expand the numerator and divide by x^4 :

$$y = \frac{(x-1)(x^2-2x)}{x^4} = \frac{x^3 - 3x^2 + 2x}{x^4} = x^{-1} - 3x^{-2} + 2x^{-3}.$$

Then use the Sum and Power Rules:

$$\begin{aligned}\frac{dy}{dx} &= -x^{-2} - 3(-2)x^{-3} + 2(-3)x^{-4} \\ &= -\frac{1}{x^2} + \frac{6}{x^3} - \frac{6}{x^4}.\end{aligned}$$

Second- and Higher-Order Derivatives

If $y = f(x)$ is a differentiable function, then its derivative $f'(x)$ is also a function. If f' is also differentiable, then we can differentiate f' to get a new function of x denoted by f'' . So $f'' = (f')'$. The function f'' is called the **second derivative** of f because it is the derivative of the first derivative. Notationally,

$$f''(x) = \frac{d^2y}{dx^2} = \frac{d}{dx} \left(\frac{dy}{dx} \right) = \frac{dy'}{dx} = y'' = D^2(f)(x) = D_x^2 f(x).$$

The symbol D^2 means the operation of differentiation is performed twice.

If $y = x^6$, then $y' = 6x^5$ and we have

$$y'' = \frac{dy'}{dx} = \frac{d}{dx}(6x^5) = 30x^4.$$

Thus $D^2(x^6) = 30x^4$.

If y'' is differentiable, its derivative, $y''' = dy''/dx = d^3y/dx^3$ is the **third derivative** of y with respect to x . The names continue as you imagine, with

$$y^{(n)} = \frac{d}{dx}y^{(n-1)} = \frac{d^n y}{dx^n} = D^n y$$

denoting the **n th derivative** of y with respect to x for any positive integer n .

We can interpret the second derivative as the rate of change of the slope of the tangent to the graph of $y = f(x)$ at each point. You will see in the next chapter that the second derivative reveals whether the graph bends upward or downward from the tangent line as we move off the point of tangency. In the next section, we interpret both the second and third derivatives in terms of motion along a straight line.

EXAMPLE 14 Finding Higher Derivatives

The first four derivatives of $y = x^3 - 3x^2 + 2$ are

First derivative: $y' = 3x^2 - 6x$

Second derivative: $y'' = 6x - 6$

Third derivative: $y''' = 6$

Fourth derivative: $y^{(4)} = 0$.

The function has derivatives of all orders, the fifth and later derivatives all being zero.

Exercise 1.2:

Find the first and the second derivatives of the following functions:

1. $w = 3z^7 - 7z^3 + 21z^2$

2. $y = \frac{4x^3}{3} - x + 2e^x$

3. $y = 4 - 2x - x^{-3}$

4. $r = \frac{1}{3s^2} - \frac{5}{2s}$

5. Find the derivative of the function

$$y = (3 - x^2)(x^3 - x + 1)$$

- a) by using product rule
- b) by multiplying factors to produce a sum of simpler terms

Find the derivative of the function

6. $y = \frac{2x + 5}{3x - 2}$

7. $y = \frac{x^2 + 3e^x}{2e^x - x}$

8. $v = \frac{1 + x - 4\sqrt{x}}{x}$

9. $r = \frac{1}{3s^2} - \frac{5}{2s}$

Find the first and the second derivatives of the functions:

10. $w = 3z^2e^z$

11. $u = \frac{(x^2 + x)(x^2 - x + 1)}{x^4}$