```
n1 = 2;
g1 = 3;
n2 = 1;
g2 = 2;
n3 = 1;
g3 = 2;


(*Consistencies solution for the choice of pentaquark *)

γ12bar = 0.4974508021184714; (* relationship between light and anti-charm quark *)
β12 = 0.2485750240167111; (* relationship between light and charm quark *)
β22bar = 0.882615032977455; (* relationship between charm and anti-charm quark *)

(*Defining Constants*)

xx = 0.346;
(*This is the value of αs which is αs=g^2/(ħc),
used from the mesonic fit as done in meson calculations.*)
alfas = xx;
(*aa=.178;*)
 zz = 306.;(*This is the mass of light quark like up, down*)
 aa = 0.197; (*This is the ratio of light mass to the heavy mass of quarks *)
yy = 107.6;
B0 = yy; (*Bag Constant *)
a1 = 1 / zz; (*This is α=ħ/(m1 c), where m1 is the mass of light quark. here,
reduced planck's constant and c both are set to be 1.*)
P = N[Pi, 7];
alfa1 = 1.;(* Ratio of mass of light quarks, BA, 13 September *)
alfa2 = aa; (* Ratio of mass of light quark to the heavy quark,
BA, 13 September *)

η = (n1 * g1 + n3 * g3 + n2 * g2)/5;
theta = -6 η/(5 (5 η - 1)); (*Just a multiplication factor for a particular value of eta *)
Q1 = -0.5; (*Q1 is the constant in the differential equation  d^2 f1/dx^2 =
```

Q1 $\frac{f1^{\frac{3}{2}}}{\sqrt{x}}$ for the outer region*)

```
Q3 = -1.4247837854483154;
(*Q3 is the constant in the differential equation  d^2 f3/dx^2 =
```

Q3 $\frac{f3^{\frac{3}{2}}}{\sqrt{x}}$ in the middle region. This is obtained by solving two

```
  consistency conditions along with other constant γ12bar. *)
Q2 = -3.6375498477951878 ;
(*Qbar is the constant in the differential equation  d^2 f3/dx^2 =
```

Q2 $\frac{f3^{\frac{3}{2}}}{\sqrt{x}}$ in the inner region. This is obtained by solving three

```
    consistency conditions along with other constants β13 & β23. *)
g = alfas;(*Why  g = alfas in this case ?*)
(* I think there should be g = /sqrt{alfas} in order to satisfy the multiplication
  facor in the expreesion of kinetic energy i.e. 'c', BA, 13 September *)
```

$$R = \frac{3\,a}{(8 * alfas)} * \left(\frac{3\,\pi\,\eta}{2}\right)^{\frac{2}{3}};$$(*Page number 54, it is inverse in the copy though *)

$$c = \frac{12}{5\,\pi}\,\frac{4}{3}\,g^2 - \frac{4}{3}\,\frac{alfas}{a}\left(\frac{3\,\pi\,\eta}{2}\right)^{\frac{1}{3}};$$

```
(*Multiplication factor in the expression of kinetic energy *)
```

$$d = \frac{4}{3}\,g^2 - \frac{4}{3}\,\frac{alfas}{a}\left(\frac{3\,\pi\,\eta}{2}\right)^{\frac{1}{3}}\,\frac{18}{25\,\pi}\,\frac{\eta}{\left(\eta - \frac{1}{5}\right)};$$ (* $\frac{18g^2}{5(5\eta\,-1\,)}$.16 π^2.$\left(\frac{3a}{8alfas}\right)$. $\frac{R^2}{(6\pi^2)^2}$,

```
The value of R is from page number 54 which is written inverse here,
and (3a/8alfas) is inverse in copy, BA, 13 September *)
(*Multiplication factor in the expression of Potential energy *)


(*Let us define the three functions for three regions *)

f3[x3_?NumericQ, x2_?NumericQ, u2_?NumericQ , s2_?NumericQ] :=
  Abs[x3 * y3'[x3] - y3[x3] - n3 * Q3/(3 η) - β22bar * n2 * Q2/(3 η) + β22bar^(3/2) * n2 * Q3/(3 η)]
```

```
(*Appendix from Kinetic energy*) /.
 First@NDSolve[{y3''[x] == (Q3 * Abs[y3[x]^(3/2)] / x^(1/2)), y3'[x2] == β22bar * s2 ,
    y3[x2] == β22bar * u2 (* It shows how the function f3
     (x3) changes on other regions*)}, y3, {x, x2, x3}];
(* s2 is the first derivative of function f_2(x) *)
(*u2 is the function f_2(x) *)
(*This is the function f3 defined on the
  middle region. It has the differential equation  d^2 f3/dx^2 =
 Q3 f3^(3/2)/Sqrt[x] in for x3<x<x2 which is middle region. On this region, f1(x)=γ12 f2(x).  *)
f1[x1_?NumericQ, x3_?NumericQ, s3_?NumericQ, u3_?NumericQ] :=
  Abs[x1 * y1'[x1] - y1[x1] - n1 * Q1/(3 η) - γ12bar * n3 * Q3/(3 η) +
    γ12bar^(3/2) * n3 * Q1/(3 η) - (γ12bar^(3/2) * β22bar^(3/2) - β12^(3/2)) * n2 * Q1/(3 η) +
    γ12bar * β22bar^(3/2) * n2 * Q3/(3 η) - γ12bar * β22bar * n2 * Q2/(3 η)]
  (*Appendix from Kinetic energy*) /. First@NDSolve[{y1''[x] == (Q1 * Abs[y1[x]^(3/2)] / x^(1/2)),
    y1'[x3] == γ12bar * s3, (* s3 is the first derivative of function f_2bar(x) *)
    (* It means at the boundary x3 the function df1 changes to other function *)
    y1[x3] == γ12bar * u3 (* u3 is the function f_2bar(x) *)
    (*It means at the boundary x3 the function f1 changes to other function*)
   }, (* It means at the boundary x3 the function f1 changes to other function *)
   y1, {x, x3, x1}]; (*This is the function defined in outer region for x2<
  x<x1 so called f1. It has differential equation  d^2 f1/dx^2 =
 Q1 f1^(3/2)/Sqrt[x] for the outer region which is solved by using boundary
   condition of (d/dx) f1(x)|x2 = γ12bar (d/dx) f2(x)|x2 and f1(x3)=γ12 f2(x3)  *)
```

```
f2[xmax_?NumericQ, m2_?NumericQ] :=
  Module[{xx = xmax},
   solution2 = NDSolve[
     {y2''[x] == (Q2 * Abs[y2[x]^(3/2)]) / x^(1/2), y2'[1.*^-9] == m2, y2[1.*^-9] == 0.}, y2, {x, 1.*^-9, xx}];
   m = NIntegrate[Sqrt[x] * Evaluate[y2[x] /. solution2]^(3/2),
     {x, 1.*^-9, xx}, Method → {Automatic, "SymbolicProcessing" → 0}];
   z = m[[1]];
   z = z - n2 / (3. * η);
   (*z=z-n/(A)*);
   Return[Abs[z]]; Remove["NDSolve`*$*$1"]];
(*This is the function f3 defined on the inner region. It has the differential
  equation  d^2 f3/dx^2 = Q3 f3^(3/2)/Sqrt[x]  for 0<x<x_3 inner region. On this region,
f_1(x)=β_13 f_3(x) and f_2(x)= β_23 f_3(x).    *)

(*Start the program*)

m2 = 14;
For[j = 1., j ≤ 1., j++, B = (B0)^4;
  a = a1; (* It must be to satisfy the multiplication
   factor used in kinetic energy i.e. c, BA, 13 September *)
  (*Now for the first pentaquark,
  eta=1 which in total contains 5 quarks which can be written by *)

 mqn = If[j > 1, m2 + 2, m2]; (*case2new*)(*mqn=If[j>1,m2+4,m2];*)
  m2 = mqn;
  topi = 0.01;
  m2B = m2;
  Energy = 10^9;
  Energy = 10^9;
  Energy2plus = 10^9;
```

```
H = 10^9 + 1;
SN = 0;
nloop = 1;
For[n = 1, n ≤ 3, n++, While[Energy ≤ H, F = H;
  H = Energy;
  HX = Energy6bare;
  HY = Energy2plus; m2 = m2 + SN * 10^(-n);
  x2 = 2.*^-8; (*Print[m2];*)(*was 0 .5 1.66*)
  FindMinimum[f2[x2, m2], {x2, topi}];
  (*remove this function and investigate if it is actually required BS 4/3/2021*)
  x2 = x2 /. Last[FindMinimum[f2[x2, m2], {x2, topi}]];
  solution2 = NDSolve[{y2''[x] == (Q2 * Abs[y2[x]^(3/2)]) / x^(1/2),
     y2'[1.*^-9] == m2, y2[1.*^-9] == 0.}, y2, {x, 1.*^-9, x2}];
  t1 = Evaluate[y2[x2] /. solution2];
  t4 = Evaluate[y2'[x2] /. solution2];
  u2 = t1[[1]]; (* u2 is the function f_2(x) *)
  s2 = t4[[1]]; (* s2 is the first derivative of function f_2(x) *)
  If[u2 < 0., u2 = 0.];


  x3 = x2 + 0.1;
  FindMinimum[{f3[x3, x2, u2, s2]}, {x3, x2}];
  x3 = x3 /. Last[FindMinimum[{f3[x3, x2, u2, s2]}, {x3, x2}]];
  solution3 = NDSolve[{y3''[x] == (Q3 * Abs[y3[x]^(3/2)]) / x^(1/2),
     y3'[x2] == β22bar * s2 , y3[x2] == β22bar * u2}, y3, {x, x2, x3}];
  t2 = Evaluate[y3[x3] /. solution3];
  t5 = Evaluate[y3'[x3] /. solution3];
  u3 = t2[[1]]; (* u3 is the function f_2bar(x) *)
  s3 = t5[[1]]; (* s3 is the first derivative of function f_2bar(x) *)
  If[u3 < 0., u3 = 0.];

  x1 = x3 + 0.1;
  FindMinimum[{f1[x1, x3, s3, u3]}, {x1, x3}];
  x1 = x1 /. Last[FindMinimum[{f1[x1, x3, s3, u3]}, {x1, x3}]];
```

```
solution1 = NDSolve[{y1''[x] == (Q1 * Abs[y1[x]^(3/2)] / x^(1/2)),
    y1'[x3] == γ12bar * s3 , y1[x3] == γ12bar * u3 }, y1, {x, x3, x1}];
t3 = Evaluate[y1[x1] /. solution1];
t6 = Evaluate[y1'[x1] /. solution1];
u1 = t3[[1]]; (* u1 is the function f_1(x) *)
s1 = t6[[1]]; (* s1 is the first derivative of function f_1(x) *)
If[u1 < 0., u1 = 0.];


(*Integrations *)(* These integrations change values but outer and outer 1
 doesnot because they depend on x and derivates of x (BA, 11 September)*)

normmx2 = x2 s2 - u2 ;
normx2 = n2 Q2 / (3 η) ;
(*normx2barr and normx2bar should be equal *)
normmx3 = x3 s3 - u3 ;
normx3 = n3 Q3 / (3 η) + β22bar n2 Q2 / (3 η) - (β22bar)^(3/2) n2 Q3 / (3 η);
(*normmx2 and normx2 should be equal *)
normmx1 = x1 s1 - u1 ;
normx1 = n1 Q1 / (3 η) + γ12bar n3 Q3 / (3 η) - (γ12bar)^(3/2) n3 Q1 / (3 η) +
  ((γ12bar)^(3/2) (β22bar)^(3/2) - (β12)^(3/2)) n2 Q1 / (3 η) - γ12bar (β22bar)^(3/2) n2 Q3 / (3 η) + γ12bar β22bar n2 Q2 / (3 η) ;
(*normmx1 and normx1 should be equal *)

h1 = NIntegrate[Evaluate[y2[x] /. solution2]^2.5 / (x^0.5),
  {x, 0, x2}, Method → {Automatic, "SymbolicProcessing" → 0}];
h2 = NIntegrate[Evaluate[y3[x] /. solution3]^2.5 / (x^0.5), {x, x2, x3},
  Method → {Automatic, "SymbolicProcessing" → 0}];
h3 = NIntegrate[Evaluate[y1[x] /. solution1]^2.5 / (x^0.5), {x, x3, x1},
  Method → {Automatic, "SymbolicProcessing" → 0}];
h4 = h1[[1]];
h5 = h2[[1]];
```

```
h6 = h3[[1]] ;

AB = 4/7 Sqrt[x2] (u2)^(5/2) - 5 n2/(21 η) s2 ;

CD = 4/7 Sqrt[x3] (u3)^(5/2) - 4/7 (β22bar)^(5/2) Sqrt[x2] (u2)^(5/2) -
   5/(7 Q3) s3 (n3 Q3/(3 η) + β22bar n2 Q2/(3 η) - (β22bar)^(3/2) n2 Q3/(3 η)) + 5/(7 Q3) (β22bar)^2 n2 Q2/(3 η) s2 ;

CCD = 4/7 Sqrt[x3] (u3)^(5/2) - 4/7 Sqrt[x2] (β22bar)^(5/2) (u2)^(5/2) - 5/(7 Q3) s3 normmx3 +
   5/(7 Q3) (β22bar)^2 normmx2 s2 ;

EF = 4/7 Sqrt[x1] (u1)^(5/2) - 4/7 (γ12bar)^(5/2) Sqrt[x3] (u3)^(5/2) -
   5/(7 Q1) s1 (n1 Q1/(3 η) + γ12bar n3 Q3/(3 η) - (γ12bar)^(3/2) n3 Q1/(3 η) + n2 Q1/(3 η) ((γ12bar)^(3/2) (β22bar)^(3/2) - (β12)^(3/2)) -
      (γ12bar) (β22bar)^(3/2) n2 Q3/(3 η) + (γ12bar) (β22bar) n2 Q2/(3 η)) +
   5/(7 Q1) (γ12bar)^2 s3 (n3 Q3/(3 η) + β22bar n2 Q2/(3 η) - (β22bar)^(3/2) n2 Q3/(3 η)) ;
(* h4 and AB should be equal *)
(* h5 and CD should be equal *)
(* h6 and EF should be equal *)


K222 = n2/(3 η Q2) s2 - 1/Q2 h4 ;

K22 = 4 n2/(7 η Q2) s2 - 4/(7 Q2) Sqrt[x2] (u2)^(5/2) ;

(*K2bar2barr and K2bar2bar should be equal *)

K32 = - 4/(7 Q2) β22bar^1.5 Sqrt[x2] (u2)^(5/2) +
```

$$\frac{n2}{3\,\eta}\left(\frac{\beta 22bar^{1.5}}{Q2}-\frac{\beta 22bar}{Q3}+\frac{5*\beta 22bar^{1.5}}{7\,Q2}\right)s2+\frac{n2}{3\,\eta\,Q3}s3\,;$$

$$K332=\frac{n2\,(\beta 22bar)^{\frac{3}{2}}}{3\,\eta\,Q2}s2-\frac{n2\,\beta 22bar}{3\,\eta\,Q3}s2+\frac{n2}{3\,\eta\,Q3}s3-\frac{(\beta 22bar)^{\frac{3}{2}}}{Q2}h4;$$

(*K2bar2 and K2barr2 should be equal *)

$$K21=-\frac{4}{7\,Q2}\beta 12^{1.5}\sqrt{x2}\;(u2)^{\frac{5}{2}}+\frac{n2}{3\,\eta}\left(\frac{5}{7\,Q2}\beta 12^{1.5}+\frac{\beta 12^{1.5}}{Q2}-\frac{\gamma 12bar^{1.5}*\beta 22bar}{Q3}\right)s2+$$
$$\frac{n2}{3\,\eta}\left(\frac{\gamma 12bar^{1.5}}{Q3}-\frac{\gamma 12bar}{Q1}\right)s3+\frac{n2}{3\,\eta\,Q1}s1\,;$$

$$K221=\frac{n2}{3\,\eta\,Q2}(\beta 12)^{\frac{3}{2}}s2+\frac{n2\,(\gamma 12bar)^{\frac{3}{2}}}{3\,\eta\,Q3}s3-$$
$$\frac{n2}{3\,\eta\,Q3}(\gamma 12bar)^{\frac{3}{2}}\beta 22bar\,s2+\frac{n2}{3\,\eta\,Q1}s1-\frac{n2}{3\,\eta\,Q1}\gamma 12bar\,s3-\frac{(\beta 12)^{\frac{3}{2}}}{Q2}h4\,;$$

(*K2bar1 and K2barr1 should be equal *)

$$K31=\frac{4}{7}\beta 22bar^{1.5}\left(\frac{\gamma 12bar^{1.5}*\beta 22bar}{Q3}-\frac{\beta 12^{1.5}}{Q2}\right)\sqrt{x2}\;(u2)^{\frac{5}{2}}-\frac{4}{7\,Q3}\gamma 12bar^{1.5}\sqrt{x3}\;(u3)^{\frac{5}{2}}+$$
$$\frac{n2}{3\,\eta}\beta 22bar\left(\frac{5}{7\,Q2}\beta 22bar^{\frac{1}{2}}*\beta 12^{1.5}-\frac{\gamma 12bar^{1.5}*\beta 22bar^{1.5}}{Q3}-\frac{5\,Q2}{7\,Q3^2}\gamma 12bar^{1.5}*\right.$$
$$\left.\beta 22bar+\frac{Q2}{Q3^2}\gamma 12bar^{1.5}*\beta 22bar+\frac{1}{Q2}\beta 12^{1.5}*\beta 22bar^{\frac{1}{2}}-\frac{\beta 12^{1.5}}{Q3}\right)s2+$$
$$\frac{1}{3\,\eta\,Q3}\left(n3*\gamma 12bar^{1.5}-\frac{n3\,Q3}{Q1}\gamma 12bar+n2*\beta 12^{1.5}-\frac{n2\,Q2}{Q3}\gamma 12bar^{1.5}*\beta 22bar+\right.$$
$$\left.\frac{5}{7}\gamma 12bar^{1.5}\left(n3+\frac{n2\,Q2}{Q3}\beta 22bar-n2*\beta 22bar^{1.5}\right)\right)s3+\frac{n3}{3\,\eta\,Q1}s1\,;$$

$$(*K122 = \frac{n2}{3\,\eta}(\gamma 12bar)^{\frac{3}{2}}(\beta 22bar)^{\frac{3}{2}}\left[\frac{1}{Q3}s3 - \frac{\beta 22bar}{Q3}s2\right] - \frac{(\beta 12)^{\frac{3}{2}}(\beta 22bar)^{\frac{3}{2}}}{Q2}h4 - \frac{(\gamma 12bar)^{\frac{3}{2}}}{Q3}h5$$

$$- \frac{n2\,Q2}{3\,\eta\,Q3}(\gamma 12bar)^{\frac{3}{2}}\beta 22bar\left[\frac{1}{Q3}s3 - \frac{\beta 22bar}{Q3}s2\right] + \frac{n3}{3\,\eta\,Q1}s1 - \frac{n3}{3\,\eta\,Q1}\gamma 12bar\,s3$$

$$+ (\beta 12)^{\frac{3}{2}}\frac{n2}{3\,\eta}\left[\frac{(\beta 22bar)^{\frac{3}{2}}}{Q2}s2 + \frac{1}{Q3}s3 - \frac{\beta 22bar}{Q3}s2\right] +$$

$$\frac{n3}{3\,\eta\,Q3}(\gamma 12bar)^{\frac{3}{2}}s3 - \frac{n2}{3\,\eta\,Q3}(\gamma 12bar)^{\frac{3}{2}}(\beta 22bar)^{\frac{3}{2}}s3\ ;\ *)$$

$$K331 = \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}s3 - \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}\beta 22bar\,s2 - \frac{(\beta 12)^{\frac{3}{2}}(\beta 22bar)^{\frac{3}{2}}}{Q2}h4 - \frac{(\gamma 12bar)^{\frac{3}{2}}}{Q3}h5 -$$

$$\frac{n2\,Q2}{3\,\eta\,Q3^2}(\gamma 12bar)^{\frac{3}{2}}\beta 22bar\,s3 + \frac{n2\,Q2}{3\,\eta\,Q3^2}(\gamma 12bar)^{\frac{3}{2}}(\beta 22bar)^2\,s2 + \frac{n2}{3\,\eta}(\beta 22bar)^{\frac{3}{2}}$$

$$\left(\frac{\beta 12^{\frac{3}{2}}}{Q2} - \frac{(\gamma 12bar)^{\frac{3}{2}}\beta 22bar}{Q3}\right)s2 + \frac{n3}{3\,\eta}\gamma 12bar\left(\frac{\gamma 12bar^{\frac{1}{2}}}{Q3} - \frac{1}{Q1}\right)s3 + \frac{n3}{3\,\eta\,Q1}s1\ ;$$

(*K12 and K122 should be equal*)

$$K33 = \frac{4}{7}\beta 22bar^2\left(\frac{\beta 22bar^{\frac{1}{2}}}{Q3} - \frac{\beta 22bar}{Q2}\right)\sqrt{x2}\,(u2)^{\frac{5}{2}} -$$

$$\frac{4}{7\,Q3}\sqrt{x3}\,(u3)^{\frac{5}{2}} + \frac{n2}{3\,\eta}\beta 22bar^2\left(\beta 22bar\frac{12}{7\,Q2} + \frac{2}{7}\frac{Q2}{Q3^2} - \frac{2*\beta 22bar^{\frac{1}{2}}}{Q3}\right)s2 +$$

$$\left(\frac{12\,n3}{21\,\eta\,Q3} + \beta 22bar\frac{n2}{3\,\eta\,Q3}\left(\beta 22bar^{\frac{1}{2}} - \frac{2}{7}\frac{Q2}{Q3} - \frac{5}{7}\beta 22bar^{\frac{1}{2}}\right)\right)s3\ ;$$

$$K333 = \frac{n2\,Q2}{3\,\eta\,Q3}\left(\frac{(\beta 22bar)^{\frac{3}{2}}}{Q2} - \frac{\beta 22bar}{Q3}\right)s3 - \frac{n2\,Q2\,\beta 22bar}{3\,\eta\,Q3}\left(\frac{(\beta 22bar)^{\frac{3}{2}}}{Q2} - \frac{\beta 22bar}{Q3}\right)s2 -$$

$$\frac{(\beta 22bar)^3}{Q2}h4 - \frac{1}{Q3}h5 + (\beta 22bar)^3\frac{n2}{3\,\eta\,Q2}s2 - (\beta 22bar)^{\frac{5}{2}}\frac{n2}{3\,\eta\,Q3}s2 + \frac{n3}{3\,\eta\,Q3}s3\ ;$$

(*K22 and K222 should be equal *)

(*THese Y and Z are to check the value of integral of power $\frac{5}{2}$ in the outer region. Which proved to be false yet. I need to work on it extensively.*)

$$K11 = \frac{4}{7}\left(\frac{\beta22bar^{\frac{5}{2}} * \gamma12bar^3}{Q3} - \frac{\beta12^3}{Q2}\right)\sqrt{x2}\ (u2)^{\frac{5}{2}} + \frac{4}{7}\left(\frac{\gamma12bar^{\frac{5}{2}}}{Q1} - \frac{\gamma12bar^3}{Q3}\right)\sqrt{x3}\ (u3)^{\frac{5}{2}} - \frac{4}{7\,Q1}\sqrt{x1}\ (u1)^{\frac{5}{2}} + \left(\frac{n2}{3\,\eta}\left(\frac{\beta12^3}{Q2} - \frac{\beta12^{\frac{3}{2}} * \gamma12bar^{\frac{3}{2}} * \beta22bar}{Q3} + \frac{Q2}{Q3^2}\gamma12bar^3 * \beta22bar^2 - \frac{\beta12^{1.5} * \gamma12bar^{1.5} * \beta22bar}{Q3}\right) + \frac{5\,n2}{21\,\eta}\left(\frac{\beta12^3}{Q2} - \frac{Q2}{Q3^2}\beta22bar^2 * \gamma12bar^3\right)\right)s2 + \left(\frac{n2}{3\,\eta}\left(\frac{\beta12^{1.5} * \gamma12bar^{1.5}}{Q3} - \frac{\beta12^{1.5} * \gamma12bar}{Q1} - \frac{\beta22bar^{1.5} * \gamma12bar^3}{Q3} + \frac{\beta22bar^{1.5} * \gamma12bar^{\frac{5}{2}}}{Q1} + \frac{\beta12^{1.5} * \gamma12bar^{1.5}}{Q3} - \frac{Q2}{Q3^2}\gamma12bar^3 * \beta22bar - \frac{\beta12^{1.5} * \gamma12bar}{Q1} + \frac{\gamma12bar^{\frac{5}{2}} * \beta22bar^{1.5}}{Q1} + \frac{Q2}{Q1^2}\gamma12bar^2 * \beta22bar - \frac{Q3}{Q1^2}\beta22bar^{1.5} * \gamma12bar^2\right) + \frac{n3}{3\,\eta}\left(\frac{\gamma12bar^3}{Q3} - \frac{2 * \gamma12bar^{\frac{5}{2}}}{Q1} + \frac{Q3}{Q1^2}\gamma12bar^2\right) + \frac{5}{21\,\eta}\left(\frac{n3}{Q3}\gamma12bar^3 + \frac{n2\ Q2}{Q3^2}\gamma12bar^3 * \beta22bar - \frac{n2}{Q3}\beta22bar^{1.5} * \gamma12bar^3 - \frac{n3\ Q3}{Q1^2}\gamma12bar^2 - \frac{n2\ Q2}{Q1^2}\gamma12bar^2 * \beta22bar + \frac{n2\ Q3}{Q1^2}\gamma12bar^2 * \beta22bar^{1.5}\right)\right) * s3 + \left(\frac{n1}{3\,\eta\,Q1} - \frac{n3\,Q3}{3\,\eta\,Q1^2}\gamma12bar + \frac{n3}{3\,\eta\,Q1}\gamma12bar^{1.5} + \frac{n2}{3\,\eta\ Q1}\right.$$

$$\left(\beta 12^{1.5} - \gamma 12bar^{1.5} * \beta 22bar^{1.5} - \frac{Q2}{Q1}\beta 22bar * \gamma 12bar + \frac{Q3}{Q1}\beta 22bar^{1.5} * \gamma 12bar\right) + \frac{5}{21\,\eta}\left(\frac{n1}{Q1} + \frac{n3\ Q3}{Q1^2}\gamma 12bar - \frac{n3}{Q1}\gamma 12bar^{1.5} + \frac{n2}{Q1}\gamma 12bar^{1.5} * \beta 22bar^{1.5} - \frac{n2}{Q1}\beta 12^{1.5} - \frac{n2\ Q3}{Q1^2}\beta 22bar^{1.5} * \gamma 12bar + \frac{n2\ Q2}{Q1^2}\gamma 12bar\ \beta 22bar\right)\Bigg) * s1;$$

$$K111 = \frac{-(\beta 12)^3}{Q2}h4 - \frac{(\gamma 12bar)^3}{Q3}h5 - \frac{1}{Q1}h6 + \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}(\gamma 12bar)^{\frac{3}{2}}s3 - \frac{n2\,Q2}{3\,\eta\,Q3^2}(\gamma 12bar)^3\,\beta 22bar\,s3 - \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}(\gamma 12bar)^{\frac{3}{2}}\beta 22bar\,s2 + \frac{n2\,Q2}{3\,\eta\,Q3^2}(\gamma 12bar)^3(\beta 22bar)^2 s2 + \left(\frac{n2}{3\,\eta\,Q1}(\beta 12)^{\frac{3}{2}} + \frac{n3}{3\,\eta\,Q1}(\gamma 12bar)^{\frac{3}{2}} - \frac{n3\,Q3}{3\,\eta\,Q1^2}\gamma 12bar - \frac{n2}{3\,\eta\,Q1}(\gamma 12bar)^{\frac{3}{2}}(\beta 22bar)^{\frac{3}{2}} - \frac{n2\,Q2}{3\,\eta\,Q1^2}\beta 22bar\,\gamma 12bar + \frac{n2\,Q3}{3\,\eta\,Q1^2}(\beta 22bar)^{\frac{3}{2}}\gamma 12bar\right)s1 - \left(\frac{n2}{3\,\eta\,Q1}(\beta 12)^{\frac{3}{2}}\gamma 12bar + \frac{n3}{3\,\eta\,Q1}(\gamma 12bar)^{\frac{5}{2}} - \frac{n3\,Q3}{3\,\eta\,Q1^2}(\gamma 12bar)^2 - \frac{n2}{3\,\eta\,Q1}(\gamma 12bar)^{\frac{5}{2}}(\beta 22bar)^{\frac{3}{2}} - \frac{n2\,Q2}{3\,\eta\,Q1^2}(\gamma 12bar)^2\,\beta 22bar + \frac{n2\,Q3}{3\,\eta\,Q1^2}(\beta 22bar)^{\frac{3}{2}}(\gamma 12bar)^2\right)s3 + \frac{n2}{3\,\eta}\frac{(\beta 12)^3}{Q2}s2 - \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}(\gamma 12bar)^{\frac{3}{2}}\beta 22bar\,s2 + \frac{n2}{3\,\eta\,Q3}(\beta 12)^{\frac{3}{2}}(\gamma 12bar)^{\frac{3}{2}}s3 - \frac{n2}{3\,\eta\,Q1}(\beta 12)^{\frac{3}{2}}\gamma 12bar\,s3 + \frac{n3}{3\,\eta\,Q3}(\gamma 12bar)^3 s3 - \frac{n2}{3\,\eta\,Q3}(\beta 22bar)^{\frac{3}{2}}(\gamma 12bar)^3 s3 - \frac{n3}{3\,\eta\,Q1}(\gamma 12bar)^{\frac{5}{2}}s3 + \frac{n2}{3\,\eta\,Q1}(\beta 22bar)^{\frac{3}{2}}(\gamma 12bar)^{\frac{5}{2}}s3 + \frac{n1}{3\,\eta\,Q1}s1;$$

(*K11 and K111 should be equal *)

$$T1 = c\left(\frac{4}{7}\left(alfa1 * g1\left(\beta 12^{\frac{5}{2}} - \gamma 12bar^{\frac{5}{2}} * \beta 22bar^{\frac{5}{2}}\right) + alfa2 * g2\right)\sqrt{x2}\ (u2)^{\frac{5}{2}} +$$

$$\frac{4}{7}\,\text{alfa2}*\text{g3}\sqrt{\text{x3}}\,(\text{u3})^{\frac{5}{2}}+\frac{4}{7}\,\text{alfa1}*\text{g1}\sqrt{\text{x1}}\,(\text{u1})^{\frac{5}{2}}+$$

$$\frac{5\,\text{n2}}{21\,\eta}\left(\beta 22\text{bar}^{2}\,\frac{\text{Q2}}{\text{Q3}}\left(\text{alfa1}*\text{g1}*\gamma 12\text{bar}^{\frac{5}{2}}+\text{alfa2}*\text{g3}\right)-\right.$$

$$\left.\text{alfa1}*\text{g1}*\beta 12^{\frac{5}{2}}-\text{alfa2}*\text{g3}*\beta 22\text{bar}^{\frac{5}{2}}-\text{alfa2}*\text{g2}\right)\text{s2}+$$

$$\frac{5}{21\,\eta}\left(\frac{\gamma 12\text{bar}^{2}*\text{alfa1}*\text{g1}}{\text{Q1}}\left(\text{n3}*\text{Q3}+\beta 22\text{bar}*\text{n2}*\text{Q2}-\beta 22\text{bar}^{1.5}*\text{n2}*\text{Q3}\right)-\right.$$

$$\left.\left(\text{alfa1}*\text{g1}*\gamma 12\text{bar}^{\frac{5}{2}}+\text{alfa2}*\text{g3}\right)\left(\text{n3}+\beta 22\text{bar}\,\frac{\text{n2}*\text{Q2}}{\text{Q3}}-\beta 22\text{bar}^{1.5}*\text{n2}\right)\right)\text{s3}-$$

$$\frac{5\,\text{alfa1}\,\text{g1}}{21\,\eta}\left(\text{n1}+\gamma 12\text{bar}\,\frac{\text{n3}\,\text{Q3}}{\text{Q1}}+\gamma 12\text{bar}^{1.5}\left(\beta 22\text{bar}^{1.5}*\text{n2}-\text{n3}\right)-\right.$$

$$\left.\beta 12^{1.5}*\text{n2}-\gamma 12\text{bar}*\beta 22\text{bar}^{1.5}\,\frac{\text{n2}\,\text{Q3}}{\text{Q1}}+\gamma 12\text{bar}\,\beta 22\text{bar}\,\frac{\text{n2}\,\text{Q2}}{\text{Q1}}\right)\text{s1}\Bigg);$$

$$\text{V1}=-\text{d}\left(\frac{(\text{g1}\,\text{n1}-1)}{\text{n1}}\,\text{g1}\,\text{K11}+\frac{(\text{g3}\,\text{n3}-1)}{\text{n3}}\,\text{g3}\,\text{K33}+\right.$$

$$\left.\frac{(\text{g2}\,\text{n2}-1)}{\text{n2}}\,\text{g2}\,\text{K22}+2\,\text{g1}\,\text{g3}\,\text{K31}+\frac{4}{3}\,\text{g2}\,\text{g1}\,\text{K21}+\frac{4}{3}\,\text{g2}\,\text{g3}\,\text{K32}\right);$$

```
(*These T1 and V1 are obtained
 by integrating manually which is being verified.*)
Energy6bare = T1 + V1 + 4 * P / 3 * R ^ 3 * x1 ^ 3 * B + n1 * g1 * 306 + n3 * g3 * 1553 + n2 * g2 * 1553;
Energy = Energy6bare;
Volumeenergy = 4 * P / 3 * R ^ 3 * x1 ^ 3 * B;
Energywithoutmass = T1 + V1 + 4 * P / 3 * R ^ 3 * x1 ^ 3 * B;
Print["R=  ", R];
Print["normx2 (computationally)=  ", normmx2];
Print["normx2 (by hand)=  ", normx2];
Print["normx3 (computationally)=  ", normmx3];
Print["normx3 (by hand)=  ", normx3];
Print["normx1 (computationally)=  ", normmx1];
Print["normx1 (by hand)=  ", normx1];
Print["0 to x2 (h4 computationally)=  ", h4];
Print["0 to x2 (h4 by hand)=  ", AB];
```

```
 Print["x2 to x3 (h5 computationally)=  ", h5];
 Print["x2 to x3 (h5 by hand)=  ", CD];
 Print["x2 to x3 (h5 by handdddd)=  ", CCD];
 Print["x3 to x1 (h6 computationally)=  ", h6];
 Print["x3 to x1 (h6 by hand)=  ", EF];
 Print["K22  (computationally) = ", K222];
 Print["K22  (by hand) = ", K22];
 Print["K32  (computationally) = ", K332];
 Print["K32  (by hand) = ", K32];
 Print["K21  (computationally) = ", K221];
 Print["K21  (by hand) =", K21];
 Print["K31  (computationally) = ", K331];
 Print["K31  (by hand) = ", K31];
 Print["K33  (computationally) = ", K333];
 Print["K33  (by hand) = ", K33];
 Print["K11  (computationally) = ", K111];
 Print["K11  (by hand) = ", K11];
 Print["x2= ", x2];
 Print["x3= ", x3];
 Print["x1=  ", x1];
  Print["T by hand=", T1];
 Print["V by hand=", V1];
 Print["m2=", m2];
 Print["Energy without mass = ", Energywithoutmass];
 Print["Energy=  ", Energy];
 If[n == 1, If[nloop == 1, nloop = nloop + 1; SN = 1. * 10^(-1);
   Energy11 = Energy, If[nloop == 2, nloop = nloop + 1;
    H = 10^9 + 1;
    m2 = m2B;
    If[Energy - Energy11 < 0, SN = 1., SN = -1.]]]];
If[n < 3, m2 = m2 - SN * 10^(-n); Energy = F; H = F];
If[n == 3, Energy6bare = HX;
 Print["x1=", x1];
 Print["x3=", x3];
 Print["x2=", x2];
 Print["Kinetic Energy=", T1];
 Print["Potential Energy=", V1];
 Print["Volume Energy=", Volumeenergy];
]]];

(*Code for Plotting function *)
Plot1 = Plot[Evaluate[((y2[x] /.solution2)/ x)^(3 / 2)],
```

```
    {x, .00001, x2}, (*%PlotRange→{{0,0.65},{0,79.5}},*)
    (*Solution 1 = f3(x3), solution3 = f3(x3), solution3 = f1(x1) *)
    PlotRange → All,
    AxesLabel → {StyleForm["x", Italic, FontSize → 17]
       , StyleForm["(f2(x)/x)^(3/2)", Italic, FontSize → 17]},
    PlotStyle -> {Thickness[0.01], Green}, AxesStyle → Black,
    PlotLabel → None, LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large];
Plot2 = Plot[Evaluate[((β22bar * y2[x] /.solution2) / x)^(3 / 2)], {x, .00001, x2},
  (*%PlotRange→{{0,.65},{0,79.5}},*)
  PlotRange → All, AxesLabel → {StyleForm["x", Italic, FontSize → 17],
    StyleForm["(f2 bar(x)/x)^(3/2)", Italic, FontSize → 17]},
  PlotStyle -> {Thickness[0.01], Orange}, AxesStyle → Black, PlotLabel → None,
  LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large]
Plot3 = Plot[Evaluate[((y3[x] /.solution3) / x)^(3 / 2)],
  {x, x2, x3}, (*PlotRange→{{0,.65},{0,79.5}},*)
  PlotRange → All,
  AxesLabel → {StyleForm["x", Italic, FontSize → 17],
    StyleForm["(f2 bar(x)/x)^(3/2)", Italic, FontSize → 17]},
  PlotStyle -> {Thickness[0.01], Orange}, AxesStyle → Black, PlotLabel → None,
  LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large]
Plot4 = Plot[Evaluate[((β12 * y2[x] /.solution2) / x)^(3 / 2)], {x, .00001, x2},
  (*%PlotRange→{{0,.65},{0,79.5}},*)
  PlotRange → All, AxesLabel →
   {StyleForm["x", Italic, FontSize → 17], StyleForm["(f1(x)/x)^(3/2)", Italic, FontSize → 17]},
  PlotStyle -> {Thickness[0.01], Blue}, AxesStyle → Black, PlotLabel → None,
  LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large]
Plot5 = Plot[Evaluate[((γ12bar * y3[x] /. solution3) / x)^(3 / 2)],
  {x, x2, x3}, (*PlotRange→{{0,.65},{0,79.5}},*)
  PlotRange → All,
  AxesLabel →
   {StyleForm["x", Italic, FontSize → 17], StyleForm["(f1(x)/x)^(3/2)", Italic, FontSize → 17]},
  PlotStyle -> {Thickness[0.01], Blue}, AxesStyle → Black, PlotLabel → None,
  LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large]
Plot6 = Plot[Evaluate[((y1[x] /. solution1) / x)^(3 / 2)],
  {x, x3, x1}, (*PlotRange→{{0,.65},{0,79.5}},*)
  PlotRange → All,
  AxesLabel →
   {StyleForm["x", Italic, FontSize → 17], StyleForm["(f1(x)/x)^(3/2)", Italic, FontSize → 17]},
  PlotStyle -> {Thickness[0.01], Blue}, AxesStyle → Black, PlotLabel → None,
  LabelStyle → {GrayLevel[0], Bold}, ImageSize → Large]
Show[{Plot1}]
```

```
Show[{Plot2, Plot3}]
Show[{Plot4, Plot5, Plot6}]
Show[Plot1, Plot2, Plot3, Plot4, Plot5, Plot6,
 AxesLabel → {StyleForm["x", Italic, FontSize → 17],
   StyleForm["(f(x)/ x\!\(\*SuperscriptBox[\(\), \(3/ 2\)]\)", Italic, FontSize → 17]}]
```