

Predictive Pattern Recognition Techniques Towards Spatiotemporal Representation of Plant Growth in Simulated and Controlled Environments: A Comprehensive Review

Mohamed Debbagh¹, Shangpeng Sun¹, and Mark Lefsrud¹

¹Department of Bioresource Engineering, McGill University, Montréal, Canada.

Abstract

Accurate predictions and representations of plant growth patterns in simulated and controlled environments are important for addressing various challenges in plant phenomics research. This review explores various works on state-of-the-art predictive pattern recognition techniques, focusing on the spatiotemporal modeling of plant traits and the integration of dynamic environmental interactions. We provide a comprehensive examination of deterministic, probabilistic, and generative modeling approaches, emphasizing their applications in high-throughput phenotyping and simulation-based plant growth forecasting. Key topics include regressions and neural network-based representation models for the task of forecasting, limitations of existing experiment-based deterministic approaches, and the need for dynamic frameworks that incorporate uncertainty and evolving environmental feedback. This review surveys advances in 2D and 3D structured data representations through functional-structural plant models and conditional generative models. We offer a perspective on opportunities for future works, emphasizing the integration of domain-specific knowledge to data-driven methods, improvements to available datasets, and the implementation of these techniques toward real-world applications.

1 Introduction

With the growing prevalence of challenges in global food systems, including restricted access to fresh produce for remote populations and issues faced within traditional cultivation practices, outlooks on controlled environment agriculture (CEA) present potential solutions through precision methods of localized food production [1–3]. Controlled environments within crop production systems provide opportunities to observe and cultivate plants with diverse characteristics, or phenotypes, making it suitable for specialized applications from trait selection for cultivar development and biopharming [4–8], to food production in extreme and resource scarce environments [9–11]. Moreover, the ability to accurately predict and represent the phenotypic expression of a plant and its trajectory at various points in its growth cycle demonstrates a fundamental step towards advancing simulation and modeling research within CEA.

Recent developments in computational and algorithmic theories, coupled with innovations in data collection and fusion techniques, have opened new avenues for exploring plant phenotypes and growth patterns within simulated and controlled environments [12]. These environments allow for the precise manipulation of variables such as light, temperature, humidity, and nutrient availability, enabling researchers to study the dynamic interactions that influence plant development. By leveraging techniques in predictive pattern recognition, models can be developed to accurately capture and represent plant growth behaviors.

1.1 Background

1.1.1 Rebuttal: Comparing Controlled and Open Field Agricultural Systems

To contextualize the focus of this review, it is necessary to contrast the differences in the observational parameters between controlled environment and open field agricultural systems. Open field environments are characterized by

1.1.2 Predictive Modeling in Agriculture

Predictive models within plant research are often dominated by a frequentist approach to statistical inference, where empirical observations and static conditions are tested against hypotheses to determine the probable outcome through confidence intervals or point estimations. This statistical approach is favored within the field for its simplicity in analysis and interpretation. Frequentist methods, including various forms of regression modeling, provide robust tools for analyzing experimental data and have become the standard in plant research due to their well-established procedures and ease of use [13, 14]. However, this experiment-centric approach is often limited when applied toward plant observation tasks that involve complex data structures, as with analyzing sequential and spatiotemporal images collected within the domain of high-throughput plant phenotyping (HTPP). Moreover, plant growth modeling and forecasting is an ill-posed problem where the trajectory of a plant’s development does not have a unique solution. Dynamic growing conditions and stochastic elements within plant development are prevalent due to the inherent variability and uncertainty that exists in biological systems. For example, plant growth is influenced by numerous interacting factors, including genetic, environmental, and management variables, which can lead to highly complex and non-linear relationships. Thus methods for regularization of the problem should be explored.

Within the context of simulations, computational models aim to predict plant traits, such as biomass accumulation, crop yield, presence of diseases, and overall growth dynamics, during various development stages under varying environmental conditions [15]. Some classical methods have approached this task through a combination of process models that simulate physiological principles and data-driven empirical models, typically obtained through a frequentist framework, to form various subsystems of an overarching hybrid model, offering a detailed understanding of how plants respond to different environmental stimuli at a field scale. Models such as the Crop Environment Resource Synthesis (CERES) [16] and the Decision Support System for Agrotechnology Transfer (DSSAT) have been extensively used to forecast plant growth by integrating weather data, soil conditions, and crop management practices [17, 18]. These models have proven instrumental in




Figure 1: Overview of two modeling paradigms in plant growth research. The Frequentist approach emphasizes fixed data collection, hypothesis testing, and multivariate inference within an experimental-centric framework. In contrast, the probabilistic approach employs dynamic data integration, iterative updates to prior distributions, and posterior evaluation for decision-making, enabling data-driven adaptability over time.

evaluating the potential impacts of climate change on crop productivity, guiding decision-making in traditional agriculture. However, these methods often demonstrate the various limitations of an experiment-centric approach.

The first limitation of this classical framework is that the underlying inference models are typically discriminative. The parameters of these models are treated as fixed variables that characterize boundaries used to determine the most probable outcome based on the frequencies of occurrence in sample data. This lack of flexibility in describing data distributions makes these models less effective in representing systems that involve stochastic processes, as with the various biological processes that constitute plant growth. As a result, the deterministic process models derived from these empirical outcomes are often sensitive to initial conditions and are constrained to rigid scopes of inference, making it unsuitable for distant future predictions and the synthesis of realistic data. The second limitation is that the outputs of such models are typically expressed through low-dimensional or scalar values, such as yield, biomass, and vegetative index. While this allows for simple comparative analysis and assessments of seasonality of data points, insights toward spatial information and links

between input conditions and plant structural developments are lost. The third limitation is that frequentist-based inference models do not intrinsically incorporate mechanisms to update parameters and integrate new information dynamically. This paradigm makes these models less effective for sequence analysis, as they provide no insights on prior information and treat parameters and datasets as static. In contrast, probabilistic approaches, such as those based on Bayesian inference (BI), explicitly quantify uncertainties and dynamically update with new data [19]. The static nature of the classical approach to modeling plant growth can lead to the underestimation of uncertainty, as the underlying assumptions about the "true" distribution of data may change, compounded by the cumulative effect of decisions made during earlier stages of inference in sequence analysis [20]. An overview of these dynamics between modeling paradigms are illustrated in Figure 1.

In this review, we explore these paradigms and assess the body of works available for various modeling tasks that require spatiotemporal analysis and pattern recognition of plant traits during its growth and development. We further propose insights to advance future works within this field and provide a perspective towards a model-centric probabilistic approach to plant growth pattern recognition and modeling through structured 2D and 3D data representations.

2 Rationale and Approach

Recent state-of-the-art (SOTA) developments made within the field of HTPP have led to the establishment of data collection platforms and protocols for multimodal datasets, which include image sensor and 3D point cloud data [21–25]. With the introduction of these complex datasets and multi-modal data fusion techniques, plant phenomic research has shifted towards computer vision (CV) and sensor-based methods for efficient assessment of plant traits when compared to traditional, labor-intensive approaches [26, 27]. Advances in CV, particularly through the use of deep learning (DL) models such as convolutional neural networks (CNN), have enabled the extraction of phenotypic traits such as leaf area, plant height, and disease symptoms from high-resolution images with good accuracy [28]. 3D imaging techniques, including structure-from-motion (SfM), stereo vision, and light detection and ranging (LiDAR) scanning, provide methods for processing spatial information to produce reconstructed 3D representation of plant architectures [29].

While these datasets are applied towards methods for assessing crop quality and quantifying biophysical properties throughout various stages of a plant’s growth cycle, the scope of phenomics research is often limited to classification tasks such as detection and segmentation, with limited work done on CV-based predictive forecasting. In contrast, this review focuses on work that explore the spatiotemporal components of plant phenomics research and the representation of plant traits through time.

2.1 Literature Overview

A rigorous screening process was applied to articles indexed within the following publication databases; SCOPUS, Web of Science (WOS), PubMed, and Google Scholar. A total of 50 method papers on plant growth modeling and 22 temporal dataset papers were selected and reviewed (Figure 2). These




Figure 2: Breakdown of publications reviewed, categorized by modeling approaches and dataset types. The figure highlights the distribution of works across regression models, latent representation models, functional-structural plant models, and computer vision-based techniques, as well as their focus on temporal scalar, 2D image-based, and 3D point cloud datasets.

papers include articles published within the last 10 years, 2015-2025, with a particular focus on assessing plant traits over time, where temporal analysis is a key component to the method or dataset. Articles in which traits are observed sparsely over time, for example traits observed once at the beginning of an experiment and once after harvest, are excluded from this review, opting for consistent, relatively high temporal resolution tracking of traits over time.

2.1.1 Plant Growth Representations and Datasets

A repository of recently published datasets on various temporal plant growth dynamics was compiled for this review and is shown in Table 1. This repository establishes several representations of observable plant traits which can be categorized into broader segments and provides insights towards the current focus of plant phenomics research. These data representations include: 3D representations of plant structures, typically expressed through 3D point clouds; 2D representations of plant structures expressed through images; and finally, the most common amongst a broader spectrum of classical methods, is 1D representation of various traits expressed through quantitative scalar data.

2.1.2 Limitations

Due to the exploratory nature of this review, the number of works available for a full systemic review of novel and SOTA techniques within this developing field is limited. This review is constrained to published research and datasets, which may lead to an inherent bias towards studies and datasets that have achieved publication rather than those that may be equally relevant but remain in preprint stages, unpublished, or private.

2.2 Scope of this Review

This review serves as a comprehensive examination of the field of pattern recognition in plant growth. We offer a prospective that utilizes the available SOTA methods towards a probabilistic approach through image-based simulation and generative modeling. The review is organised as such:

- We define plant growth modeling through various temporal paradigms and examine the prominent methods explored for scalar traits.
- We examine the application of these modeling paradigms towards 2D and 3D representations.
- We explore progress made within the developing field of probabilistic and generative computer-vision that utilize image-based forecasting.
- We provide a prospective towards the current state of research and propose a path for future works within the field.

3 Temporal Modeling Paradigms of Quantitative Traits

The abstraction of various plant biophysical properties into 1D quantitative traits provides a way to produce model parameters with functional meaning for a wide application of plant processes. Within plant phenomics, these applications can range from the molecular level with gene-protein expressions and phenotype mapping; the plant level in which we observe various plant-environment interactions; and the phenological level, where cyclical patterns of vegetative traits over a population within an ecosystem are mapped. In typical HTPP methods, scalar quantitative traits are inferred from high dimensional data such as RGB and multi-spectral images collected over time [51–54]. These low dimensional quantitative traits can be assessed through time and represented through various temporal modeling paradigms.

3.1 Regression Models

A common approach to simulate and predict the trajectory of these traits through time is through some form of regression-based inference model. Much of the works covered in this review describe methods for pattern recognition of plant growth that adhere to this framework. Phenotypic traits, denoted as y , can be expressed by explicitly fitting one of various forms of regressions (Equations 2-4). We can define the configuration of independent parameters that contribute to the expression of these traits, as $X = \{x_i \mid i \in \{1, 2, \dots, N\}\}$ for N parameters. For instance, y can represent a phenotypic trait such as plant height or leaf area, and x_i represents the expression level of gene i . In order to model these effects continuously through time, t , a generic regression model is given by Equation 1. This regression approach provides an empirical basis for fitting discrete observation data and determining the fixed components of models such as regression and spline coefficients, β and γ , respectively. Discrepancies between observed data and modeled outputs can be described as noise through error term, ϵ .

Table 1: Comprehensive list of temporal plant research datasets, published between 2015 and 2025, capturing various growth dynamics and traits.

Dataset	Crop	Traits	Data Type
Sun et al. (Soybean-MVS) [30]	Soybean (<i>Glycine max</i> var. <i>DN251</i> , <i>DN252</i> , <i>HN48</i> , <i>HN51</i>)	Segmented plant structure (leaf shape, stem, branches)	Point cloud (3D)
Schunck et al. (Pheno4D) [31]	Maize (<i>Zea mays</i>), <i>Solanum lycopersicum</i>	Segmented plant structure (leaf shape, stem, branches)	Point cloud (3D)
Sultan et al. (Shoots and Roots) [32]	Maize (<i>Zea mays</i>), <i>Solanum lycopersicum</i>), rice (<i>Oryza sativa</i>), Bala, Arabidopsis (<i>Arabidopsis thaliana</i>)	Root system traits and structure	Point cloud (3D)
Uchiyama et al. (KO-MATSUNA) [33]	Komatsuna (<i>Brassica rapa</i> var. <i>perviridis</i>)	Plant structure	Multi-view RGBD images and annotated masks (2D/3D)
Jayasuriya et al. [34]	Capsicum (<i>Capsicum annuum</i> L.)	Plant structure	RGBD images (2D/3D)
Kierdorf et al. (GrowliFlower) [33]	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)	Plant structure	RGB and multi-spectral images (2D)
Shadrin et al. [35]	Lettuce (<i>Lactuca sativa</i>)	Leaf growth and leaf segmentation	Top-down RGB images and annotated masks (2D), and environmental sensor readings (tabular)
Genze et al. [36]	Maize (<i>Zea mays</i> L.), Sorghum (<i>Sorghum bicolor</i>)	Plant structure	Top-down RGB images and annotated masks (2D)
Thoday-Kennedy et al. [37]	Safflower (<i>Carthamus tinctorius</i> L.)	Plant morphology and salt response	Multi-view RGB images (2D)
Larm et al. [38]	Maize (<i>Zea mays</i>), Wheat (<i>Triticum aestivum</i>)	Root system traits and structure	RGB images and ground-penetrating radar map (2D), soil sensor data (tabular)
Parmentier et al. [39]	General vegetation in Arctic ecosystems	Vegetative phenology and climate interactions	RGB images (2D) and environmental vegetative index data (tabular)
Craig et al. [40]	Maize (<i>Zea mays</i> L.)	Various agronomic traits and vegetative index	Scalar (tabular)
Dépigny et al. (Plantain-Optim) [41]	Plantain (<i>Musa acuminata</i> × <i>Musa balbisiana</i>)	Various agronomic traits and biomass	Scalar (tabular)
Ngaba et al. [42]	Sugarcane (<i>Saccharum officinarum</i>)	Various agronomic traits and biomass	Scalar (tabular)
Newman et al. [43]	Multiple field-crop species	Long-term agronomic traits and biomass	Scalar (tabular)
Campbell et al. [44]	Maize (<i>Zea mays</i>)	Photosynthetic function and vegetative index	Scalar (tabular)
Manjarrez-Sanchez et al. [45]	Greenhouse Tomato (<i>Lycopersicon esculentum</i>)	Photosynthetic function and agronomic traits	Scalar (tabular)
Niu et al. [46]	Maize (<i>Zea mays</i>)	Long-term phenological and climate interactions	Scalar (tabular)
Zhu et al. [47]	Multiple field-crop species	Long-term phenological and climate interactions	Scalar (tabular)
Mergner et al. [48]	Arabidopsis (<i>Arabidopsis thaliana</i>)	Gene expression	Scalar (tabular)
Castro et al. [49]	Camu-camu (<i>Myrciaria dubia</i>)	Gene expression	Scalar (tabular)
Peng et al. [50]	Peanut (<i>Arachis hypogaea</i>)	Gene expression	Scalar (tabular)

$$y(t) = f(t) + \epsilon(t) \quad (1)$$

$$f(t) = \sum_{i=0}^n \beta_i t^i \quad (\text{Polynomial}) \quad (2)$$

$$f(t) = \beta_0 + \beta_1 t + \sum_{j=1}^K \gamma_j S_j(t) \quad (\text{Spline}) \quad (3)$$

$$f(t) = \frac{1}{1 + e^{-(\beta_0 + \beta_1 t)}} \quad (\text{Logistic}) \quad (4)$$

174 The most common of these include non-linear regressions models that capture the typical patterns
 175 seen within plant growth traits. Following the conventions of a frequentest approach, sources of
 176 variation amongst growth patterns are hypothesized and made explicit before the collection of data.
 177 Non-experimental parameters, such as environmental and spatial variations, are standardized or
 178 corrected. Two works, Perez et al. [55] and Kar et al.[56], have proposed the implementation of
 179 spatial analysis of trials with splines (SPATS) for correcting spatial variations amongst experimental
 180 groups as a data pre-processing step before performing temporal modeling. The fixed but unknown
 181 statistics that describe the data are obtained empirically to fit the regression function for longitudinal
 182 trait predictions. The most prevalent implementations for modeling non-linear time-series patterns
 183 of scalar growth traits are spline-based forms, with common application of the model demonstrated
 184 in works such as Momen et al. [57] and Pauli et al. [58]. Polynomial forms, demonstrated in works
 185 such as Baba et al. [59] and Morales et al. [60], account for the majority of regressions used for
 186 longitudinal trait prediction.

187 **3.1.1 Separating Effects using Hierarchical Models**

188 To further disentangle the individual and compounding effects of various factors on plant growth
 189 trajectories, a hierarchical regression structure can be employed to represent plant traits for different
 190 levels of variation. This can be particularly useful for complex systems representing plant growth
 191 that is determined by multiple cumulative factors. For instance, plant traits such as height or leaf
 192 area can be modeled as responses to a combination of fixed and random effects. Fixed effects capture
 193 the consistent factors, such as the overall population-wide trends in growth, while random effects
 194 model the variation at each hierarchical level including genotype-specific effects and individual plant-
 195 specific variations. The random effects can vary across time, capturing the longitudinal nature of
 196 plant growth, while still maintaining a structure that allows for partial pooling of information across
 197 the hierarchy.

198 One such application of this is demonstrated by Equation 5 and shown in Figure 3. A hierarchical
 199 logistic regression is employed to represent plant growth at the population level, genotype level, and
 200 individual plant level. A generic form logistic transformation (Equation 4) can be applied towards
 201 the compounding fixed population effects, f_p , the random effects at the genotype level, $f_{pg}(t)$, and




Figure 3: Hierarchical logistic regression modeling of typical plant growth with fixed and random effects. This hierarchical approach highlights variability at multiple levels, with the logistic model capturing both population-wide trends and genotype-specific and plant-specific variations in plant traits such as height, biomass, or leaf area.

the random effects of each individual plants, $f_{pgi}(t)$, to describe the development of plant traits over time, $y(t)$. In this example β , L , and t_0 in Equations 6-8 are model parameters to be determined in order to fit our regressions to corresponding categorized data.

$$\log \left(\frac{y_{ijt}(t)}{L_{ij} - y_{ijt}(t)} \right) = f_p(t) + f_{pg}(t) + f_{pgi}(t) + \epsilon_{ijt}, \quad \epsilon_{ijt} \sim \mathcal{N}(0, \sigma^2) \quad (5)$$

$$f_p(t) = \beta_{p,0} + \beta_{p,1} \cdot (t - t_0) \quad (\text{Fixed Population Effects}) \quad (6)$$

$$f_{pg}(t) = \beta_{pg,0_i} + \beta_{pg,1_i} \cdot (t - t_0) \quad (\text{Random Genotype Effects}) \quad (7)$$

$$f_{pgi}(t) = \beta_{pgi,0_{ij}} + \beta_{pgi,1_{ij}} \cdot (t - t_0) \quad (\text{Random Individual Plant Effects}) \quad (8)$$

When modeling plant traits over time, fixed effects are commonly defined by a logistic regression to capture typical growth curve patterns and to distinguish temporal trends between categorical groups, such as genotypes. A typical implementation of such models is demonstrated in works such

as Baker et al. [61], in which the effects of genotypes are separated using Bayesian routines to estimate logistic parameters. Certain instances may require the disentanglement of complex effects between groups. In such cases, the mapping of these relationships require that regression models capture higher degrees of non-linearity. In Perez et al. [62], a P-spline-based hierarchical structure (Equation 3) is employed to model the effects not just at the population and genotype level but also the effects between individual plants, experimental groups and spatio-temporal variations. This approach allows for the simultaneous representation of growth curves at various levels of variability and the incorporation of shared parameters amongst each group effects on plant growth.

3.1.2 Pitfalls of Regression-based Temporal Inference

Modeling plant growth through regressions provides meaningful interpretations of trends on labeled data points. However, temporal inferences made from these models suffer from the pitfalls of a post-hoc analysis. Following the typical experiment-centric modeling pipeline (Figure. 1), inference models are selected after data has been collected and are typically implemented to explain trends once experimental groups are found to be statistically significant. As such, various regression forms and their hyper-parameters are selected once this data is revealed in order to achieve best fit. This post-hoc selection provides an inherent risk of overfitting due to the lack of a priori model constraints, leading to interpretations that may capture transient or idiosyncratic patterns rather than robust, underlying temporal trends. As a result, post-hoc regression models lack the generalizability necessary to predict future data accurately, particularly in time-series with complex dependencies or irregular periodicity.

Another pitfall involves the interpretability of the regression parameters. Temporal inferences often rely on the assumption that parameters estimated from the data capture the true characteristics of the underlying temporal process. However, in the absence of predefined model assumptions, these parameters can be sensitive to random variations or transient anomalies in the dataset. This sensitivity can result in misleading estimates of trend strength, direction, or duration, especially in nonlinear regression models, which are more prone to parameter instability when applied post hoc. Consequently, regression-based temporal models may yield unreliable estimates that lack external validity, thereby compromising the inference drawn from these analyses.

Furthermore, temporal data often contain complex dependencies due to autocorrelation, seasonal effects, or multi-modal factors. When regression models are employed in a post-hoc framework, these dependencies are frequently under accounted for, leading to temporal confounding. Such confounding poses a significant challenge as it obscures the causative factors underlying observed trends and further limits the model’s applicability beyond the studied period. Works such as Nesteruk et al. [63] demonstrate the use of autoregressive model improved predictions when compared to other non-correlated regressions.

3.2 Latent Representation Models

Quantitative traits can be assessed through time by representing their complex temporal dependencies through some reduced latent representations, or hidden state. These representations can

be applied as a transformation on sequential inputs to derive predictive outputs. In this section we explore various methods for extracting latent representation from the temporal dependencies between scalar quantitative traits and the models used to perform sequence-based inference from these hidden states.

3.2.1 Dimensionality Reduction through Principal Component Analysis

Methods like those employed by Adak et al. [64], aim to extract key features from time-series data that correspond to a reduced representation of growth rates using principal component analysis (PCA). In this context, PCA is a pattern recognition technique that captures the temporal components of the evolution of traits, such as key moments found within vegetative indices (VI) of plants of differing genotypes. These reduced representations effectively act as growth-rate phenotypes, embodying the entirety of a plant’s growth cycle and can be incorporated as an independent variable to improve accuracy of genomic and phenomic prediction models. However, meaningful interpretations of these phenotypes, especially for simulation purposes, are lost as these representations are abstracted into a reduced latent form. Interpretations of these representations back to the time domain for inference requires a decoding process. We explore such a process for decoding a latent representation in Section 3.2.2.

3.2.2 Neural Network-based Models

More recently, with advancements in computational systems and optimization algorithms, NNs have been employed in models to perform powerful opaque mappings between inputs and outputs. As such, they have expanded the range of tools available for dimensionality reduction and feature extraction. While some early implementations of simple feed forward DL systems are employed for pattern recognition, they often perform inference on fixed time windows which result in the same pitfalls of regression systems described in Section 3.1.2. For instance, Matsumura et al. [65] employs a multi-layer perceptron (MLP) with inputs of early season parameters in the form of a fixed window to forecast end of season outputs. While this approach can capture relationships within the window, it fails to incorporate longer-term temporal dependencies beyond the window itself. This results in models that are limited in their ability to adapt to changes outside the fixed period, potentially leading to overfitting to short-term patterns and reduced generalizability to future time steps or different seasons.

Recurrent Neural Networks To resolve this issue of fixed windows, sequence-based NNs, such as recurrent neural networks (RNN) and transformers, have been the focus of recent works to assess traits through time. Two main implementations of these networks include direct sequence predictions and latent sequence encoding (Figure 4). The former utilizes NNs for sequence generation tasks such as time-series forecasting. In the context of RNNs, prior sequential information is defined by the weights of an NN and represented as a hidden state, h , which embodies all cumulative prior sequential information. This h is used to transform the next given input, X_t , to predict the next element in the sequence, Y_{t+1} (Figure 4(a)). Various recent works within plant yield forecasting implement a




Figure 4: Directed graph models (folded) of the typical recurrent neural network (RNN) pipeline for sequence prediction and encoding. (a) Shows the sequence prediction task, where inputs X are processed through recurrent connections to generate hidden states h , leading to intermediate outputs O and a final output Y . This demonstrates how RNNs can predict each subsequent element in the sequence based on prior context. (b) Shows a sequence encoding process, where X is processed into a final encoded latent variable Z , capturing the entire sequence’s context in a single vector. Z can be decoded to obtain Y .

subset of RNNs known as long short term memory (LSTM) networks to predict the overall real-time trajectory of plant growth traits [66, 67].

The second implantation of these NNs aims to capture and represent temporal relationships through a fixed latent variable, Z (Figure 4(b)). Z is obtained by processing sequential information through an RNN and is subsequently derived from h , also typically defined in the form of the weights of a MLP between the two states. This representation captures the entirety of the sequence as a context driven conditional and can be decoded to perform inference on various tasks. One such application is demonstrated by Gong et al. [68], in which end of harvest yield is forecasted from a latent embedding of sequential environmental parameters through a temporal CNN decoder. Moreover, the decoding process can also be applied to task in which it is useful to predict future sequences of the same trait within a sequence-to-sequence approach. Works such as Bashar et al. [69], show the utility of such approach using a LSTM-based networks as both a sequence encoder to capture significant temporal relationships within plant growth development into a low dimensional embedding and as a decoder to obtain the next sequence of predicted outcomes from these embeddings.

Transformers and Attention Mechanisms It is worth noting the applicability of transformers and attention mechanisms for analyzing sequential data similar to RNNs. However, unlike RNNs, transformers leverage a self-attention mechanism to process entire sequences in parallel, enabling them to capture long-range dependencies effectively as each element directly attends to all others [70].

Although few studies have applied these architectures towards temporal plant trait modeling, their ability to weigh the importance of different time steps dynamically demonstrates their potential for analyzing plant growth patterns influenced by both recent and distant environmental factors. This adaptability suggests significant opportunity for advancing plant growth forecasting, particularly for complex, multi-modal time-series data. Their use within higher-dimensional structured output predictions is explored in Section 4.2.1.

3.2.3 State-Space Models

State-space models (SSM) define a broader categorization of sequence processing approaches similar to RNNs. However, rather than learning an implicit hidden state, which captures prior information as memory, SSMs are defined by an explicit transition state model and observation model. These models can be expressed flexibly between deterministic functions for dynamical systems or as probabilistic methods for describing complex stochastic processes (Table 2). The transition state model defines how latent states, z_{t+1} , evolve over time, and their dependence with the prior input, x_t , and latent state, z_t . The observation model relates the latent state to observed variables, y_t . SSMs assume a Markov property, in which the future state of a system depends only on its current state. As such, the process has no memory beyond the current state as with RNNs. This Markov property can be relaxed with the inclusion of memory states of prior sequences in the transition state model (Section 4.2.2).

Table 2: Comparison between the components of deterministic and probabilistic state-space models.

	Deterministic SSM	Probabilistic SSM
State Transition Model	$\mathbf{z}_{t+1} = f_{\theta}(\mathbf{z}_t, \mathbf{x}_t) + \eta_t$	$\mathbf{z}_{t+1} \sim p_{\theta}(\mathbf{z}_{t+1} \mathbf{z}_t, \mathbf{x}_t)$
Process Noise	$\eta_t \approx 0$	$\eta_t \sim \mathcal{N}(0, \Sigma_{\eta})$
Observation Model [†]	$\mathbf{y}_t = g_{\theta}(\mathbf{z}_t) + \xi_t$	$\mathbf{y}_t \sim p_{\theta}(\mathbf{y}_t \mathbf{z}_t)$
Observation Noise [†]	$\xi_t \approx 0$	$\xi_t \sim \mathcal{N}(0, \Sigma_{\xi})$

[†] Models may introduce stochasticity to a core deterministic functional. In such cases, process and observation noise do not equal 0.

Within the context of plant growth modeling and phenological pattern recognition, SSMs present an approach that captures the gradual changes in traits by modeling transitions within latent growth states [71, 72]. This approach has the advantage of continuously updating estimates of latent variables based on context driven observations such as environmental factors. For instance, Viaud et al. [73] proposes that complex plant growth dynamics, including the variability from noisy environmental conditions can be better represented through a classical SSM framework, when compared to traditional models. More specifically, BI can be applied to estimate the parameters, θ , of a hidden Markov model (HMM), through a particle MCMC method. As such the stochastic elements of plant growth can be captured through a probabilistic latent transition state. In recent works, NNs have been employed as the transition state models to capture complex non-linear relationships between environmental and plant trait development. Shibata et al. [74] introduces a deep SSM designed to predict sugar content in greenhouse tomatoes from inputs representing environmental conditions. In

this work it is suggested that a semi-supervised approach can be taken and paired with reinforcement learning for dynamic control tasks in CEA, where optimal decisions must be made in response to changing conditions.

3.2.4 Pitfalls of Representation Models

Assessing plant growth patterns through latent representations provides a way to capture the complex long-term temporal dynamics of trait development. The hidden and latent states, particularly of RNNs and SSMs, determine the transformations on prior states, which can be used to interpret how the system will change through time. As such, these methods improve upon classic regression systems by providing a context driven inference paradigm that resolves various issues of post-hoc analysis and allows for real-time predictions. However, the abstraction of these temporal representations into latent forms results in challenges of interpretability.

As the prior states are transformed into latent or hidden states, the connection between the model inputs and the output space can become unclear. For instance, PCA reduces temporal data to principal components, but these components often lack a direct explainable meaning, making it difficult to infer how individual traits evolve over time directly and require some additional mapping to outputs. This challenge is further exacerbated if mappings are performed through NN-based models, where latent and output states are learned through complex nonlinear weight transformations, leading to opaque predictions that prevent practical interpretations. Moreover, determining the inverse mapping of the model solution is not directly possible. For example, given the outcome of a plant's trait at a particular time, we cannot determine the input conditions and prior states that gave rise to the outcome directly, especially if latent states are determined through NNs or other highly abstracted models.

Integrating Domain-Specific Knowledge Future work may focus on the integration of domain-specific knowledge within these pattern recognition paradigms, especially in cases where meaningful interpretation of processes is important. These knowledge-driven models typically define various plant processes based on biological principles such as photosynthesis, respiration, and growth dynamics through mechanistic models [75]. Xing-Rong et al. [76] explores a combined data-driven and knowledge-based approach to resolve issues of strict calibration constraints, and incomplete representations of complex plant dynamics that result from purely mechanistic models. Much of these hybrid models are early in their conception, utilizing simple data-driven models, such as regressions or simple MLPs, to account for variability of outputs or as direct inputs, as an estimator, to a mechanistic model. Consequently, multiple models are often employed as separate components and consolidated through a two step process; one step to obtain the data driven term, and the second step for the mechanistic output. This approach has the disadvantage of higher computational cost and potential loss of capturing hierarchical or interdependent effects, especially if training processes are separated. Thus, there are significant potential for further contributions of advanced data-driven techniques to improve methods within a hybrid paradigm.




Figure 5: High-level conceptual overview of a context driven simulation framework for structured output representation through time.

4 2D/3D Structured Representations through Time

From low-dimensional scalar traits, we move toward representing the structural elements of plant growth through higher order data structures, such as 2D images and 3D point clouds. This is particularly useful for describing the development of plant structures through time, providing a graphical representation and visual insights to plant dynamics that are normally abstracted away with vectorized quantitative traits. The spatiotemporal simulation of these attributes offers a holistic assessment of strategies exhibited by the various plant behaviors based on contextual stimuli or signals. In this section, we take the foundation of pattern recognition techniques explored in Section 3 and apply them toward tasks that represent plant morphologies as structured outputs. This typically involves the addition of some rendering process which defines our plant representation model and is driven by a contextual process (Figure 5). As such, we explore the various works on the 2D or 3D structured plant representations, as well as works that attempt to model the entire context driven spatiotemporal pipeline for simulating plant growth.

4.1 Functional-Structural Plant Models

Functional-structural plant models (FSPM) represent a class of plant representation models that construct the structural and geometric elements of a plant from its mechanistic processes [77, 78]. Particularly, these models capture the interaction between the plant’s physiological processes, such as photosynthesis, water transport, and nutrient allocation, and its development of plant organs such as leaves, branches, and roots over time [79–82]. By coupling these two aspects, FSPMs provide a comprehensive view of plant behavior in response to various dependent parameters, allowing for the visualization of resource allocation and its effect on growth patterns. More recently, empirical methods have been integrated to the development process of FSPMs to establish the relationships between features extracted from HTPP data [83]. As such, an emergence of hybrid data-driven, mechanistic FSPMs have been applied towards plant growth simulations.

4.1.1 Process-driven Differential Growth Models

One application of these FSPMs is to capture the morphological developments within plant organs through differential growth models. The development and emergence of complex 3D morphologies within plant structures through time can be represented through numerical models, such as finite element methods (FEM), to describe differential growth across localized regions. The basis of the mechanistic models that determine these differentials are typically derived through experimental observations, utilizing image-processing techniques such as optical flow to obtain growth parameters from HTPP data [84]. Changes are mapped spatially and reconstructed from the distribution of conditions which influence the final shape of plant organs. One example of such modeling is demonstrated in Huang et al. [85], where biomechanical conditions are observed to determine the shape formations of leaves and petals. In this work, the modeling of growth differentials can be used to describe and reconstruct twisting, helical twisting, and edge-waving configurations within leaf structures through FEM simulations.

More often, this approach for plant structural modeling is applied toward developing fate and growth maps of cells within individual plant organs to simulate localized development [86]. For instance, in the work done by Kierzkowski et al. [87], the dynamics of genomic-proteomics pathways of two homeobox genes and their effects on promoting, inhibiting, and differentiating plant cells are modeled through spatial growth differentials using FEM. This phenomena is modeled through time and driven by the localized stimuli of plant hormone, namely auxin, concentrations. In this case, this model is applied toward developing a deep understanding of the causal effects of these specific signal pathways through empirical methods by mapping signals through time.

4.1.2 Rule-based Lindemyer-Systems

Another common implementation of FSPMs involves the application of a rule-based modeling paradigm known as the Lindemyer-systems (L-systems). L-systems define a formal set of recursive rules that specify the configuration of plant structures based on a string of symbolic characters. These rules guide the growth and branching patterns of simulated plants through iterative processes that aim to generalize and represent the realistic biological development and propagation

421 of plant structures [88, 89]. By repeatedly applying these recursive rules, L-systems can simulate
 422 natural variations and intricate details found in plant morphology, making them a powerful tool in
 423 computational botany and generative modeling.

424 A basic example is given by Algorithm 1, which establishes a generalized execution of a simple
 425 L-system, where Figure 6 shows the visual rendering of a rule set commonly used for describing a
 426 simple branching pattern in 2D.

Algorithm 1 Generate L-System String

Require: Axiom (initial string) S , number of iterations N , set of production rules P

Ensure: Final string after applying rules

```

1: Initialize  $current \leftarrow S$ 
2: for  $i = 1$  to  $N$  do
3:    $next \leftarrow$  empty string
4:   for each character  $c$  in  $current$  do
5:     if there is a rule  $c \rightarrow replacement$  in  $P$  then
6:       Append  $replacement$  to  $next$ 
7:     else
8:       Append  $c$  to  $next$ 
9:     end if
10:  end for
11:   $current \leftarrow next$ 
12: end for
13: return  $current$ 

```




Figure 6: Demonstration of plant-like structures generated using L-Systems with varying parameters. The axiom $S = "F"$ and production rule P are iterated N times with specified angles. Examples show branching patterns for $N = 2, 3, 4$ and angles $15^\circ, 30^\circ, 40^\circ$, demonstrating the influence of parameterized rules on the rendering process. The symbols represent drawing actions.

This algorithm can be expanded to include parameterized production rules that encapsulate complex relationships at each iteration of the simulated plant branching. For example, various works have developed production rules that dictate tree organ formation based on a source-sink dynamic between existing plant organs at previous iterations, which is then subsequently rendered for the 3D simulation of various tree cultivars [90, 91]. More recent work has expanded on production rules using sink-source dynamics by incorporating context-driven systems such as environmental and management conditions, soil nutrient availability, and distribution to improve the dynamic tree simulation process [92–94].

Furthermore, works such as Hitti et al. [95] and Mußmann et al. [96] introduce an adaptation of the L-systems paradigm that captures the dynamic development of plant structures using agent-based interactions. Plant structures, represented as agents, can independently interact with various stimuli. Within such systems, agent-based feedback mechanisms, such as reinforcement learning, can be used to train model growth patterns based on external stimuli.

4.2 Computer Vision Approach to Plant Modeling

FSPMs typically rely on procedural methods for the visual rendering process of the plant simulation pipeline (Figure. 5). Although the outcomes of such models are interpretable and result in visually consistent structures, the outputs of such methods are often derived from deterministic solutions that lack stochastic variability. Such solutions tend to drift from biological realism, particularly when simulating complex interactions or environmental responses, requiring the integration of experiment data to drive realistic outputs [78]. Alternatively, some initial efforts have been made towards a




Figure 7: Directed graph model (unfolded) of a Convolutional Recurrent Neural Network (CRNN) implementation for predicting the next sequence in a plant scene. The model includes convolutional layers (ConvL₁, ConvL₂, ..., ConvL_n) for feature extraction and encoding, followed by a recurrent layer for temporal processing of sequential frame inputs, X , at $t_0, t_1, \dots, t_n$. A decoder module with deconvolution layers (DeConvL) reconstructs the output frames, Y , at t_{n+1} based on learned spatiotemporal features.

novel approach using data-driven CV for rendering the structural elements of the simulation task.

For instance, the task of plant growth modeling can be structured to fall within one of two classic problems found within the domain of predictive CV and video analysis; frame-to-frame translation and frame synthesis. Within this perspective, the structural elements of plant growth can be represented as images of a temporal scene, or frames, where the goal of our methods is to predict the pixel-wise configuration of these frames, such that the corresponding scene describes the realistic spatiotemporal development of plant growth. These methods do not rely on procedural rendering routines using pre-rendered graphics. Rather, the visual rendering process becomes a CV-based image translation task driven by empirical methods for predicting frames through features learned from high dimensional 2D or 3D HTPP data. We explore some early works which utilize such methods for predictive pattern recognition.

4.2.1 Next Frame Prediction

Some early works on CV-based plant growth forecasting include Sakurai et al. [97] and Wang et al. [98], which have proposed the adoption of convolutional recurrent neural networks (CRNN) for the task of image-to-image translation to predict the next frame of a plant growth scene given a set of prior frames from the same scene. This approach expands on the sequence-based prediction techniques described in Section 3.2.2 to assess spatial features from consecutive images. This is done by incorporating convolutional layers within a RNN framework to extract and encode features from

a set of input images, X , to a transitional hidden state h . The rendering of this hidden state is described by a decoding step which consists of an up-sampling process of h through deconvolutional layers. The typical generic implementation of the CRNN is shown in Figure 7. This approach combines pattern recognition techniques for forecasting plant parameters using RNNs with the visual rendering of plant structures through image translation methods. While CRNNs allow for spatial features to be captured through time, the deterministic mapping through convolutional and deconvolutional layers often result in artifacts and ambiguities within structures of the output images [99].

To mitigate problems in image quality, recent work have focused on improving the image generation component of model architectures, which correspond to the decoder of a CRNN. Yasrab et al. [100] proposes the use of a Generative Adversarial Network (GAN)-based network architectures for improved realistic image synthesis. Other works such as Kim et al. [101] propose multi-step networks that separate various components of the image generation task, such as shape estimation and image reconstruction, utilizing a transformer-based network for spatial pattern recognition. Although these approaches improve image generation quality, they do not address the challenges of ambiguous structures caused by a deterministic mappings. CV-based plant forecasting is an ill-posed problem, in which, there exists non-unique outcomes of a plants growth trajectory given the same initial set of frames. This is particularly evident under diverging environmental conditions. Moreover, predictions from the current models are made solely on the previous frames of the scene and further contribute to the prevalence of ambiguities during image translation.

4.2.2 Frame Synthesis Through Conditional Generative Models

A probabilistic approach can be taken to regularize the plant growth modeling problem and to guide the prediction process by incorporating stochastic elements caused by dynamic growing conditions. In such methods, a generative model is employed for the task of frame synthesis to predicted the expected frames of the plant growth scene. While generative models are typically used as data augmentation techniques for enhancing sparse agricultural datasets [102, 103], few works have explored the use of conditional generative models (CGM) for guided image synthesis. A probabilistic approach using a CGM is illustrated by a generic conditional variational autoencoder (CVAE) shown in figure 8. Instead of performing direct deterministic mappings between inputs and output frames, the objective of the CVAE is to learn a prior distribution, $p_\theta(z|x)$, where $z \sim p_\theta(z|x)$ is the latent representation of scene at a given point in time, t , directed by some conditional vector x . The prior distribution can be sampled such that it incorporates stochasticity through auxiliary variable, ϵ . The network structure explored within the domain of plant growth modeling typically consists of a probabilistic latent variable encoder and a deterministic decoder to render output frames, $\hat{y} \sim p_\theta(y|z, x)$.

Early works, including Miranda et al. [104] and Drees et al. [105], propose the adoption of a conditional GAN (cGAN), which introduces a discriminator with adversarial routines to the generic CVAE structure, for synthesizing field images and forecasting the state of the crop canopy. The conditional vector consists of fixed soil treatment embeddings set at the start of the growing season. Forecasts are made at discrete points in time driven by the initial states of the treatment conditions.




Figure 8: Conditional generative model architecture for guided plant scene simulation. The framework includes a latent encoder $F_\Phi(x, y \rightarrow z)$ that maps input data (X, Y) to a latent variable Z , and a decoder $G_\Theta(z, x \rightarrow y)$ that reconstructs ground truth (GT) outputs, Y . The latent space is defined by μ and σ^2 and incorporates auxiliary stochastic noise ϵ . Conditional vector, X , is directed by environmental parameters, $c(\tau)$, such as temperature, humidity, pH, and EC, over a time period $\tau = [0, t]$. The architecture allows for generating sampled outputs, $\hat{y}$, guided by X .

These works have demonstrated SOTA guided image generation for realistic image synthesis with variability in outputs defined by stochastic elements, or “*factors of variation*” (FOV). The frames synthesized from such models can be used in conjunction with image-based feature extraction models to provide quantitative estimation of biomass for a complete simulation process [106].

In these works, predictions are guided by a conditional vector comprised of treatment conditions, a prior frame state, and a time factor. The treatment embeddings are fixed at a particular point in time, typically set at the start of experimentation, and do not account for dynamic changes in conditions. For spatiotemporal simulation and trajectory estimation, the guidance of predictions without feedback can lead to larger variability when diverging from initial conditions due to the accumulated effects of compounding factors, including environmental conditions [107]. As such, this approach may be applicable for simulating images of end of harvest canopies under varying treatment conditions at a discrete point in time in the future, but lacks the dynamics for real-time continuous plant growth predictions useful for real world applications such as anomaly detection and intervention practices.

Debbagh et al. [108] proposes a sequence-informed approach which introduces dynamic and compounding environmental conditions as feedback to a CGM, guiding a continuous simulation process. In this work, the conditional vector, x , which guides the image generation of the CGM, is directed by a configuration of random variables, c , that describe dynamic factors that contribute to the development of the plants within the scene (i.e. environmental parameters). x is determined by encoding a sequence of these environmental parameters from the start of the simulation up to the




Figure 9: Directed graph model (folded) of a sequence-informed method for incorporating compounding environmental feedback to the conditional generative model (CGM). Environmental parameters, $c(\tau)$, are sequentially processed through an RNN encoder to obtain conditional vector, X . X guides the generation and latent encoding process of the CGM to produce output frames, Y . The generation edges correspond to the synthesis of outputs frames, Y , while the recognition edges correspond the process of learning the probability distributions of latent variable, Z . The recurrent connections allow the system to integrate sequential information for dynamic plant growth predictions.

point of inference, $c(\tau)$, where $\tau = [0, t]$. The encoding process consists of an RNN encoder, which allows the model to learn the compounding and sequential relationships between environmental conditions as a vector representation. The dependencies between each model component is illustrated by the directed graph model shown in Figure 9. This approach provides a framework that further regularizes the image generation task through dynamic context, allowing for the realistic synthesis of consecutive frames.

5 Perspectives

5.1 Data-Driven vs Experiment-Driven Approaches

Plant growth modeling has traditionally relied on experiment-driven approaches, where predefined hypotheses are tested using carefully controlled experiments. These methods provide valuable insights into specific biological processes but are often time-consuming, resource-intensive, and limited in scalability. By contrast, data-driven and model-centric approaches leverage large datasets and machine learning techniques to capture patterns and relationships that might not be immediately apparent within an experiment-driven framework [109]. The works presented in this review demonstrate a shift towards the latter, as challenges with dynamic updating and long-term predictions are prevalent in simple regression methods as described in Section 3.1.2.

A shift towards 2D image and 3D point cloud data for describing plant phenomena, as with HTPP datasets, requires model frameworks designed for interpreting and processing higher order

data-structures. This inherently moves prospective research towards the CV-based models described in Sections 4.2.1 and 4.2.2. However, the advantages of these data-driven approaches come with the limitation of reduced interpretability and a lack of the biological grounding inherent in experiment-driven methods. In contrast, FSPMs provide a robust simulation framework for structured representation that is highly controllable, but is completely reliant on experiment-driven approaches for developing knowledge-based rule sets.

Future research in this domain should prioritize the development of hybrid approaches that integrate the strengths of both paradigms. Two key directions are proposed to achieve this goal. The first involves designing model architectures that incorporate knowledge-informed mechanisms, enabling the integration of biological principles and domain expertise into data-driven frameworks. The second focuses on advancing methods for decoding and interpreting opaque models, ensuring that complex representation models provide outputs that align with biological processes and facilitate deeper understanding.

5.2 Dataset Gap

The shift toward data-driven approaches has placed datasets at the center of plant growth modeling and predictive frameworks. Large-scale, high-quality datasets are necessary for training DL models and validating their predictions. However, the effectiveness of these models are dependent on the availability, diversity, and representativeness of the data. Despite advancements in HTPP platforms, significant gaps remain in the scope, quality, and accessibility of existing datasets.

First, many datasets focus on single time points or static snapshots, which fail to capture the dynamic growth trajectories of plants over time. Existing publicly available datasets that capture longitudinal data (Table 1) are often scarce, and limited in sample frequency. This is particularly evident for datasets that capture multiple modalities of data including 2D images and 3D point clouds along with environmental parameters. Second, the diversity of crop species represented in current datasets is limited, with a heavy emphasis on staple crops such as maize (*Zea mays*) and soybean (*Glycine max*). This narrow scope restricts the ability of models to generalize across plant species with differing morphologies, such as root vegetables or perennials. Finally, the lack of consolidation of datasets across different research groups and platforms creates significant barriers to broader adoption and cross-comparative studies. Inconsistent formats, annotation practices, and metadata standards hinder the ability to combine datasets, thereby limiting the diversity and scale necessary for developing generalizable models.

To address these gaps, several considerations for developing future datasets are proposed. First, works should aim to apply standardization and open-access practices during dataset development to enable reproducibility and comparability across studies through benchmarking. High-frequency sampling should also be prioritized to produce longitudinal datasets that span entire growth cycles, improving the ability to model and predict temporal dynamics accurately. Furthermore, the development of multi-modal datasets, combining 2D images, 3D point cloud data, and environmental sensor measurements is suggested for capturing a holistic view of plant-environment interactions. Finally, integrating domain knowledge into dataset design to ensure that datasets not only capture

581 observable traits but also reflect underlying biological processes. Such biologically informed datasets
582 will enable models to generate predictions that are both empirically and biologically meaningful.

583 **5.3 Practical Applications**

584 The recent advancements in pattern recognition techniques explored in this review have significant
585 implications for practical applications. While traditional applications include crop yield forecasting
586 and climate simulations, these novel approaches expand the scope to include controlled environment
587 systems, field crop systems, and plant genomics and phenomics research. These advancements
588 allow for precise detection, simulation, and representation of plant growth processes in complex
589 environments.

590 **5.3.1 Anomaly Detection**

591 Anomaly detection and early interventions practices during crop development benefit from predictive
592 pattern recognitions techniques that incorporate dynamic feedback from environmental growing
593 conditions. The focus on the development of models that capture long-term temporal dependencies
594 through sequence-based methods, such as transformers and RNNs, provides an effective approach
595 to identifying deviations in plant growth trajectories. For instance, temporal anomalies such as
596 stunted growth or disease occurrence can be detected in advance by analyzing sequential data from
597 real-time HTPP platform measurements. Moreover, the incorporation multi-modal approaches to
598 combine imaging data with contextual environmental parameters presents a more comprehensive
599 assessment of plant health. These models improve the precision of anomaly detection and allow for
600 dynamic adjustments through intervention and control strategies. Such techniques are particularly
601 useful for mitigating potential yield losses and optimizing resource use in controlled environments
602 and field systems.

603 **5.3.2 Plant-Environment Simulation**

604 Simulation frameworks informed by generative and probabilistic models improve the ability to predict
605 plant growth under varying environmental conditions. Models like the CGMs presented in Section
606 4.2.2 enable the synthesis of realistic growth trajectories influenced by the spatial and dynamic
607 growing condition, including the distribution of light, nutrient conditions, temperature, and CO₂.
608 This provides *in silico* methods to inform the structural development process of a controlled plant
609 system and to develop strategies prior to conducting in-field operations. These frameworks allow
610 for resource optimization in controlled environments and improve the forecasting of long-term plant
611 responses to changing environments.

612 **5.3.3 Digital Twinning and Representation**

613 Digital twinning within plant phenomics typically consists of methods for 3D reconstruction of
614 plant structures taken at sparse static snapshots within its growth cycle. This is often due to
615 limitations in imaging techniques for capturing frequent 3D spatial data during its growth. The

structured representation methods explored within this review have implications on expanding this static framework to capture a continuous virtual replica of a plant system during its entire growth cycle. The continuous frame synthesis techniques explored in Debbagh et al. [108], allow for the interpolation of structured outputs between points of time. This presents an opportunity to generate digital plant structures with the guidance of few snapshots of plants. Moreover, the divergence of these plant structures from the snapshots can be explored through simulated growing conditions.

6 Conclusion

In this review, we provide a comprehensive examination of recent works and explore prospects towards advancing predictive pattern recognition techniques within plant phenomics research. These works cover various approaches that address challenges in processing and interpreting various representations of spatiotemporal plant data, which include longitudinal scalar growth traits and structured 2D images and 3D point cloud data. Within these works, novel modeling paradigms, which take advantage of recent advancements in computational systems, HTTP data collection and data fusion techniques, have presented significant improvements to the precision and accuracy of models for long-term forecasting and other simulation tasks. Notably, NN-based representation models and CGMs present a shift toward a data-driven approach to predictive pattern recognition that not only allows for dynamic updating for guiding predictions, but also allows for the synthesis of structured representation for spatial understanding of outputs.

Additionally, this review offers a perspective for opportunities to advance works within in this field. This includes perspectives on the integration of domain knowledge with data-driven approaches, and the emphasis on developing comprehensive, standardized datasets. Furthermore, we suggest practical applications within various domains to be explored in future works. The collection of these works and our perspective aims to consolidate and expand the understanding of spatiotemporal modeling frameworks and their role in advancing plant phenomics research and applications.

Acknowledgments

General

Special thanks to Anita Parmar, Ollivier Dyens, and the Building 21 community for their valuable contributions through engaging and thought-provoking discussions.

Author Contributions

M. Debbagh drafted the manuscript, prepared the figures and tables, and incorporated feedback from co-authors. S. Sun and M. Lefsrud reviewed and edited the manuscript, offering feedback to improve clarity and impact.

Funding

This work was supported by the Natural Sciences and Engineering Research Council of Canada (RGPIN-2021-03266, RGPIN-2021-03410); and Gardyn through the Mitacs Accelerate program (IT16220).

Tools

Figure 1 (<https://BioRender.com/y80s089>) and Figure 5 (<https://BioRender.com/y69z168>) were created in BioRender.com.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article.

References

- [1] Alisher Mirzabaev, Lennart Olsson, Rachel Bezner Kerr, Prajal Pradhan, Marta Guadalupe Rivera Ferre, and Hermann Lotze-Campen. *Climate Change and Food Systems*, pages 511–529. Springer International Publishing, Cham, 2023. ISBN 978-3-031-15703-5. doi:10.1007/978-3-031-15703-5_27.
- [2] C.A. O’Sullivan, G.D. Bonnett, C.L. McIntyre, Z. Hochman, and A.P. Wasson. Strategies to improve the productivity, product diversity and profitability of urban agriculture. *Agricultural Systems*, 174:133–144, 2019. ISSN 0308-521X. doi:10.1016/j.agsy.2019.05.007.
- [3] Kurt Benke and Bruce Tomkins. Future food-production systems: vertical farming and controlled-environment agriculture. *Sustainability: Science, Practice and Policy*, 13(1):13–26, 2017. doi:10.1080/15487733.2017.1394054.
- [4] Ofer Halperin, Alem Gebremedhin, Rony Wallach, and Menachem Moshelion. High-throughput physiological phenotyping and screening system for the characterization of plant–environment interactions. *The Plant Journal*, 89(4):839–850, 2017. doi:10.1111/tpj.13425.
- [5] Robert J Henry. Innovations in plant genetics adapting agriculture to climate change. *Current Opinion in Plant Biology*, 56:168–173, 2020. ISSN 1369-5266. doi:10.1016/j.pbi.2019.11.004. Biotic interactions AGRI 2019.
- [6] Anna Langstroff, Marc C Heuermann, Andreas Stahl, and Astrid Junker. Opportunities and limits of controlled-environment plant phenotyping for climate response traits. *Theoretical and Applied Genetics*, 135(1):1–16, 2022. doi:10.1007/s00122-021-03892-1.
- [7] Vishwas Anant Bapat, P. B. Kavi Kishor, Naravula Jalaja, Shri Mohan Jain, and Suprasanna Penna. Plant cell cultures: Biofactories for the production of bioactive compounds. *Agronomy*, 13(3), 2023. ISSN 2073-4395. doi:10.3390/agronomy13030858.

- [8] Yang Yang, Sedky H.A. Hassan, Mukesh Kumar Awasthi, Babu Gajendran, Monika Sharma, Min-Kyu Ji, and El-Sayed Salama. The recent progress on the bioactive compounds from algal biomass for human health applications. *Food Bioscience*, 51:102267, 2023. ISSN 2212-4292. doi:10.1016/j.fbio.2022.102267.
- [9] Mai Trinh Phuong Nguyen, Matthew Knowling, Nam N. Tran, Alexandra Burgess, Ian Fisk, Michelle Watt, Marc Escribá-Gelonch, Herve This, John Culton, and Volker Hessel. Space farming: Horticulture systems on spacecraft and outlook to planetary space exploration. *Plant Physiology and Biochemistry*, 194:708–721, 2023. ISSN 0981-9428. doi:10.1016/j.plaphy.2022.12.017.
- [10] Ryan M. Lefers, Mark Tester, and Kyle J. Lauersen. Emerging technologies to enable sustainable controlled environment agriculture in the extreme environments of middle east-north africa coastal regions. *Frontiers in Plant Science*, 11, 2020. ISSN 1664-462X. doi:10.3389/fpls.2020.00801.
- [11] Paul Zabel, Conrad Zeidler, Vincent Vrakking, Markus Dorn, and Daniel Schubert. Biomass production of the eden iss space greenhouse in antarctica during the 2018 experiment phase. *Frontiers in Plant Science*, 11, 2020. ISSN 1664-462X. doi:10.3389/fpls.2020.00656.
- [12] Bedrich Benes, Kaiyu Guan, Meagan Lang, Stephen P. Long, Jonathan P. Lynch, Amy Marshall-Colón, Bin Peng, James Schnable, Lee J. Sweetlove, and Matthew J. Turk. Multiscale computational models can guide experimentation and targeted measurements for crop improvement. *The Plant Journal*, 103(1):21–31, 2020. doi:10.1111/tpj.14722.
- [13] Reza Hoshmand. *Design of experiments for agriculture and the natural sciences*. Chapman and Hall/CRC, 2006. doi:10.1201/9781315276021.
- [14] Filmore E Bender. *Statistical methods for food and agriculture*. CRC Press, 1989. doi:10.1201/9781003073116.
- [15] Gianni Fenu and Francesca Maridina Mallocci. Forecasting plant and crop disease: An explorative study on current algorithms. *Big Data and Cognitive Computing*, 5(1), 2021. ISSN 2504-2289.
- [16] Bruno Basso, Lin Liu, and Joe T. Ritchie. A comprehensive review of the cereals-wheat, -maize and -rice models’ performances. In Donald L. Sparks, editor, *Advances in Agronomy*, volume 136 of *Advances in Agronomy*, pages 27–132. Academic Press, 2016. doi:10.1016/bs.agron.2015.11.004.
- [17] J.W Jones, G Hoogenboom, C.H Porter, K.J Boote, W.D Batchelor, L.A Hunt, P.W Wilkens, U Singh, A.J Gijsman, and J.T Ritchie. The dssat cropping system model. *European Journal of Agronomy*, 18(3):235–265, 2003. ISSN 1161-0301. doi:10.1016/S1161-0301(02)00107-7. Modelling Cropping Systems: Science, Software and Applications.

- [18] Konan Jean-Yves N’guessan, Botou Adahi, Arthur-Brice Konan-Waidhet, Satoh Masayoshi, and Nogbou Emmanuel Assidjo. Assessment of climate change impact on water requirement and rice productivity. *Rice Science*, 30(4):276–293, 2023. ISSN 1672-6308. doi:10.1016/j.rsci.2023.03.010.
- [19] David JC MacKay. *Information theory, inference and learning algorithms*. Cambridge university press, 2003. doi:10.1017/S026357470426043X.
- [20] Richard McElreath. *Statistical rethinking: A Bayesian course with examples in R and Stan*. Chapman and Hall/CRC, 2018. doi:10.1201/9781315372495.
- [21] Achim Walter, Frank Liebisch, and Andreas Hund. Plant phenotyping: from bean weighing to image analysis. *Plant methods*, 11:1–11, 2015. doi:10.1186/s13007-015-0056-8.
- [22] Daoliang Li, Chaoqun Quan, Zhaoyang Song, Xiang Li, Guanghui Yu, Cheng Li, and Akhter Muhammad. High-throughput plant phenotyping platform (ht3p) as a novel tool for estimating agronomic traits from the lab to the field. *Frontiers in Bioengineering and Biotechnology*, 8, 2021. ISSN 2296-4185. doi:10.3389/fbioe.2020.623705.
- [23] Rijad Sarić, Viet D Nguyen, Timothy Burge, Oliver Berkowitz, Martin Trtílek, James Whelan, Mathew G Lewsey, and Edhem Čustović. Applications of hyperspectral imaging in plant phenotyping. *Trends in plant science*, 27(3):301–315, 2022. doi:10.1016/j.tplants.2021.12.003.
- [24] Manuel G. Forero, Harold F. Murcia, Dehyro Méndez, and Juan Betancourt-Lozano. Lidar platform for acquisition of 3d plant phenotyping database. *Plants*, 11(17), 2022. ISSN 2223-7747. doi:10.3390/plants11172199.
- [25] Ting Wen, Jian-Hong Li, Qi Wang, Yang-Yang Gao, Ge-Fei Hao, and Bao-An Song. Thermal imaging: The digital eye facilitates high-throughput phenotyping traits of plant growth and stress responses. *Science of The Total Environment*, 899:165626, 2023. ISSN 0048-9697. doi:10.1016/j.scitotenv.2023.165626.
- [26] Sruti Das Choudhury, Ashok Samal, and Tala Awada. Leveraging image analysis for high-throughput plant phenotyping. *Frontiers in Plant Science*, 10, 2019. ISSN 1664-462X. doi:10.3389/fpls.2019.00508.
- [27] Justyna Jadwiga Olas, Franziska Fichtner, and Federico Apelt. All roads lead to growth: imaging-based and biochemical methods to measure plant growth. *Journal of Experimental Botany*, 71(1):11–21, 09 2019. ISSN 0022-0957. doi:10.1093/jxb/erz406.
- [28] Arti Singh, Baskar Ganapathysubramanian, Asheesh Kumar Singh, and Soumik Sarkar. Machine learning for high-throughput stress phenotyping in plants. *Trends in plant science*, 21(2):110–124, 2016. doi:10.1016/j.tplants.2015.10.015.

- [29] Shangpeng Sun, Changying Li, Peng W. Chee, Andrew H. Paterson, Yu Jiang, Rui Xu, Jon S. Robertson, Jeevan Adhikari, and Tariq Shehzad. Three-dimensional photogrammetric mapping of cotton bolls in situ based on point cloud segmentation and clustering. *ISPRS Journal of Photogrammetry and Remote Sensing*, 160:195–207, 2020. ISSN 0924-2716. doi:10.1016/j.isprsjprs.2019.12.011.
- [30] Yongzhe Sun, Zhixin Zhang, Kai Sun, Shuai Li, Jianglin Yu, Linxiao Miao, Zhanguo Zhang, Yang Li, Hongjie Zhao, Zhenbang Hu, Dawei Xin, Qingshan Chen, and Rongsheng Zhu. Soybean-mvs: Annotated three-dimensional model dataset of whole growth period soybeans for 3d plant organ segmentation. *Agriculture*, 13(7), 2023. ISSN 2077-0472. doi:10.3390/agriculture13071321.
- [31] David Schunck, Federico Magistri, Radu Alexandru Rosu, André Cornelißen, Nived Chebrolu, Stefan Paulus, Jens Léon, Sven Behnke, Cyrill Stachniss, Heiner Kuhlmann, and Lasse Klingbeil. Pheno4d: A spatio-temporal dataset of maize and tomato plant point clouds for phenotyping and advanced plant analysis. *PLOS ONE*, 16(8):1–18, 08 2021. doi:10.1371/journal.pone.0256340.
- [32] Sam Sultan, Joseph Snider, Adam Conn, Mao Li, Christopher N. Topp, and Saket Navlakha. A statistical growth property of plant root architectures. *Plant Phenomics*, 2020, 2020. doi:10.34133/2020/2073723. URL <http://plant3d.navlakhlab.net/roots/>.
- [33] Hideaki Uchiyama, Shunsuke Sakurai, Masashi Mishima, Daisaku Arita, Takashi Okayasu, Atsushi Shimada, and Rin-ichiro Taniguchi. An easy-to-setup 3d phenotyping platform for komatsuna dataset. In *2017 IEEE International Conference on Computer Vision Workshops (ICCVW)*, pages 2038–2045, 2017. doi:10.1109/ICCVW.2017.239.
- [34] Namal Jayasuriya, Oula Ghannoum, Wen Hu, Norbert Klaue, Weiguang Liang, and Yi Guo. Pc4c.capsi: Image data of capsicum plant growth in protected horticulture. *Data in Brief*, 55: 110735, 2024. ISSN 2352-3409. doi:10.1016/j.dib.2024.110735.
- [35] Dmitrii G. Shadrin, Victor Kulikov, and Maxim V. Fedorov. Instance segmentation for assessment of plant growth dynamics in artificial soilless conditions. In *British Machine Vision Conference*, 2018.
- [36] Nikita Genze, Wouter K Vahl, Jennifer Groth, Maximilian Wirth, Michael Grieb, and Dominik G Grimm. Manually annotated and curated dataset of diverse weed species in maize and sorghum for computer vision. *Scientific Data*, 11(1):109, 2024. doi:10.1038/s41597-024-02945-6.
- [37] Emily Thoday-Kennedy, Adam M. Dimech, Sameer Joshi, Hans D. Daetwyler, David Hudson, German Spangenberg, Matthew Hayden, and Surya Kant. An image dataset of diverse safflower (*carthamus tinctorius* l.) genotypes for salt response phenotyping. *Data in Brief*, 46:108787, 2023. ISSN 2352-3409. doi:10.1016/j.dib.2022.108787.

- [38] Lena Lärm, Felix Maximilian Bauer, Normen Hermes, Jan van der Kruk, Harry Vereecken, Jan Vanderborght, Thuy Huu Nguyen, Gina Lopez, Sabine Julia Seidel, Frank Ewert, et al. Multi-year belowground data of minirhizotron facilities in selhausen. *Scientific Data*, 10(1): 672, 2023. doi:10.1038/s41597-023-02570-9.
- [39] F.-J. W. Parmentier, L. Nilsen, H. Tømmervik, and E. J. Cooper. A distributed time-lapse camera network to track vegetation phenology with high temporal detail and at varying scales. *Earth System Science Data*, 13(7):3593–3606, 2021. doi:10.5194/essd-13-3593-2021.
- [40] Valerie Craig, Hugh Earl, John Sulik, and Elizabeth A Lee. Hyperspectral time series datasets of maize during the grain filling period. *BMC Research Notes*, 15(1):152, 2022. doi:10.1186/s13104-022-06029-9.
- [41] Sylvain Dépigny, Frédéric Tchotang, Médard Talla, Désirée Fofack, David Essomé, Jean-Pierre Ebongué, Bernard Kengni, and Thierry Lescot. The 'plantain-optim' dataset: Agronomic traits of 405 plantains every 15 days from planting to harvest. *Data in Brief*, 17:671–680, 2018. ISSN 2352-3409. doi:10.1016/j.dib.2018.01.065.
- [42] Billy Ngaba, Mathias Christina, Alizé Mansuy, Julien Chetty, Mathilde Soulé, Marion Schwartz, Benjamin Heuclin, and Sandrine Auzoux. Experimental dataset of sugarcane-cover crop intercropping trials to control weeds in reunion island. *Data in Brief*, 48:109244, 2023. ISSN 2352-3409. doi:10.1016/j.dib.2023.109244.
- [43] Saul Justin Newman and Robert T Furbank. A multiple species, continent-wide, million-phenotype agronomic plant dataset. *Scientific data*, 8(1):116, 2021. doi:10.1038/s41597-021-00898-8.
- [44] Petya Campbell, Elizabeth Middleton, Karl Huemmerich, Lauren Ward, Tommaso Julitta, Peiqi Yang, Christiaan van der Tol, Craig Daughtry, Andrew Russ, Joseph Alfieri, and William Kustas. Scaling photosynthetic function and co₂ dynamics from leaf to canopy level for maize – dataset combining diurnal and seasonal measurements of vegetation fluorescence, reflectance and vegetation indices with canopy gross ecosystem productivity. *Data in Brief*, 39:107600, 2021. ISSN 2352-3409. doi:10.1016/j.dib.2021.107600.
- [45] Jorge Manjarrez-Sanchez and Gerardo Martinez-Carrillo. Dataset of photosynthesis and photosynthetic factors measurements of greenhouse tomato. *Data in Brief*, 32:106274, 2020. ISSN 2352-3409. doi:10.1016/j.dib.2020.106274.
- [46] Q. Niu, X. Li, J. Huang, H. Huang, X. Huang, W. Su, and W. Yuan. A 30 m annual maize phenology dataset from 1985 to 2020 in china. *Earth System Science Data*, 14(6):2851–2864, 2022. doi:10.5194/essd-14-2851-2022.
- [47] M. Zhu, J. Dai, H. Wang, J. M. Alatalo, W. Liu, Y. Hao, and Q. Ge. Mapping 24 woody plant species phenology and ground forest phenology over china from 1951 to 2020. *Earth System Science Data*, 16(1):277–293, 2024. doi:10.5194/essd-16-277-2024.

- [48] Julia Mergner, Martin Frejno, Maxim Messerer, Daniel Lang, Patroklos Samaras, Mathias Wilhelm, Klaus FX Mayer, Claus Schwechheimer, and Bernhard Kuster. Proteomic and transcriptomic profiling of aerial organ development in arabidopsis. *Scientific Data*, 7(1):334, 2020. doi:10.1038/s41597-020-00678-w.
- [49] Juan C. Castro, J. Dylan Maddox, Hicler N. Rodríguez, Carlos G. Castro, Sixto A. Imán-Correa, Marianela Cobos, Jae D. Paredes, Jorge L. Marapara, Janeth Braga, and Pedro M. Adrianzén. Dataset of de novo assembly and functional annotation of the transcriptome during germination and initial growth of seedlings of myrciaria dubia "camu-camu". *Data in Brief*, 31:105834, 2020. ISSN 2352-3409. doi:10.1016/j.dib.2020.105834.
- [50] Zhenying Peng, Kai-Hua Jia, Jingjing Meng, Jianguo Wang, Jiale Zhang, Xinguo Li, and Shubo Wan. Transcriptome profiling of aerial and subterranean peanut pod development. *Scientific Data*, 11(1):364, 2024. doi:10.1038/s41597-024-03205-3.
- [51] Yufeng Ge, Geng Bai, Vincent Stoerger, and James C. Schnable. Temporal dynamics of maize plant growth, water use, and leaf water content using automated high throughput rgb and hyperspectral imaging. *Computers and Electronics in Agriculture*, 127:625–632, 2016. ISSN 0168-1699. doi:10.1016/j.compag.2016.07.028.
- [52] Mahlet T. Anche, Nicholas Morales, Nicholas S. Kaczmar, Nicholas Santantonio, Michael A. Gore, and Kelly R. Robbins. Scalable growth models for time-series multispectral images. *The Plant Phenome Journal*, 6(1):e20064, 2023. doi:10.1002/ppj2.20064.
- [53] Mahlet T Anche, Nicholas S Kaczmar, Nicolas Morales, James W Clohessy, Daniel C Ilut, Michael A Gore, and Kelly R Robbins. Temporal covariance structure of multi-spectral phenotypes and their predictive ability for end-of-season traits in maize. *Theoretical and Applied Genetics*, 133:2853–2868, 2020. doi:10.1007/s00122-020-03637-6.
- [54] Shangpeng Sun, Changying Li, Andrew H Paterson, Yu Jiang, Rui Xu, Jon S Robertson, John L Snider, and Peng W Chee. In-field high throughput phenotyping and cotton plant growth analysis using lidar. *Frontiers in Plant Science*, 9:16, 2018. doi:10.3389/fpls.2018.00016.
- [55] Diana M. Pérez-Valencia, María Xosé Rodríguez-Álvarez, Martin P. Boer, Lukas Kronenberg, Andreas Hund, Llorenç Cabrera-Bosquet, Emilie J. Millet, and Fred A. van Eeuwijk. A two-stage approach for the spatio-temporal analysis of high-throughput phenotyping data. *Scientific Reports*, 12(1):3177, 2022. ISSN 2045-2322. doi:10.1038/s41598-022-06935-9.
- [56] Soumyashree Kar, Vincent Garin, Jana Kholová, Vincent Vadez, Surya S. Durbha, Ryokei Tanaka, Hiroyoshi Iwata, Milan O. Urban, and J. Adinarayana. Spatemhtp: A data analysis pipeline for efficient processing and utilization of temporal high-throughput phenotyping data. *Frontiers in Plant Science*, 11, 2020. ISSN 1664-462X. doi:10.3389/fpls.2020.552509.
- [57] Mehdi Momen, Malachy T Campbell, Harkamal Walia, and Gota Morota. Predicting Longitudinal Traits Derived from High-Throughput Phenomics in Contrasting Environments Using

- 856 Genomic Legendre Polynomials and B-Splines. *G3 Genes—Genomes—Genetics*, 9(10):3369–
857 3380, 10 2019. ISSN 2160-1836. doi:10.1534/g3.119.400346.
- 858 [58] Duke Pauli, Pedro Andrade-Sanchez, A Elizabete Carmo-Silva, Elodie Gazave, Andrew N
859 French, John Heun, Douglas J Hunsaker, Alexander E Lipka, Tim L Setter, Robert J Strand,
860 Kelly R Thorp, Sam Wang, Jeffrey W White, and Michael A Gore. Field-Based High-
861 Throughput Plant Phenotyping Reveals the Temporal Patterns of Quantitative Trait Loci
862 Associated with Stress-Responsive Traits in Cotton. *G3 Genes—Genomes—Genetics*, 6(4):
863 865–879, 04 2016. ISSN 2160-1836. doi:10.1534/g3.115.023515.
- 864 [59] Toshimi Baba, Mehdi Momen, Malachy T. Campbell, Harkamal Walia, and Gota Morota.
865 Multi-trait random regression models increase genomic prediction accuracy for a temporal
866 physiological trait derived from high-throughput phenotyping. *PLOS ONE*, 15(2):1–17, 02
867 2020. doi:10.1371/journal.pone.0228118.
- 868 [60] Nicolas Morales, Mahlet T Anche, Nicholas S Kaczmar, Nicholas Lepak, Pengzun Ni,
869 Maria Cinta Romay, Nicholas Santantonio, Edward S Buckler, Michael A Gore, Lukas A
870 Mueller, and Kelly R Robbins. Spatio-temporal modeling of high-throughput multispectral
871 aerial images improves agronomic trait genomic prediction in hybrid maize. *Genetics*, 227(1):
872 iyae037, 03 2024. ISSN 1943-2631. doi:10.1093/genetics/iyae037.
- 873 [61] R L Baker, W F Leong, S Welch, and C Weinig. Mapping and Predicting Non-Linear
874 Brassica rapa Growth Phenotypes Based on Bayesian and Frequentist Complex Trait Es-
875 timation. *G3 Genes—Genomes—Genetics*, 8(4):1247–1258, 04 2018. ISSN 2160-1836.
876 doi:10.1534/g3.117.300350.
- 877 [62] Diana Marcela Perez-Valencia, Maria Xose Rodriguez-Alvarez, Martin P Boer, and Fred A
878 van Eeuwijk. A one-stage approach for the spatio-temporal analysis of high-throughput phe-
879 notyping data. *Journal of Agricultural, Biological and Environmental Statistics*, pages 1–23,
880 2024.
- 881 [63] Sergey Nesteruk, Dmitrii Shadrin, Vladislav Kovalenko, Antonio Rodríguez-Sánchez, and
882 Andrey Somov. Plant growth prediction through intelligent embedded sensing. In *2020*
883 *IEEE 29th International Symposium on Industrial Electronics (ISIE)*, pages 411–416, 2020.
884 doi:10.1109/ISIE45063.2020.9152399.
- 885 [64] Alper Adak, Aaron J DeSalvio, Mustafa A Arik, and Seth C Murray. Field-based high-
886 throughput phenotyping enhances phenomic and genomic predictions for grain yield and
887 plant height across years in maize. *G3 Genes—Genomes—Genetics*, 14(7):jkae092, 05 2024.
888 doi:10.1093/g3journal/jkae092.
- 889 [65] K. Matsumura, C. F. Gaitan, K. Sugimoto, A. J. Cannon, and W. W. Hsieh. Maize yield
890 forecasting by linear regression and artificial neural networks in jilin, china. *The Journal of*
891 *Agricultural Science*, 153(3):399–410, 2015. doi:10.1017/S0021859614000392.

- [66] Joon-Woo Lee, Taewon Moon, and Jung-Eek Son. Development of growth estimation algorithms for hydroponic bell peppers using recurrent neural networks. *Horticulturae*, 7(9):284, 2021. doi:10.3390/horticulturae7090284.
- [67] Bashar Alhnaity, Simon Pearson, Georgios Leontidis, and Stefanos Kollias. Using deep learning to predict plant growth and yield in greenhouse environments. In *International Symposium on Advanced Technologies and Management for Innovative Greenhouses: GreenSys2019 1296*, pages 425–432, 2019. doi:10.17660/ActaHortic.2020.1296.55.
- [68] Liyun Gong, Miao Yu, Shouyong Jiang, Vassilis Cutsuridis, and Simon Pearson. Deep learning based prediction on greenhouse crop yield combined tcn and rnn. *Sensors*, 21(13), 2021. ISSN 1424-8220. doi:10.3390/s21134537.
- [69] Bashar Alhnaity, Stefanos Kollias, Georgios Leontidis, Shouyong Jiang, Bert Schamp, and Simon Pearson. An autoencoder wavelet based deep neural network with attention mechanism for multi-step prediction of plant growth. *Information Sciences*, 560:35–50, 2021. ISSN 0020-0255. doi:10.1016/j.ins.2021.01.037.
- [70] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N. Gomez, Lukasz Kaiser, and Illia Polosukhin. Attention is all you need. In *Proceedings of the 31st International Conference on Neural Information Processing Systems, NIPS’17*, pages 6000–6010, Red Hook, NY, USA, 2017. Curran Associates Inc. ISBN 9781510860964. doi:10.48550/arXiv.1706.03762.
- [71] Marie Auger-Méthé, Ken Newman, Diana Cole, Fanny Empacher, Rowenna Gryba, Aaron A. King, Vianey Leos-Barajas, Joanna Mills Flemming, Anders Nielsen, Giovanni Petris, and Len Thomas. A guide to state-space modeling of ecological time series. *Ecological Monographs*, 91(4):e01470, 2021. doi:10.1002/ecm.1470.
- [72] Dmitrii Shadrin, Tatiana Podladchikova, George Ovchinnikov, Artem Pavlov, Mariia Pukalchik, and Andrey Somov. Kalman filtering for accurate and fast plant growth dynamics assessment. In *2020 IEEE International Instrumentation and Measurement Technology Conference (I2MTC)*, pages 1–6, 2020. doi:10.1109/I2MTC43012.2020.9129053.
- [73] Gautier Viaud, Yuting Chen, and Paul-Henry Cournède. Full bayesian inference in hidden markov models of plant growth. *The Annals of Applied Statistics*, 16(4):2352–2368, 2022. doi:10.1214/21-AOAS1594.
- [74] S. Shibata, R. Mizuno, and H. Mineno. Semisupervised deep state-space model for plant growth modeling. *Plant Phenomics*, 2020, 2020. doi:10.34133/2020/4261965.
- [75] Pauliina Schiestl-Aalto, Liisa Kulmala, Harri Mäkinen, Eero Nikinmaa, and Annikki Mäkelä. Cassia – a dynamic model for predicting intra-annual sink demand and interannual growth variation in scots pine. *New Phytologist*, 206(2):647–659, 2015. doi:10.1111/nph.13275.

- [76] Xing-Rong Fan, Meng-Zhen Kang, Ep Heuvelink, Philippe de Reffye, and Bao-Gang Hu. A knowledge-and-data-driven modeling approach for simulating plant growth: A case study on tomato growth. *Ecological Modelling*, 312:363–373, 2015. ISSN 0304-3800. doi:10.1016/j.ecolmodel.2015.06.006.
- [77] Christophe Godin and Hervé Sinoquet. Functional–structural plant modelling. *New Phytologist*, 166(3):705–708, 2005. doi:10.1111/j.1469-8137.2005.01445.x.
- [78] J. Vos, J. B. Evers, G. H. Buck-Sorlin, B. Andrieu, M. Chelle, and P. H. B. de Visser. Functional–structural plant modelling: a new versatile tool in crop science. *Journal of Experimental Botany*, 61(8):2101–2115, 12 2009. ISSN 0022-0957. doi:10.1093/jxb/erp345.
- [79] Jonas R Coussement, Tom De Swaef, Peter Lootens, Isabel Roldán-Ruiz, and Kathy Steppe. Introducing turgor-driven growth dynamics into functional–structural plant models. *Annals of Botany*, 121(5):849–861, 01 2018. ISSN 0305-7364. doi:10.1093/aob/mcx144.
- [80] Jochem B. Evers. *Simulating Crop Growth and Development Using Functional-Structural Plant Modeling*, pages 219–236. Springer Netherlands, Dordrecht, 2016. ISBN 978-94-017-7291-4. doi:10.1007/978-94-017-7291-4_8.
- [81] Xiujuan Wang, Jing Hua, Mengzhen Kang, Haoyu Wang, and Philippe Reffye. Functional–structural plant model ”greenlab”: A state-of-the-art review. *Plant Phenomics*, 6:0118, 2024. doi:10.34133/plantphenomics.0118.
- [82] Yin Hoon Chew, Bénédicte Wenden, Anna Flis, Virginie Mengin, Jasper Taylor, Christopher L. Davey, Christopher Tindal, Howard Thomas, Helen J. Ougham, Philippe de Reffye, Mark Stitt, Mathew Williams, Robert Muetzelfeldt, Karen J. Halliday, and Andrew J. Millar. Multiscale digital *Arabidopsis* predicts individual organ and whole-organism growth. *Proceedings of the National Academy of Sciences*, 111(39):E4127–E4136, 2014. doi:10.1073/pnas.1410238111.
- [83] Sriram Parasurama, Darshi Banan, Kyungdahm Yun, Sharon Doty, and Soo-Hyung Kim. Bridging time-series image phenotyping and functional–structural plant modeling to predict adventitious root system architecture. *Plant Phenomics*, 5:0127, 2023. doi:10.34133/plantphenomics.0127.
- [84] Shogo Nagano, Shogo Moriyuki, Kazumasa Wakamori, Hiroshi Mineno, and Hirokazu Fukuda. Leaf-movement-based growth prediction model using optical flow analysis and machine learning in plant factory. *Frontiers in Plant Science*, 10, 2019. ISSN 1664-462X. doi:10.3389/fpls.2019.00227.
- [85] Changjin Huang, Zilu Wang, David Quinn, Subra Suresh, and K. Jimmy Hsia. Differential growth and shape formation in plant organs. *Proceedings of the National Academy of Sciences*, 115(49):12359–12364, 2018. doi:10.1073/pnas.1811296115.
- [86] Gabriella Mosca, Milad Adibi, Soeren Strauss, Adam Runions, Aleksandra Sapala, and Richard S. Smith. *Modeling Plant Tissue Growth and Cell Division*, pages 107–138. Springer

- International Publishing, Cham, 2018. ISBN 978-3-319-99070-5. doi:10.1007/978-3-319-99070-5_7.
- [87] Daniel Kierzkowski, Adam Runions, Francesco Vuolo, Sören Strauss, Rena Lymbouridou, Anne-Lise Routier-Kierzkowska, David Wilson-Sánchez, Hannah Jenke, Carla Galinha, Gabriella Mosca, Zhongjuan Zhang, Claudia Canales, Raffaele Dello Ioio, Peter Huijser, Richard S. Smith, and Miltos Tsiantis. A growth-based framework for leaf shape development and diversity. *Cell*, 177(6):1405–1418.e17, 2019. ISSN 0092-8674. doi:10.1016/j.cell.2019.05.011.
- [88] Przemyslaw Prusinkiewicz and Aristid Lindenmayer. *The algorithmic beauty of plants*. Springer Science & Business Media, 2012.
- [89] Przemyslaw Prusinkiewicz, Mikolaj Cieslak, Pascal Ferraro, and Jim Hanan. *Modeling Plant Development with L-Systems*, pages 139–169. Springer International Publishing, Cham, 2018. ISBN 978-3-319-99070-5. doi:10.1007/978-3-319-99070-5_8.
- [90] M. T. Allen, P. Prusinkiewicz, and T. M. DeJong. Using l-systems for modeling source–sink interactions, architecture and physiology of growing trees: the l-peach model. *New Phytologist*, 166(3):869–880, 2005. doi:10.1111/j.1469-8137.2005.01348.x.
- [91] TM DeJong, D Da Silva, C Negron, M Cieslak, and P Prusinkiewicz. The l-almond model: a functional-structural virtual tree model of almond tree architectural growth, carbohydrate dynamics over multiple years. In *X International Symposium on Modelling in Fruit Research and Orchard Management 1160*, pages 43–50, 2015. doi:10.17660/ActaHortic.2017.1160.7.
- [92] Theodore DeJong. Simulating fruit tree growth, structure, and physiology using l-systems. *Crop Science*, 62(6):2091–2106, 2022. doi:10.1002/csc2.20847.
- [93] Rika Rokhana, Wiwiet Herulambang, and Rarasmaya Indraswari. Machine learning and polynomial – l system algorithm for modeling and simulation of glycine max (l) merrill growth. In *2020 International Electronics Symposium (IES)*, pages 463–467, 2020. doi:10.1109/IES50839.2020.9231935.
- [94] Le Yang, Panpan Wu, Zhengkang Zuo, Lan Long, Junlin Shi, and Yutang Liu. Eroots: A three-dimensional dynamic growth model of rice roots coupled with soil. *Biosystems Engineering*, 244:122–133, 2024. ISSN 1537-5110. doi:10.1016/j.biosystemseng.2024.06.002.
- [95] Yasmeeen Hitti, Ionelia Buzatu, Manuel Del Verme, Mark Lefsrud, Florian Golemo, and Audrey Durand. Growspace: A reinforcement learning environment for plant architecture. *Computers and Electronics in Agriculture*, 217:108613, 2024. ISSN 0168-1699. doi:10.1016/j.compag.2024.108613.
- [96] Marc Mußmann, Daniel Nicolas Hofstadler, and Sebastian von Mammen. An agent-based, interactive simulation model of root growth. volume ALIFE 2024: Proceedings of the 2024 Artificial Life Conference of *Artificial Life Conference Proceedings*, page 7, 07 2024. doi:10.1162/isal.a.00718.

- [97] Shunsuke Sakurai., Hideaki Uchiyama., Atshushi Shimada., and Rin-Ichiro Taniguchi. Plant growth prediction using convolutional lstm. In *Proceedings of the 14th International Joint Conference on Computer Vision, Imaging and Computer Graphics Theory and Applications (5: VISAPP)*, pages 105–113. INSTICC, 2019. ISBN 978-989-758-354-4. doi:10.5220/0007404901050113.
- [98] Chunying Wang, Weiting Pan, Xubin Song, Haixia Yu, Junke Zhu, Ping Liu, and Xiang Li. Predicting plant growth and development using time-series images. *Agronomy*, 12(9), 2022. ISSN 2073-4395. doi:10.3390/agronomy12092213.
- [99] Diederik P Kingma and Max Welling. Auto-encoding variational bayes. *arXiv preprint arXiv:1312.6114*, 2013. doi:10.48550/arXiv.1312.6114.
- [100] Robail Yasrab, Jincheng Zhang, Polina Smyth, and Michael P. Pound. Predicting plant growth from time-series data using deep learning. *Remote Sensing*, 13(3), 2021. ISSN 2072-4292. doi:10.3390/rs13030331.
- [101] Taehyeon Kim, Sang-Ho Lee, and Jong-Ok Kim. A novel shape based plant growth prediction algorithm using deep learning and spatial transformation. *IEEE Access*, 10:37731–37742, 2022. doi:10.1109/ACCESS.2022.3165211.
- [102] Yinglun Li, Xiaohai Zhan, Shouyang Liu, Hao Lu, Ruibo Jiang, Wei Guo, Scott Chapman, Yufeng Ge, Benoit Solan, Yanfeng Ding, and Frédéric Baret. Self-supervised plant phenotyping by combining domain adaptation with 3d plant model simulations: Application to wheat leaf counting at seedling stage. *Plant Phenomics*, 5:0041, 2023. doi:10.34133/plantphenomics.0041.
- [103] Yuzhen Lu, Dong Chen, Ebenezer Olaniyi, and Yanbo Huang. Generative adversarial networks (gans) for image augmentation in agriculture: A systematic review. *Computers and Electronics in Agriculture*, 200:107208, 2022. ISSN 0168-1699. doi:10.1016/j.compag.2022.107208.
- [104] Miro Miranda, Lukas Drees, and Ribana Roscher. Controlled multi-modal image generation for plant growth modeling. In *2022 26th International Conference on Pattern Recognition (ICPR)*, pages 5118–5124, 2022. doi:10.1109/ICPR56361.2022.9956115.
- [105] Lukas Drees, Laura Verena Junker-Frohn, Jana Kierdorf, and Ribana Roscher. Temporal prediction and evaluation of brassica growth in the field using conditional generative adversarial networks. *Computers and Electronics in Agriculture*, 190:106415, 2021. ISSN 0168-1699. doi:10.1016/j.compag.2021.106415.
- [106] Johannes Leonhardt, Lukas Drees, Peter Jung, and Ribana Roscher. Probabilistic biomass estimation with conditional generative adversarial networks. In Björn Andres, Florian Bernard, Daniel Cremers, Simone Frintrop, Bastian Goldlücke, and Ivo Ihrke, editors, *Pattern Recognition*, pages 479–494, Cham, 2022. Springer International Publishing. ISBN 978-3-031-16788-1.
- [107] Lukas Drees, Dereje T. Demie, Madhuri R. Paul, Johannes Leonhardt, Sabine J. Seidel, Thomas F. Döring, and Ribana Roscher. Data-driven crop growth simulation on time-varying

- 1036 generated images using multi-conditional generative adversarial networks. *Plant Methods*, 20
1037 (1), 06 2024. ISSN 1746-4811. doi:10.1186/s13007-024-01205-3.
- 1038 [108] Mohamed Debbagh, Yixue Liu, Zhouzhou Zheng, Xintong Jiang, Shangpeng Sun, and Mark
1039 Lefsrud. Generative plant growth simulation from sequence-informed environmental condi-
1040 tions. In *Artificial Neural Networks in Pattern Recognition*, pages 308–319, Cham, 2024.
1041 Springer Nature Switzerland. ISBN 978-3-031-71602-7. doi:10.1007/978-3-031-71602-7_26.
- 1042 [109] Ribana Roscher, Lukas Roth, Cyrill Stachniss, and Achim Walter. Data-centric digital agri-
1043 culture: A perspective, 2023.