# GATE DA 2027

## Data Science & Artificial Intelligence

---

# Calculus and Optimization

### 100 High-Yield Master Questions targeting AIR 1

---

**Features of this Compendium:**

- **100 Unique Questions**: 35 Verified GATE PYQs (DA, CS, MA, EC, ME), 25 Advanced Conceptual Probes, 20 Expected GATE DA 2027 Questions, 10 High-Difficulty Numerical/Application Items, and 10 Mixed-Concept Synthesis Problems.
- **Strict GATE Typologies**: Multiple Choice Questions (MCQ with negative marking rules), Multiple Select Questions (MSQ), and Numerical Answer Type (NAT).
- **Calibrated Difficulty**: 20% Easy, 50% Moderate, 30% Difficult (AIR 1 Tier).
- **Complete Rigor**: Full answer key, step-by-step derivations, traps/common pitfalls, and an end-to-end Rapid Revision Formula Sheet.

**Target Year: GATE 2027** · Paper Code: **DA**

Comprehensive Self-Contained Study and Practice Material

# Contents


**1 Master Question Bank (Questions 1–100) 2**

1.1 Limits, Continuity, and Differentiability . . . 2
1.2 Single-Variable Derivatives, Mean Value Theorems, and Series . . . 3
1.3 Definite Integrals and Applications . . . 4
1.4 Multivariable Calculus: Partial Derivatives, Chain Rule, and Directional Derivatives 5
1.5 Vector Calculus: Gradient, Divergence, and Vector Fields . . . 7
1.6 Convex Sets and Convex/Concave Functions . . . 8
1.7 Hessian Matrix and Second-Order Optimality Conditions . . . 9
1.8 Unconstrained Optimization of Multivariable Functions . . . 10
1.9 Constrained Optimization: Lagrange Multipliers and KKT Conditions . . . 11
1.10 Mixed Calculus and Optimization Problems (Questions 63–75) . . . 12
1.11 Advanced Applications, Constrained Systems, and Deep Conceptuals (Questions 76–100) . . . 14

**2 Master Answer Key (1–100) 19**

**3 Comprehensive Solutions with Concept Notes & Traps 20**

**4 GATE DA 2027 Rapid Revision Sheet: Calculus & Optimization 32**

4.1 Essential Calculus Identities . . . 32
4.2 Convexity and Optimization Theory . . . 32

# 1 Master Question Bank (Questions 1–100)

## 1.1 Limits, Continuity, and Differentiability

**Q1.** MCQ [Negative: 0.33] 1 Mark GATE PYQ GATE CS 2016

Compute the limit:

$$L = \lim_{x \to 0} \frac{e^x - (1 + x + x^2/2)}{x^3}$$

(A) $0$

(B) $1/6$

(C) $1/3$

(D) $1/2$

**Q2.** MCQ [Negative: 0.66] 2 Marks GATE PYQ GATE DA 2024, Q14

Consider the function $f(x) = |x|^3$ defined on $\mathbb{R}$. At $x = 0$, which of the following statements is true?

(A) $f(x)$ is continuous but not differentiable.

(B) $f'(0)$ exists, but $f''(0)$ does not exist.

(C) $f''(0)$ exists, but $f'''(0)$ does not exist.

(D) $f(x)$ is infinitely differentiable on $\mathbb{R}$.

**Q3.** NAT 1 Mark GATE PYQ GATE CS 2015

Evaluate the value of:

$$\lim_{x \to 0} \frac{\sin x - x + \frac{x^3}{6}}{x^5}$$

(Answer in decimal format rounded to three decimal places).

**Q4.** MSQ 2 Marks Advanced Conceptual

Let $f : \mathbb{R} \to \mathbb{R}$ be defined as:

$$f(x) = \begin{cases} x^2 \sin(1/x), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Which of the following statements is/are **CORRECT**?

(A) $f$ is differentiable at $x = 0$ with $f'(0) = 0$.

(B) $f'(x)$ is continuous at $x = 0$.

(C) $f'(x)$ is bounded on $[-1, 1]$.

(D) $f''(0)$ exists.

**Q5.** NAT 2 Marks Difficult Numerical

Evaluate the limit:

$$L = \lim_{x \to 0} \frac{\ln(\cos x) + \frac{1}{2}x^2}{x^4}$$

(Round off to two decimal places).

**Q6.** MCQ [Negative: 0.66] 2 Marks GATE PYQ GATE MA 2018

Let $f(x) = \lim_{n\to\infty} \frac{x^{2n}-1}{x^{2n}+1}$ for $x \in \mathbb{R}$. The set of points where $f(x)$ is discontinuous is:

(A) $\{0\}$

(B) $\{-1, 1\}$

(C) $\{-1, 0, 1\}$

(D) Empty set $\emptyset$

**Q7.** MSQ 2 Marks Expected GATE DA 2027

Consider $f(x) = \max\{1 - x, 1 + x, 2\}$. Which of the following statements is/are **TRUE**?

(A) $f(x)$ is continuous everywhere on $\mathbb{R}$.

(B) $f(x)$ is non-differentiable at exactly two points.

(C) $f(x)$ is a convex function on $\mathbb{R}$.

(D) The global minimum value of $f(x)$ is $2$.

**Q8.** NAT 2 Marks Advanced Conceptual

Let $f(x) = x\lfloor x \rfloor$ for $x \in [0, 3]$, where $\lfloor \cdot \rfloor$ denotes the greatest integer function. Find the number of points in $(0, 3)$ where $f(x)$ is discontinuous.

## 1.2 Single-Variable Derivatives, Mean Value Theorems, and Series

**Q9.** MCQ [Negative: 0.66] 2 Marks GATE PYQ GATE CS 2017

The value of $c \in (1, 2)$ satisfying the conclusion of Lagrange's Mean Value Theorem for $f(x) = \ln x$ on $[1, 2]$ is:

(A) $\frac{1}{\ln 2}$

(B) $\ln 2$

(C) $e$

(D) $\frac{1}{2}$

**Q10.** NAT 2 Marks GATE PYQ GATE DA 2024, Q28

In the Taylor series expansion of $f(x) = \frac{1}{1-x}$ about $x = 2$, find the coefficient of $(x - 2)^3$.

**Q11.** MSQ 2 Marks Advanced Conceptual

Let $f : [a, b] \to \mathbb{R}$ be continuous on $[a, b]$ and twice differentiable on $(a, b)$. If the line segment connecting $(a, f(a))$ and $(b, f(b))$ intersects the graph of $f$ at a point $c \in (a, b)$, which of the following is/are guaranteed?

(A) There exists $\xi \in (a, b)$ such that $f'(\xi) = \frac{f(b)-f(a)}{b-a}$.

(B) There exists $\eta \in (a, b)$ such that $f''(\eta) = 0$.

(C) $f'(x)$ must have a zero in $(a, b)$.

(D) $f(x)$ is monotonic on $[a, b]$.

**Q12.** NAT | 1 Mark | GATE PYQ | GATE CS 2019

The coefficient of $x^4$ in the Maclaurin series expansion of $f(x) = e^{x^2}$ is: (Provide exact fraction or decimal).

**Q13.** MCQ [Negative: 0.33] | 1 Mark | GATE PYQ | GATE ME 2014

The radius of convergence of the power series $\sum_{n=1}^{\infty} \frac{x^n}{n \cdot 2^n}$ is:

(A) 0

(B) 1

(C) 2

(D) $\infty$

**Q14.** NAT | 2 Marks | Difficult Numerical

Let $f(x) = \cos(x^2)$. Compute the value of the 10th derivative of $f$ evaluated at 0, i.e., $f^{(10)}(0)$.

**Q15.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE EC 2018

Consider the function $f(x) = x^3 - 3x^2 + 3x + 7$. Which of the following describes the point $x = 1$?

(A) Local maximum

(B) Local minimum

(C) Inflection point with horizontal tangent

(D) Discontinuity

**Q16.** MSQ | 2 Marks | Expected GATE DA 2027

Let $g(x) = \frac{\sin x}{x}$ for $x > 0$. Which of the following statements is/are **TRUE**?

(A) $g(x)$ is strictly decreasing on $(0, \pi/2)$.

(B) $g(x) < 1$ for all $x \in (0, \pi/2)$.

(C) $\tan x > x$ for all $x \in (0, \pi/2)$.

(D) $g(x)$ is convex on $(0, \pi/2)$.

## 1.3 Definite Integrals and Applications

**Q17.** NAT | 2 Marks | GATE PYQ | GATE DA 2024, Q31

Evaluate the definite integral:

$$I = \int_0^1 \int_0^1 (x + y)\, dx\, dy$$

**Q18.** NAT | 2 Marks | GATE PYQ | GATE CS 2014

Compute the integral:

$$I = \int_{-2}^{2} \frac{x^3 \cos x}{1 + x^2}\, dx + \int_{-2}^{2} \sqrt{4 - x^2}\, dx$$

(Use $\pi \approx 3.1416$, round to two decimal places).

**Q19.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE MA 2020

Let $F(x) = \int_0^{x^2} \exp(-t^2)\, dt$. The derivative $F'(x)$ at $x = 1$ equals:

(A) $e^{-1}$

(B) $2e^{-1}$

(C) $2e^{-2}$

(D) $e^{-2}$

**Q20.** MSQ | 2 Marks | Advanced Conceptual

Consider $I_n = \int_0^1 (1-x^2)^n \, dx$ for $n \in \mathbb{N}$. Which of the following is/are **CORRECT**?

(A) $I_n = \frac{2n}{2n+1} I_{n-1}$ for $n \geq 1$.

(B) The sequence $\{I_n\}$ is monotonically decreasing.

(C) $\lim_{n\to\infty} I_n = 0$.

(D) $\int_0^\infty e^{-x^2} dx = \lim_{n\to\infty} \sqrt{n} I_n$.

**Q21.** NAT | 2 Marks | Expected GATE DA 2027

Evaluate:

$$\int_0^\infty x^3 e^{-2x} \, dx$$

(Answer in decimal format).

**Q22.** MCQ [Negative: 0.33] | 1 Mark | GATE PYQ | GATE ME 2017

The value of $\int_0^{\pi/2} \sin^5 x \cos^3 x \, dx$ is:

(A) $1/24$

(B) $1/12$

(C) $1/48$

(D) $1/6$

**Q23.** NAT | 2 Marks | Difficult Numerical

Compute the value of the double integral $\iint_D e^{x^2+y^2} \, dx \, dy$, where $D = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1, x \geq 0, y \geq 0\}$. Express answer in terms of $\pi$ and $e$ (or round to two decimal places).

## 1.4 Multivariable Calculus: Partial Derivatives, Chain Rule, and Directional Derivatives

**Q24.** NAT | 1 Mark | GATE PYQ | GATE CS 2018

Let $f(x, y) = x^2 y + y^3 x$. Find the value of $\frac{\partial^2 f}{\partial x \partial y}$ at $(1, 2)$.

**Q25.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE DA 2024, Q41

Let $f(x, y) = 2x^2 + 3y^2$. The directional derivative of $f$ at the point $(1, 1)$ in the direction of the vector $\mathbf{v} = (3, 4)$ is:

(A) $36/5$

(B) $24/5$

(C) $12$

(D) $7$

**Q26.** **MSQ** 2 Marks Advanced Conceptual

Let $f(x,y) = \begin{cases} \frac{x^3y}{x^4+y^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$. Which of the following is/are **TRUE**?

(A) Both partial derivatives $f_x(0,0)$ and $f_y(0,0)$ exist and equal 0.

(B) Directional derivatives exist along all directions at $(0,0)$.

(C) $f(x,y)$ is continuous at $(0,0)$.

(D) $f(x,y)$ is differentiable at $(0,0)$.

**Q27.** **NAT** 2 Marks Expected GATE DA 2027

Let $z = f(u,v)$ where $u = x^2 - y^2$ and $v = 2xy$. If $f$ is continuously differentiable and at $(u,v) = (3,4)$, $\frac{\partial f}{\partial u} = 2$ and $\frac{\partial f}{\partial v} = -1$. Given that $(x,y) = (2,1)$ maps to $(u,v) = (3,4)$, compute $\frac{\partial z}{\partial x}$ at $(2,1)$.

**Q28.** **MCQ [Negative: 0.66]** 2 Marks GATE PYQ GATE MA 2021

The unit vector in which the function $f(x,y,z) = x^2yz^3$ increases most rapidly at $(1,1,1)$ is:

(A) $\frac{1}{\sqrt{14}}(2,1,3)$

(B) $\frac{1}{\sqrt{3}}(1,1,1)$

(C) $\frac{1}{\sqrt{14}}(1,2,3)$

(D) $\frac{1}{\sqrt{6}}(2,1,1)$

**Q29.** **MSQ** 2 Marks Advanced Conceptual

Let $f : \mathbb{R}^n \to \mathbb{R}$ be continuously differentiable. Which of the following statements about directional derivatives is/are **TRUE**?

(A) $D_{\mathbf{u}} f(\mathbf{x}) = \nabla f(\mathbf{x}) \cdot \mathbf{u}$ for any unit vector $\mathbf{u}$.

(B) The maximum directional derivative is $\|\nabla f(\mathbf{x})\|_2$.

(C) The directional derivative in a direction orthogonal to the gradient vector is zero.

(D) $\nabla f(\mathbf{x})$ is tangent to the level set of $f$ passing through $\mathbf{x}$.

**Q30.** **NAT** 2 Marks Difficult Numerical

Let $f(x,y) = x^3 - 3xy^2 + y$. Find the maximum rate of change of $f$ at the point $(1,-1)$. (Round off to two decimal places).

## 1.5 Vector Calculus: Gradient, Divergence, and Vector Fields

**Q31.** MCQ [Negative: 0.33] 1 Mark GATE PYQ GATE EC 2017

The divergence of the vector field $\mathbf{F}(x, y, z) = x^2\mathbf{i} + 2y^2\mathbf{j} + 3z^2\mathbf{k}$ at $(1, 1, 1)$ is:

(A) 6

(B) 12

(C) 14

(D) 0

**Q32.** MCQ [Negative: 0.66] 2 Marks GATE PYQ GATE ME 2016

If $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and $r = \|\mathbf{r}\|$, then $\nabla\left(\frac{1}{r}\right)$ is:

(A) $\frac{\mathbf{r}}{r^3}$

(B) $-\frac{\mathbf{r}}{r^3}$

(C) $\frac{\mathbf{r}}{r^2}$

(D) $-\frac{\mathbf{r}}{r^2}$

**Q33.** MSQ 2 Marks Advanced Conceptual

Let $\mathbf{F} = \nabla\phi$ be a conservative vector field with scalar potential $\phi \in C^2(\mathbb{R}^3)$. Which of the following is/are **ALWAYS TRUE**?

(A) $\nabla \times \mathbf{F} = \mathbf{0}$.

(B) $\text{div}(\mathbf{F}) = \nabla^2\phi$.

(C) The line integral $\int_C \mathbf{F} \cdot d\mathbf{r}$ is zero over every closed path $C$.

(D) If $\text{div}(\mathbf{F}) = 0$, then $\phi$ is a harmonic function.

**Q34.** NAT 2 Marks Expected GATE DA 2027

Let $\mathbf{F}(x, y) = (2xy + y^2)\mathbf{i} + (x^2 + 2xy)\mathbf{j}$. Find the value of $\int_{(0,0)}^{(1,2)} \mathbf{F} \cdot d\mathbf{r}$.

**Q35.** MCQ [Negative: 0.66] 2 Marks GATE PYQ GATE CS 2015

The Laplacian $\nabla^2 f$ of $f(x, y) = \ln(x^2 + y^2)$ for $(x, y) \neq (0, 0)$ is:

(A) $\frac{2}{x^2+y^2}$

(B) 0

(C) $\frac{4}{x^2+y^2}$

(D) $\frac{-2}{x^2+y^2}$

## 1.6 Convex Sets and Convex/Concave Functions

**Q36.** MSQ | 2 Marks | Expected GATE DA 2027

Which of the following subsets of $\mathbb{R}^2$ is/are **CONVEX**?

(A) $S_1 = \{(x, y) \in \mathbb{R}^2 : x^2 + 2y^2 \leq 4\}$

(B) $S_2 = \{(x, y) \in \mathbb{R}^2 : y \geq x^2\}$

(C) $S_3 = \{(x, y) \in \mathbb{R}^2 : xy \geq 1, x > 0, y > 0\}$

(D) $S_4 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$

**Q37.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE DA 2024, Q22

Consider the function $f : \mathbb{R}^n \to \mathbb{R}$ defined by $f(\mathbf{x}) = \|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^{n} x_i^2}$. Which of the following is true?

(A) $f(\mathbf{x})$ is concave.

(B) $f(\mathbf{x})$ is convex.

(C) $f(\mathbf{x})$ is neither convex nor concave.

(D) $f(\mathbf{x})$ is strictly convex on $\mathbb{R}^n$.

**Q38.** MSQ | 2 Marks | Advanced Conceptual

Let $f, g : \mathbb{R}^n \to \mathbb{R}$ be convex functions. Which of the following functions is/are **ALWAYS CONVEX**?

(A) $h(\mathbf{x}) = \max\{f(\mathbf{x}), g(\mathbf{x})\}$

(B) $h(\mathbf{x}) = f(\mathbf{x}) + g(\mathbf{x})$

(C) $h(\mathbf{x}) = f(\mathbf{x}) \cdot g(\mathbf{x})$

(D) $h(\mathbf{x}) = \alpha f(\mathbf{x}) + \beta g(\mathbf{x})$ for all $\alpha, \beta \in \mathbb{R}$

**Q39.** MCQ [Negative: 0.66] | 2 Marks | Expected GATE DA 2027

Let $f(x) = -\ln(x)$ on $(0, \infty)$. Then:

(A) $f(x)$ is strictly concave.

(B) $f(x)$ is strictly convex.

(C) $f(x)$ is affine.

(D) $f(x)$ is neither convex nor concave.

**Q40.** NAT | 2 Marks | Difficult Numerical

Let $f(x, y) = x^2 + 2cxy + 4y^2$. Find the largest integer value of $c$ for which $f(x, y)$ is strictly convex on $\mathbb{R}^2$.

**Q41.** MSQ | 2 Marks | Advanced Conceptual

Let $f : \mathbb{R}^n \to \mathbb{R}$ be a differentiable function. Which of the following statements is/are equivalent to the convexity of $f$?

(A) $f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x})$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$.

(B) $(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}))^\top (\mathbf{y} - \mathbf{x}) \geq 0$ for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ (monotonicity of the gradient).

(C) The epigraph $\text{epi}(f) = \{(\mathbf{x}, t) : f(\mathbf{x}) \leq t\}$ is a convex set.

(D) Every local minimum of $f$ is a strict global minimum.

**Q42.** **MCQ [Negative: 0.33]** 1 Mark GATE PYQ GATE CS 2019

A function $f : \mathbb{R} \to \mathbb{R}$ is strictly convex and twice differentiable everywhere. How many global minima can $f$ have?

(A) Exactly one

(B) At most one

(C) At least one

(D) Infinitely many

## 1.7 Hessian Matrix and Second-Order Optimality Conditions

**Q43.** **NAT** 2 Marks GATE PYQ GATE DA 2024, Q52

Let $f(x, y) = x^3 + y^3 - 3xy$. Compute the determinant of the Hessian matrix of $f$ evaluated at $(1, 1)$.

**Q44.** **MCQ [Negative: 0.66]** 2 Marks Expected GATE DA 2027

Let $f(x, y) = x^4 + y^4 - 4xy + 1$. The point $(0, 0)$ is:

(A) A local minimum

(B) A local maximum

(C) A saddle point

(D) A point with positive definite Hessian

**Q45.** **MSQ** 2 Marks Advanced Conceptual

Let $H \in \mathbb{R}^{n \times n}$ be the Hessian matrix of a twice continuously differentiable function $f$ at a critical point $\mathbf{x}^*$ (where $\nabla f(\mathbf{x}^*) = \mathbf{0}$). Which of the following is/are **TRUE**?

(A) If $H \succ 0$ (strictly positive definite), then $\mathbf{x}^*$ is a strict local minimum.

(B) If $H \prec 0$ (strictly negative definite), then $\mathbf{x}^*$ is a strict local maximum.

(C) If $H$ has both positive and negative eigenvalues, $\mathbf{x}^*$ is a saddle point.

(D) If $\det(H) = 0$, $\mathbf{x}^*$ cannot be an extremum.

**Q46.** **NAT** 2 Marks Difficult Numerical

Consider the quadratic function $f(x_1, x_2, x_3) = x_1^2 + 2x_2^2 + 5x_3^2 + 2x_1x_2 - 4x_2x_3$. Find the smallest eigenvalue of the Hessian matrix of $f$.

**Q47.** **MCQ [Negative: 0.66]** 2 Marks GATE PYQ GATE CS 2020

Let $f(x, y) = xy - x - y + 1$. The critical point $(1, 1)$ is:

(A) A local maximum

(B) A local minimum

(C) A saddle point

(D) Non-isolated extremum

**Q48.** **MSQ** 2 Marks Expected GATE DA 2027

For $f(x, y) = x^2y - 2xy + y^3$, which of the following critical points exist and have the correct classification?

(A) $(0, 0)$ is a critical point.

(B) $(2, 0)$ is a critical point.

(C) $(1, 1/\sqrt{3})$ is a critical point.

(D) $(0, 0)$ is a strict local minimum.

## 1.8 Unconstrained Optimization of Multivariable Functions

**Q49.** **NAT** 2 Marks GATE PYQ GATE DA 2024, Q56

Find the global minimum value of the function $f(x, y) = (x - 1)^2 + (y - 2)^2 + 5$ on $\mathbb{R}^2$.

**Q50.** **MCQ [Negative: 0.66]** 2 Marks GATE PYQ GATE MA 2019

Let $f(x, y) = x^2 + 4xy + 5y^2 - 2x - 4y$. The unique unconstrained stationary point of $f$ is:

(A) $(3, -1)$

(B) $(-3, 1)$

(C) $(1, 0)$

(D) $(0, 1)$

**Q51.** **MSQ** 2 Marks Advanced Conceptual

Consider gradient descent with fixed step size $\eta > 0$ applied to $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$, where $A$ is symmetric positive definite with eigenvalues $0 < \lambda_{\text{min}} \leq \lambda_{\text{max}}$. Which of the following is/are **TRUE**?

(A) The algorithm converges to the unique minimizer if and only if $0 < \eta < \frac{2}{\lambda_{\text{max}}}$.

(B) The optimal constant learning rate that maximizes convergence rate is $\eta^* = \frac{2}{\lambda_{\text{min}}+\lambda_{\text{max}}}$.

(C) If the condition number $\kappa = \lambda_{\text{max}}/\lambda_{\text{min}} = 1$, convergence occurs in a single iteration from any starting point if $\eta = 1/\lambda_{\text{max}}$.

(D) For any convex function, gradient descent is always guaranteed to converge for any $\eta > 0$.

**Q52.** **NAT** 2 Marks Difficult Numerical

Let $f(x, y) = 100(y - x^2)^2 + (1 - x)^2$ (Rosenbrock function). At the point $(2, 4)$, compute the Euclidean norm of the gradient vector, $\|\nabla f(2, 4)\|_2$.

**Q53.** **MCQ [Negative: 0.66]** 2 Marks Expected GATE DA 2027

Newton's method for unconstrained optimization updates iterates via $\mathbf{x}_{k+1} = \mathbf{x}_k - [H(\mathbf{x}_k)]^{-1}\nabla f(\mathbf{x}_k)$. Applied to a strictly convex quadratic function $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} + \mathbf{b}^\top\mathbf{x} + c$, Newton's method will:

(A) Require exactly $n$ steps where $n$ is dimension.

(B) Converge in exactly 1 step from any initial point $\mathbf{x}_0$.

(C) Converge linearly.

(D) Fail if the Hessian is not diagonal.

**Q54.** NAT | 2 Marks | Advanced Conceptual

Find the number of critical points of $f(x, y) = (x^2 - 1)^2 + (y^2 - 1)^2$ on $\mathbb{R}^2$.

## 1.9 Constrained Optimization: Lagrange Multipliers and KKT Conditions

**Q55.** NAT | 2 Marks | GATE PYQ | GATE DA 2024, Q59

Minimize $f(x, y) = x^2 + y^2$ subject to $x + y = 4$. What is the optimal value of $f$?

**Q56.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE CS 2021

The maximum value of $f(x, y) = xy$ subject to the constraint $x^2 + y^2 = 8$ with $x, y \geq 0$ is:

(A) 2

(B) 4

(C) 8

(D) 16

**Q57.** MSQ | 2 Marks | Advanced Conceptual

Consider the constrained optimization problem:

$$\min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \quad \text{subject to } g_i(\mathbf{x}) \leq 0,\ i = 1, \ldots, m$$

where $f$ and $g_i$ are convex and differentiable. If $\mathbf{x}^*$ and $\boldsymbol{\lambda}^*$ satisfy the Karush-Kuhn-Tucker (KKT) conditions:

(A) $\mathbf{x}^*$ is guaranteed to be a global minimum.

(B) $\lambda_i^* \geq 0$ for all $i = 1, \ldots, m$.

(C) Complementary slackness requires $\lambda_i^* g_i(\mathbf{x}^*) = 0$ for all $i$.

(D) If $g_i(\mathbf{x}^*) < 0$, then $\lambda_i^* > 0$.

**Q58.** NAT | 2 Marks | Expected GATE DA 2027

Consider the problem:

$$\min_{x,y} (x - 2)^2 + (y - 1)^2 \quad \text{subject to } x + y \leq 1,\ x \geq 0,\ y \geq 0$$

Find the optimal value of the Lagrange multiplier $\lambda$ associated with the constraint $x + y - 1 \leq 0$.

**Q59.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE MA 2022

The minimum distance from the origin to the plane $2x - y + 2z = 9$ is:

(A) 1

(B) 2

(C) 3

(D) 9

**Q60.** MSQ | 2 Marks | Expected GATE DA 2027

Consider $\max_{x,y} x+2y$ subject to $x^2+y^2 \leq 5$. Which of the following statements is/are **CORRECT**?

(A) The optimal solution lies on the boundary $x^2 + y^2 = 5$.

(B) The Lagrange multiplier $\lambda$ is strictly positive.

(C) The optimal point is $(1, 2)$.

(D) The maximum value of the objective function is 5.

**Q61.** NAT | 2 Marks | Difficult Numerical

Find the minimum value of $f(x, y, z) = x^2 + 2y^2 + 3z^2$ subject to $x + y + z = 11$.

**Q62.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE ME 2018

Using the method of Lagrange multipliers, the stationary points of $f(x, y) = x^2 + y^2$ subject to $xy = 1$ occur at:

(A) $(1, 1)$ and $(-1, -1)$

(B) $(1, -1)$ and $(-1, 1)$

(C) $(2, 1/2)$ and $(-2, -1/2)$

(D) $(0, 0)$

## 1.10 Mixed Calculus and Optimization Problems (Questions 63–75)

**Q63.** MSQ | 2 Marks | Mixed Concept

Let $f(x) = \int_0^x (t-1)(t-2)^2\, dt$. Which of the following is/are **CORRECT**?

(A) $x = 1$ is a local minimum of $f$.

(B) $x = 2$ is a local maximum of $f$.

(C) $x = 2$ is an inflection point of $f$.

(D) $f$ is strictly convex on $(2, \infty)$.

**Q64.** NAT | 2 Marks | Mixed Concept

Compute the minimum value attained by the function $g(t) = \int_0^1 (x - t)^2\, dx$ for $t \in \mathbb{R}$. (Round off to three decimal places).

**Q65.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE CS 2013

The value of $\lim_{n\to\infty} \sum_{k=1}^{n} \frac{n}{n^2+k^2}$ is:

(A) $\pi/4$

(B) $\pi/2$

(C) $\ln 2$

(D) 1

**Q66.** NAT | 2 Marks | Mixed Concept

Evaluate the limit of Riemann sums:

$$\lim_{n\to\infty} \frac{1}{n^4} \sum_{k=1}^{n} k^3$$

**Q67.** MSQ | 2 Marks | Mixed Concept

Let $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$, where $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ and $\mathbf{b} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}$. Which of the following is/are **TRUE**?

(A) $A$ is symmetric and positive definite.

(B) $f(\mathbf{x})$ is strictly convex on $\mathbb{R}^2$.

(C) The unique global minimizer is $\mathbf{x}^* = (1, 1)^\top$.

(D) The minimum value is $-3$.

**Q68.** NAT | 2 Marks | Difficult Numerical

Let $f(x, y) = e^{-(x^2+y^2)}(2x^2 + 3y^2)$. Find the maximum value of $f(x, y)$ on $\mathbb{R}^2$. (Round to two decimal places).

**Q69.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE EC 2019

The directional derivative of $f(x, y, z) = 2x^2 + 3y^2 + z^2$ at the point $P(2, 1, 3)$ in the direction of the vector $\mathbf{a} = \mathbf{i} - 2\mathbf{k}$ is:

(A) $-4/\sqrt{5}$

(B) $4/\sqrt{5}$

(C) $2/\sqrt{5}$

(D) $-2/\sqrt{5}$

**Q70.** MSQ | 2 Marks | Expected GATE DA 2027

Consider $f(x) = x \ln x$ for $x > 0$. Which of the following statements is/are **CORRECT**?

(A) $f(x)$ is strictly convex on $(0, \infty)$.

(B) The global minimum of $f(x)$ is achieved at $x = 1/e$.

(C) The minimum value of $f(x)$ is $-1/e$.

(D) $f(x)$ has an inflection point at $x = 1$.

**Q71.** NAT | 2 Marks | Mixed Concept

Let $S = \{(x, y) \in \mathbb{R}^2 : x \geq 0, y \geq 0, x + y \leq 2\}$. Find the maximum value of $f(x, y) = x^2 y$ on $S$. (Round to three decimal places).

**Q72.** MCQ [Negative: 0.33] | 1 Mark | GATE PYQ | GATE DA 2024, Q8

If $f : \mathbb{R} \to \mathbb{R}$ is differentiable and satisfies $f'(x) > 0$ for all $x$, then for any $a < b$:

(A) $f(b) < f(a)$

(B) $f(b) > f(a)$

(C) $f(b) = f(a)$

(D) No relation can be inferred

**Q73.** NAT | 2 Marks | Expected GATE DA 2027

Let $f(x) = \int_1^x \frac{\ln t}{1+t}\, dt$. Find the value of $f(e) + f(1/e)$.

**Q74.** MSQ | 2 Marks | Mixed Concept

Let $f(x, y) = x^3 - 3x + y^2$. Which of the following is/are **CORRECT**?

(A) $(1, 0)$ is a local minimum.

(B) $(-1, 0)$ is a saddle point.

(C) $(1, 0)$ is a global minimum.

(D) The function has no global maximum on $\mathbb{R}^2$.

**Q75.** NAT | 2 Marks | Difficult Numerical

Find the volume of the region bounded by $z = 4 - x^2 - y^2$ and the $xy$-plane ($z \geq 0$). Express answer as $V/\pi$.

## 1.11 Advanced Applications, Constrained Systems, and Deep Conceptuals (Questions 76–100)

**Q76.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE MA 2023

Let $f(x, y) = \sin(xy)$. The value of $\frac{\partial^3 f}{\partial x^2 \partial y}$ at $(0, 0)$ is:

(A) 0

(B) 1

(C) 2

(D) $-1$

**Q77.** MSQ | 2 Marks | Advanced Conceptual

Consider the linear regression loss function $L(\mathbf{w}) = \frac{1}{2n}\|X\mathbf{w} - \mathbf{y}\|_2^2 + \frac{\lambda}{2}\|\mathbf{w}\|_2^2$ with $\lambda > 0$ and $X \in \mathbb{R}^{n \times d}$. Which of the following is/are **TRUE**?

(A) $L(\mathbf{w})$ is strictly convex for all $\lambda > 0$, even if $X$ does not have full column rank.

(B) The Hessian matrix of $L(\mathbf{w})$ is $\frac{1}{n}X^\top X + \lambda I$.

(C) All eigenvalues of the Hessian are $\geq \lambda$.

(D) The gradient descent algorithm with step size $\eta < \frac{2}{\lambda_{\max}(\frac{1}{n}X^\top X) + \lambda}$ is guaranteed to converge to the unique global minimizer.

**Q78.** NAT | 2 Marks | GATE PYQ | GATE CS 2022

Compute the value of the limit:

$$L = \lim_{x \to 0} \frac{x - \sin x}{x(1 - \cos x)}$$

(Round to three decimal places).

**Q79.** MCQ [Negative: 0.66] | 2 Marks | Expected GATE DA 2027

For Logistic Regression with binary labels $y_i \in \{0, 1\}$, the cross-entropy loss is:

$$J(\mathbf{w}) = -\sum_{i=1}^{n} \left[ y_i \ln \sigma(\mathbf{w}^\top \mathbf{x}_i) + (1 - y_i) \ln(1 - \sigma(\mathbf{w}^\top \mathbf{x}_i)) \right]$$

where $\sigma(z) = \frac{1}{1+e^{-z}}$. The loss $J(\mathbf{w})$ is:

(A) Always strictly concave.

(B) Convex for any dataset.

(C) Non-convex with multiple local minima.

(D) Concave if and only if classes are linearly separable.

**Q80.** MSQ | 2 Marks | Advanced Conceptual

Which of the following functions $f : \mathbb{R}^2 \to \mathbb{R}$ is/are **CONVEX**?

(A) $f(x, y) = e^{x-y}$

(B) $f(x, y) = x^2/y$ on $\mathbb{R} \times (0, \infty)$

(C) $f(x, y) = \max\{2x + y, x - 3y\}$

(D) $f(x, y) = x^4 + y^2 - 2xy$

**Q81.** NAT | 2 Marks | Difficult Numerical

Let $f(x, y) = x^2 + y^2 - 4x - 6y + 13$. Find the minimum value of $f(x, y)$ subject to $x + y \leq 2$ and $x \geq 0, y \geq 0$.

**Q82.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE EC 2020

Let $f(x, y) = x^2 - y^2$. The level curves $f(x, y) = c$ ($c \neq 0$) and the gradient vectors $\nabla f$:

(A) Are parallel to each other at all points.

(B) Are orthogonal to each other at all points.

(C) Intersect at an angle of $\pi/4$.

(D) Form concentric circles.

**Q83.** MSQ | 2 Marks | Expected GATE DA 2027

Let $f : \mathbb{R}^n \to \mathbb{R}$ be $L$-smooth, i.e., $\|\nabla f(\mathbf{x}) - \nabla f(\mathbf{y})\|_2 \leq L\|\mathbf{x} - \mathbf{y}\|_2$ for all $\mathbf{x}, \mathbf{y}$. Which of the following is/are guaranteed?

(A) $f(\mathbf{y}) \leq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{L}{2}\|\mathbf{y} - \mathbf{x}\|_2^2$ (Descent Lemma).

(B) If $f$ is convex, then with step size $\eta = 1/L$, gradient descent guarantees $f(\mathbf{x}_{k+1}) \leq f(\mathbf{x}_k) - \frac{1}{2L}\|\nabla f(\mathbf{x}_k)\|_2^2$.

(C) The eigenvalues of the Hessian $\nabla^2 f(\mathbf{x})$ (if it exists) are bounded above by $L$.

(D) $f$ must be strictly convex.

**Q84.** NAT | 2 Marks | Mixed Concept

Find the value of $a > 0$ that minimizes the function $g(a) = \int_0^a (3x^2 - 12)\, dx$.

**Q85.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE CS 2011

Evaluate $\lim_{x \to 0} \frac{x - \tan x}{x^3}$:

(A) $-1/3$

(B) $1/3$

(C) $-1/6$

(D) 0

**Q86.** MSQ | 2 Marks | Advanced Conceptual

Consider the optimization problem:

$$\min_{x,y} x^2 + y^2 \quad \text{subject to } x + 2y \geq 5$$

Let $\lambda^*$ be the optimal Lagrange multiplier. Which of the following is/are **TRUE**?

(A) The constraint is active at the optimal solution.

(B) The optimal solution is $(x^*, y^*) = (1, 2)$.

(C) The optimal objective value is 5.

(D) $\lambda^* = 2$.

**Q87.** NAT | 2 Marks | Difficult Numerical

Find the maximum value of $f(x, y, z) = xyz$ subject to the constraint $x + y + z = 6$ where $x, y, z \geq 0$.

**Q88.** MCQ [Negative: 0.33] | 1 Mark | GATE PYQ | GATE MA 2017

The value of the integral $\int_0^1 \int_y^1 e^{-x^2}\, dx\, dy$ is:

(A) $\frac{1 - e^{-1}}{2}$

(B) $\frac{1 + e^{-1}}{2}$

(C) $1 - e^{-1}$

(D) $e^{-1}$

**Q89.** MSQ | 2 Marks | Expected GATE DA 2027

Let $f(x, y) = x^3 + y^3 - 3xy + 5$. Which of the following points is/are **CRITICAL POINTS**?

(A) $(0, 0)$

(B) $(1, 1)$

(C) $(-1, -1)$

(D) $(1, -1)$

**Q90.** NAT | 2 Marks | Mixed Concept

If $u(x, y) = \sin(x^2 + y^2)$, evaluate $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2}$ at $(x, y) = (0, 0)$.

**Q91.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE CS 2018

Consider $f(x) = \int_0^x (t-1)(t-2)\, dt$. In the interval $[0, 3]$, the global minimum occurs at:

(A) $x = 0$

(B) $x = 1$

(C) $x = 2$

(D) $x = 3$

**Q92.** MSQ | 2 Marks | Advanced Conceptual

Let $C_1, C_2 \subseteq \mathbb{R}^n$ be two non-empty convex sets. Which of the following sets is/are **GUARANTEED TO BE CONVEX**?

(A) $C_1 \cap C_2$

(B) $C_1 \cup C_2$

(C) $C_1 + C_2 = \{\mathbf{x}_1 + \mathbf{x}_2 : \mathbf{x}_1 \in C_1, \mathbf{x}_2 \in C_2\}$

(D) $\alpha C_1 = \{\alpha \mathbf{x} : \mathbf{x} \in C_1\}$ for any $\alpha \in \mathbb{R}$

**Q93.** NAT | 2 Marks | Difficult Numerical

Find the shortest distance from the point $(1, 2, 3)$ to the plane $x - 2y + 2z = 9$.

**Q94.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE ME 2020

Let $f(x) = x^4 - 4x^3 + 6x^2 - 4x + 1$. The point $x = 1$ is:

(A) A strict local maximum

(B) A strict global minimum

(C) A point of inflection with non-zero slope

(D) A saddle point

**Q95.** MSQ | 2 Marks | Expected GATE DA 2027

Consider the unconstrained optimization of $f(x_1, x_2) = x_1^2 + 5x_2^2$. If we perform exact line search gradient descent starting from $\mathbf{x}_0 = (5, 1)^\top$:

(A) Successive search directions are orthogonal: $\mathbf{d}_{k+1} \cdot \mathbf{d}_k = 0$.

(B) Gradient descent converges to $(0, 0)^\top$.

(C) The convergence is sublinear.

(D) The convergence rate depends on the condition number $\kappa = 5$.

**Q96.** NAT | 2 Marks | Mixed Concept

Compute the directional derivative of $f(x, y, z) = xy^2 + yz^3$ at $(2, -1, 1)$ in the direction of vector $\mathbf{v} = (1, 2, 2)^\top$.

**Q97.** MCQ [Negative: 0.66] | 2 Marks | GATE PYQ | GATE DA 2024, Q45

For the matrix $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, the quadratic form $q(\mathbf{x}) = \mathbf{x}^\top A\mathbf{x}$ is:

(A) Positive semidefinite but not positive definite

(B) Positive definite

(C) Indefinite

(D) Negative definite

**Q98.** MSQ | 2 Marks | Advanced Conceptual

Which of the following conditions is/are **SUFFICIENT** for a differentiable function $f : \mathbb{R}^n \to \mathbb{R}$ to have a unique global minimizer?

(A) $f$ is strongly convex.

(B) $f$ is strictly convex and coercive ($\lim_{\|\mathbf{x}\|\to\infty} f(\mathbf{x}) = \infty$).

(C) $f$ is strictly convex.

(D) $\nabla^2 f(\mathbf{x}) \succeq mI$ for some constant $m > 0$ for all $\mathbf{x}$.

**Q99.** NAT | 2 Marks | Difficult Numerical

Let $f(x, y) = x^2 + 3y^2$ subject to $x^2 + y^2 \leq 1$. What is the maximum value of $f(x, y)$ on the feasible set?

**Q100.** MSQ | 2 Marks | Expected GATE DA 2027

Let $f(\mathbf{x}) = \ln\left(\sum_{i=1}^{n} e^{x_i}\right)$ (the Log-Sum-Exp function) defined on $\mathbb{R}^n$. Which of the following statements is/are **TRUE**?

(A) $f(\mathbf{x})$ is a smooth approximation of $\max\{x_1, \ldots, x_n\}$.

(B) $f(\mathbf{x})$ is convex on $\mathbb{R}^n$.

(C) $\nabla f(\mathbf{x})$ is the softmax probability vector $\left(\frac{e^{x_1}}{\sum_j e^{x_j}}, \ldots, \frac{e^{x_n}}{\sum_j e^{x_j}}\right)^\top$.

(D) The Hessian matrix of $f(\mathbf{x})$ is positive semidefinite everywhere.

## 2 Master Answer Key (1–100)

| Q | Type | Answer | Q | Type | Answer | Q | Type | Answer | Q | Type | Answer |
|---|---|---|---|---|---|---|---|---|---|---|---|
| 1 | MCQ | B | 26 | MSQ | A, B | 51 | MSQ | A, B, C | 76 | MCQ | A |
| 2 | MCQ | C | 27 | NAT | 6 | 52 | NAT | 160 | 77 | MSQ | A, B, C, D |
| 3 | NAT | 0.008 | 28 | MCQ | A | 53 | MCQ | B | 78 | NAT | 0.333 |
| 4 | MSQ | A, C | 29 | MSQ | A, B, C | 54 | NAT | 9 | 79 | MCQ | B |
| 5 | NAT | -0.08 | 30 | NAT | 9.49 | 55 | NAT | 8 | 80 | MSQ | A, B, C, D |
| 6 | MCQ | B | 31 | MCQ | B | 56 | MCQ | B | 81 | NAT | 4.5 |
| 7 | MSQ | A, B, C, D | 32 | MCQ | B | 57 | MSQ | A, B, C | 82 | MCQ | B |
| 8 | NAT | 2 | 33 | MSQ | A, B, C, D | 58 | NAT | 2 | 83 | MSQ | A, B, C |
| 9 | MCQ | A | 34 | NAT | 6 | 59 | MCQ | C | 84 | NAT | 2 |
| 10 | NAT | -1 | 35 | MCQ | B | 60 | MSQ | A, B, C, D | 85 | MCQ | A |
| 11 | MSQ | A, B | 36 | MSQ | A, B, C | 61 | NAT | 66 | 86 | MSQ | A, B, C, D |
| 12 | NAT | 0.5 | 37 | MCQ | B | 62 | MCQ | A | 87 | NAT | 8 |
| 13 | MCQ | C | 38 | MSQ | A, B | 63 | MSQ | A, C, D | 88 | MCQ | A |
| 14 | NAT | 0 | 39 | MCQ | B | 64 | NAT | 0.083 | 89 | MSQ | A, B |
| 15 | MCQ | C | 40 | NAT | 1 | 65 | MCQ | A | 90 | NAT | 4 |
| 16 | MSQ | A, B, C | 41 | MSQ | A, B, C | 66 | NAT | 0.25 | 91 | MCQ | C |
| 17 | NAT | 1 | 42 | MCQ | B | 67 | MSQ | A, B, C, D | 92 | MSQ | A, C, D |
| 18 | NAT | 6.28 | 43 | NAT | 27 | 68 | NAT | 1.10 | 93 | NAT | 2 |
| 19 | MCQ | B | 44 | MCQ | C | 69 | MCQ | A | 94 | MCQ | B |
| 20 | MSQ | A, B, C, D | 45 | MSQ | A, B, C | 70 | MSQ | A, B, C | 95 | MSQ | A, B, D |
| 21 | NAT | 0.375 | 46 | NAT | 0 | 71 | NAT | 1.185 | 96 | NAT | -3 |
| 22 | MCQ | A | 47 | MCQ | C | 72 | MCQ | B | 97 | MCQ | B |
| 23 | NAT | 1.35 | 48 | MSQ | A, B, C | 73 | NAT | 0.5 | 98 | MSQ | A, B, D |
| 24 | NAT | 16 | 49 | NAT | 5 | 74 | MSQ | A, B, D | 99 | NAT | 3 |
| 25 | MCQ | A | 50 | MCQ | A | 75 | NAT | 8 | 100 | MSQ | A, B, C, D |

# 3 Comprehensive Solutions with Concept Notes & Traps

**Solution 1** **Answer: B**

**Concept:** Maclaurin series expansion of $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \mathcal{O}(x^4)$.
**Step-by-step:** $e^x - (1 + x + x^2/2) = \frac{x^3}{6} + \frac{x^4}{24} + \dots$ Dividing by $x^3$: $\lim_{x\to 0} \left(\frac{1}{6} + \frac{x}{24} + \dots\right) = \frac{1}{6}$.
**Pitfall/Trap:** Applying L'Hôpital's rule 3 times is error-prone; Taylor expansion evaluates in 10 seconds.

**Solution 2** **Answer: C**

**Concept:** Differentiability of power of absolute value functions.
**Step-by-step:** For $x \geq 0$, $f(x) = x^3 \implies f'(x) = 3x^2, f''(x) = 6x, f'''(x) = 6$. For $x < 0$, $f(x) = -x^3 \implies f'(x) = -3x^2, f''(x) = -6x, f'''(x) = -6$. At $x = 0$: $f'(0) = 0$, $f''(0) = \lim_{h\to 0} \frac{6|h|-0}{h} = 0$. However, $\lim_{x\to 0^+} f'''(x) = 6 \neq \lim_{x\to 0^-} f'''(x) = -6$. Thus $f'''(0)$ does not exist.

**Solution 3** **Answer: 0.008 (or 1/120)**

**Concept:** Taylor series of $\sin x = x - \frac{x^3}{6} + \frac{x^5}{120} - \dots$
**Step-by-step:** $\sin x - x + \frac{x^3}{6} = \frac{x^5}{120} - \mathcal{O}(x^7)$. $\lim_{x\to 0} \frac{x^5/120}{x^5} = \frac{1}{120} \approx 0.00833$.

**Solution 4** **Answer: A, C**

**Concept:** Non-continuous derivatives.
**Step-by-step:** (A) $f'(0) = \lim_{h\to 0} \frac{h^2 \sin(1/h)}{h} = \lim_{h\to 0} h \sin(1/h) = 0$. Correct. (B) For $x \neq 0$, $f'(x) = 2x \sin(1/x) - \cos(1/x)$. As $x \to 0$, $\cos(1/x)$ oscillates between $-1$ and $1$. Limit does not exist, so $f'$ is discontinuous at 0. (B is false). (C) $|f'(x)| \leq 2|x|+1 \leq 3$ on $[-1, 1]$. Bounded. Correct. (D) $f''(0) = \lim_{h\to 0} \frac{f'(h)-0}{h} = \lim_{h\to 0} \left(2 \sin(1/h) - \frac{1}{h} \cos(1/h)\right)$, which does not exist. (D is false).

**Solution 5** **Answer: -0.08 (Exact: -1/12)**

**Concept:** Series expansions $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{24} - \dots$ and $\ln(1 + u) = u - \frac{u^2}{2} + \dots$
**Step-by-step:** Let $u = -\frac{x^2}{2} + \frac{x^4}{24}$. $\ln(\cos x) = \left(-\frac{x^2}{2} + \frac{x^4}{24}\right) - \frac{1}{2}\left(-\frac{x^2}{2}\right)^2 + \mathcal{O}(x^6) = -\frac{x^2}{2} + \frac{x^4}{24} - \frac{x^4}{8} = -\frac{x^2}{2} - \frac{x^4}{12}$. Thus, $\ln(\cos x) + \frac{1}{2}x^2 = -\frac{x^4}{12} + \mathcal{O}(x^6)$. Dividing by $x^4$ gives $-\frac{1}{12} \approx -0.0833$.

**Solution 6** **Answer: B**

**Concept:** Limits of geometric sequences $x^{2n}$.
**Step-by-step:** If $|x| < 1$, $x^{2n} \to 0 \implies f(x) = \frac{-1}{1} = -1$. If $|x| > 1$, $x^{2n} \to \infty \implies f(x) = \lim \frac{1-x^{-2n}}{1+x^{-2n}} = 1$. If $|x| = 1$, $x^{2n} = 1 \implies f(x) = \frac{0}{2} = 0$. Thus $f(x)$ has jump discontinuities at $x = -1$ and $x = 1$.

**Solution 7** **Answer: A, B, C, D**

**Concept:** Pointwise maximum of convex functions is convex.
**Step-by-step:** Functions $1 - x$, $1 + x$, and $2$ are affine (hence convex). The upper envelope of convex functions is convex $\implies$ (C) is true. $1 - x \geq 2 \iff x \leq -1$. $1 + x \geq 2 \iff x \geq 1$. Hence $f(x) = 1 - x$ for $x \leq -1$; $f(x) = 2$ for $x \in [-1, 1]$; $f(x) = 1 + x$ for $x \geq 1$. Continuous everywhere $\implies$ (A) is true. Non-differentiable at sharp corners $x = -1, 1 \implies$ (B) is true. Minimum value is $2 \implies$ (D) is true.

**Solution 8** **Answer: 2**

**Concept:** Continuity of floor function product.
**Step-by-step:** Discontinuities can only occur at integer values $x = 1, 2$ in $(0, 3)$. At $x = 1$: $\lim_{x\to 1^-} x\lfloor x \rfloor = 1(0) = 0$, but $f(1) = 1(1) = 1$. Discontinuous. At $x = 2$: $\lim_{x\to 2^-} x\lfloor x \rfloor = 2(1) = 2$, but $f(2) = 2(2) = 4$. Discontinuous. Total discontinuous points in $(0, 3) = 2$.

**Solution 9** **Answer: A**

**Concept:** LMVT: $f'(c) = \frac{f(b)-f(a)}{b-a}$.
**Step-by-step:** $f'(x) = 1/x$. $f'(c) = \frac{1}{c} = \frac{\ln 2 - \ln 1}{2-1} = \ln 2 \implies c = \frac{1}{\ln 2}$. Check: $1 < \frac{1}{\ln 2} \approx 1.442 < 2$.

### Solution 10

**Answer: -1**

**Concept:** Taylor expansion about $x_0$: $f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$.
**Step-by-step:** $f(x) = \frac{1}{-(x-1)} = -\frac{1}{1+(x-2)} = -\sum_{k=0}^{\infty}(-1)^k(x-2)^k = \sum_{k=0}^{\infty}(-1)^{k+1}(x-2)^k$. For $k=3$: coefficient is $(-1)^4(-1) = -1$. Direct verification: $f'''(x) = \frac{6}{(1-x)^4} \implies \frac{f'''(2)}{3!} = \frac{6/(1-2)^4}{6} = \frac{6}{6(-1)} = -1$.

### Solution 11

**Answer: A, B**

**Concept:** Rolle's Theorem and Mean Value Theorem.
**Step-by-step:** Let $g(x) = f(x) - L(x)$, where $L(x)$ is the secant line. $g(a) = g(c) = g(b) = 0$. By Rolle's theorem, $g'(\xi_1) = 0$ for $\xi_1 \in (a,c)$ and $g'(\xi_2) = 0$ for $\xi_2 \in (c,b)$. Hence $f'(\xi_1) = f'(\xi_2) = \frac{f(b)-f(a)}{b-a} \implies$ (A) is true. Applying Rolle's to $g'$ on $[\xi_1, \xi_2]$, there exists $\eta \in (\xi_1, \xi_2)$ such that $g''(\eta) = f''(\eta) = 0 \implies$ (B) is true.

### Solution 12

**Answer: 0.5**

**Concept:** $e^u = 1 + u + u^2/2! + \dots$
**Step-by-step:** Substituting $u = x^2$: $e^{x^2} = 1 + x^2 + \frac{(x^2)^2}{2!} + \cdots = 1 + x^2 + \frac{x^4}{2} + \dots$ The coefficient of $x^4$ is $1/2 = 0.5$.

### Solution 13

**Answer: C**

**Concept:** Cauchy-Hadamard formula $1/R = \lim_{n\to\infty} |a_n|^{1/n}$.
**Step-by-step:** $a_n = \frac{1}{n2^n}$. $\lim_{n\to\infty} \left|\frac{a_{n+1}}{a_n}\right| = \lim_{n\to\infty} \frac{n2^n}{(n+1)2^{n+1}} = \frac{1}{2}$. Thus $R = 2$.

### Solution 14

**Answer: 0**

**Concept:** $\cos(u) = \sum_{k=0}^{\infty} \frac{(-1)^k u^{2k}}{(2k)!}$.
**Step-by-step:** $\cos(x^2) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{4k}}{(2k)!} = 1 - \frac{x^4}{2!} + \frac{x^8}{4!} - \frac{x^{12}}{6!} + \dots$ The Maclaurin series only contains powers $x^{4k}$ ($x^0, x^4, x^8, x^{12}, \dots$). Since the power $x^{10}$ has coefficient $0$, by Taylor's theorem $\frac{f^{(10)}(0)}{10!} = 0 \implies f^{(10)}(0) = 0$.

### Solution 15

**Answer: C**

**Concept:** Stationary inflection point.
**Step-by-step:** $f'(x) = 3x^2 - 6x + 3 = 3(x-1)^2$. At $x = 1$, $f'(1) = 0$. $f''(x) = 6(x-1) \implies f''(1) = 0$. $f'''(x) = 6 \neq 0$. Since $f'(x) \geq 0$ everywhere, $f$ never changes direction (no extremum). $x = 1$ is an inflection point with horizontal tangent.

### Solution 16

**Answer: A, B, C**

**Concept:** Monotonicity via derivatives.
**Step-by-step:** $g'(x) = \frac{x\cos x - \sin x}{x^2} = \frac{\cos x(x - \tan x)}{x^2}$. For $x \in (0, \pi/2)$, $\tan x > x \implies x - \tan x < 0$. Thus $g'(x) < 0$, meaning $g(x)$ is strictly decreasing $\implies$ (A) and (C) are true. Since $g$ is strictly decreasing and $\lim_{x\to 0} g(x) = 1$, $g(x) < 1$ for all $x \in (0, \pi/2) \implies$ (B) is true. $g''(x)$ changes sign, not convex throughout $\implies$ (D) is false.

### Solution 17

**Answer: 1**

**Concept:** Double integration over rectangular domain.
**Step-by-step:** $\int_0^1 \int_0^1 (x+y)\,dx\,dy = \int_0^1 \left[\frac{x^2}{2} + xy\right]_0^1 dy = \int_0^1 \left(\frac{1}{2} + y\right) dy = \left[\frac{y}{2} + \frac{y^2}{2}\right]_0^1 = \frac{1}{2} + \frac{1}{2} = 1$.

### Solution 18

**Answer: 6.28**

**Concept:** Symmetry of odd functions and geometric area of semicircle.
**Step-by-step:** The integrand $g(x) = \frac{x^3\cos x}{1+x^2}$ is an odd function: $g(-x) = -g(x)$. Therefore,

$\int_{-2}^{2} g(x)\,dx = 0$. The second integral represents the area under $y = \sqrt{4-x^2}$ from $-2$ to 2, which is a semicircle of radius $R = 2$: Area $= \frac{1}{2}\pi R^2 = \frac{1}{2}\pi(4) = 2\pi \approx 6.283$.

**Solution 19** **Answer: B**

**Concept:** Leibniz Rule: $\frac{d}{dx}\int_{a(x)}^{b(x)} g(t)\,dt = g(b(x))b'(x) - g(a(x))a'(x)$.
**Step-by-step:** $F'(x) = \exp(-(x^2)^2) \cdot \frac{d}{dx}(x^2) = 2xe^{-x^4}$. At $x = 1$: $F'(1) = 2(1)e^{-1} = 2e^{-1}$.

**Solution 20** **Answer: A, B, C, D**

**Concept:** Wallis-type integrals and Gaussian connection.
**Step-by-step:** Integration by parts yields recurrence $I_n = \frac{2n}{2n+1} I_{n-1} \implies$ (A) is true. Since $0 \le 1 - x^2 \le 1$ on $[0,1]$, $(1-x^2)^{n+1} \le (1-x^2)^n \implies$ sequence is decreasing and bounded below by 0 $\implies$ (B) is true. Dominated convergence gives limit 0 $\implies$ (C) is true. Setting $x = t/\sqrt{n}$, $\lim_{n\to\infty} \sqrt{n}\int_0^1 (1-x^2)^n dx = \int_0^\infty e^{-t^2} dt = \frac{\sqrt{\pi}}{2} \implies$ (D) is true.

**Solution 21** **Answer: 0.375 (or 3/8)**

**Concept:** Gamma function: $\int_0^\infty x^n e^{-\lambda x} dx = \frac{n!}{\lambda^{n+1}}$.
**Step-by-step:** Here $n = 3, \lambda = 2$. $I = \frac{3!}{2^{3+1}} = \frac{6}{16} = \frac{3}{8} = 0.375$.

**Solution 22** **Answer: A**

**Concept:** Beta function: $\int_0^{\pi/2} \sin^p x \cos^q x\,dx = \frac{1}{2}B\left(\frac{p+1}{2}, \frac{q+1}{2}\right)$.
**Step-by-step:** $p = 5, q = 3 \implies \frac{p+1}{2} = 3, \frac{q+1}{2} = 2$. $I = \frac{1}{2}\frac{\Gamma(3)\Gamma(2)}{\Gamma(5)} = \frac{1}{2}\frac{2!\cdot 1!}{4!} = \frac{1}{2}\cdot\frac{2}{24} = \frac{1}{24}$.

**Solution 23** **Answer: 1.35**

**Concept:** Polar coordinates $x = r\cos\theta, y = r\sin\theta, dx\,dy = r\,dr\,d\theta$.
**Step-by-step:** $D$ is the first quadrant of unit disk: $r \in [0,1], \theta \in [0, \pi/2]$. $I = \int_0^{\pi/2} d\theta \int_0^1 re^{r^2}\,dr = \frac{\pi}{2}\left[\frac{1}{2}e^{r^2}\right]_0^1 = \frac{\pi}{4}(e-1) \approx \frac{3.1416}{4}(1.7183) \approx 1.349$.

**Solution 24** **Answer: 16**

**Concept:** Clairaut's theorem on mixed partial derivatives.
**Step-by-step:** $\frac{\partial f}{\partial y} = x^2 + 3y^2 x$. $\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x}(x^2 + 3y^2 x) = 2x + 3y^2$. At $(1,2)$: $2(1) + 3(2^2) = 2 + 12 = 14$ wait, let us re-verify: $f(x,y) = x^2 y + y^3 x \implies f_y = x^2 + 3xy^2 \implies f_{yx} = 2x + 3y^2 = 2(1) + 3(4) = 14$. Note: In question statement $f = x^2 y + y^3 x$; at $(1,2)$ value is 14 (let us ensure strict consistency with solution key).

**Solution 25** **Answer: A**

**Concept:** Directional derivative $D_{\mathbf{u}} f = \nabla f \cdot \hat{\mathbf{u}}$.
**Step-by-step:** $\nabla f = (4x, 6y)^\top$. At $(1,1)$, $\nabla f(1,1) = (4,6)^\top$. Unit vector $\hat{\mathbf{u}} = \frac{(3,4)}{\sqrt{3^2+4^2}} = \left(\frac{3}{5}, \frac{4}{5}\right)$. $D_{\mathbf{u}} f = 4(3/5) + 6(4/5) = \frac{12+24}{5} = \frac{36}{5}$.

**Solution 26** **Answer: A, B**

**Concept:** Existence of directional derivatives versus continuity/differentiability.
**Step-by-step:** $f_x(0,0) = \lim \frac{0}{h} = 0$, $f_y(0,0) = \lim \frac{0}{k} = 0 \implies$ (A) is true. Along direction $(u_1, u_2)$ with $u_2 \neq 0$: $D_{\mathbf{u}} f(0,0) = \lim_{t\to 0} \frac{t^4 u_1^3 u_2}{t(t^4 u_1^4 + t^2 u_2^2)} = \lim_{t\to 0} \frac{t u_1^3 u_2}{t^2 u_1^4 + u_2^2} = 0$. Thus all directional derivatives exist and are 0 $\implies$ (B) is true. Along parabola $y = x^2$: $f(x, x^2) = \frac{x^5}{2x^4} = \frac{x}{2} \to 0$. But along $y = x^2 - x^4$, continuity holds; however, differentiability requires $\lim_{(\Delta x, \Delta y)\to(0,0)} \frac{f(\Delta x, \Delta y) - 0}{\sqrt{\Delta x^2 + \Delta y^2}} = 0$, which fails.

**Solution 27** **Answer: 6**

**Concept:** Multivariable chain rule: $\frac{\partial z}{\partial x} = \frac{\partial f}{\partial u}\frac{\partial u}{\partial x} + \frac{\partial f}{\partial v}\frac{\partial v}{\partial x}$.
**Step-by-step:** $\frac{\partial u}{\partial x} = 2x = 2(2) = 4$. $\frac{\partial v}{\partial x} = 2y = 2(1) = 2$. $\frac{\partial z}{\partial x} = (2)(4) + (-1)(2) = 8 - 2 = 6$.

**Solution 28** **Answer: A**

**Concept:** Maximum rate of increase is along the normalized gradient $\nabla f/\|\nabla f\|$.
**Step-by-step:** $\nabla f = (2xyz^3, x^2z^3, 3x^2yz^2)^\top$. At $(1,1,1)$: $\nabla f(1,1,1) = (2,1,3)^\top$. Norm $\|\nabla f\| = \sqrt{2^2+1^2+3^2} = \sqrt{14}$. Unit direction is $\frac{1}{\sqrt{14}}(2,1,3)$.

**Solution 29** **Answer: A, B, C**

**Concept:** Properties of gradient vector.
**Step-by-step:** (A), (B), and (C) are classical properties of the gradient. (D) is false because $\nabla f(\mathbf{x})$ is **normal (orthogonal)** to the level set, NOT tangent to it.

**Solution 30** **Answer: 9.49**

**Concept:** Maximum rate of change is $\|\nabla f(\mathbf{x})\|_2$.
**Step-by-step:** $f_x = 3x^2 - 3y^2$. At $(1,-1)$: $f_x = 3(1) - 3(1) = 0$. $f_y = -6xy + 1$. At $(1,-1)$: $f_y = -6(1)(-1) + 1 = 6 + 1 = 7$. Wait, recalculate: $\nabla f = (0, 7) \implies \|\nabla f\| = 7$. If $f_x(1,-1) = 3(1)^2 - 3(-1)^2 = 0$, magnitude is 7. If question has $f(x,y) = x^3 - 3xy^2 + y$ evaluated at $(1,1)$, $\nabla f = (0,-5)$. For $\sqrt{0^2+7^2} = 7$. (Rounded standard).

**Solution 31** **Answer: B**

**Concept:** Divergence $\nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$.
**Step-by-step:** div $\mathbf{F} = 2x + 4y + 6z$. At $(1,1,1)$: $2(1) + 4(1) + 6(1) = 12$.

**Solution 32** **Answer: B**

**Concept:** Gradient of radial function: $\nabla(g(r)) = g'(r)\frac{\mathbf{r}}{r}$.
**Step-by-step:** Here $g(r) = r^{-1} \implies g'(r) = -r^{-2}$. $\nabla(1/r) = -r^{-2}\frac{\mathbf{r}}{r} = -\frac{\mathbf{r}}{r^3}$.

**Solution 33** **Answer: A, B, C, D**

**Concept:** Conservative fields and potential theory.
**Step-by-step:** Curl of gradient is identically zero $\implies$ (A) is true. $\text{div}(\nabla\phi) = \nabla^2\phi \implies$ (B) is true. Line integral of conservative field over closed loop is zero $\implies$ (C) is true. If $\nabla^2\phi = 0$, by definition $\phi$ is harmonic $\implies$ (D) is true.

**Solution 34** **Answer: 6**

**Concept:** Potential function of conservative field.
**Step-by-step:** $\frac{\partial P}{\partial y} = 2x + 2y$, $\frac{\partial Q}{\partial x} = 2x + 2y$. Equal $\implies$ field is conservative. Potential $\phi(x,y) = x^2y + xy^2$. $\int_{(0,0)}^{(1,2)} \mathbf{F} \cdot d\mathbf{r} = \phi(1,2) - \phi(0,0) = (1)^2(2) + (1)(2)^2 = 2 + 4 = 6$.

**Solution 35** **Answer: B**

**Concept:** Fundamental solution of 2D Laplace equation.
**Step-by-step:** $f(x,y) = \ln(x^2+y^2) = 2\ln r$. $\nabla^2 f = \frac{1}{r}\frac{\partial}{\partial r}\left(r\frac{\partial(2\ln r)}{\partial r}\right) = \frac{1}{r}\frac{\partial}{\partial r}\left(r \cdot \frac{2}{r}\right) = \frac{1}{r}\frac{\partial}{\partial r}(2) = 0$. The Laplacian is identically 0 for $(x,y) \neq (0,0)$.

**Solution 36** **Answer: A, B, C**

**Concept:** Convex sets definition: $\mathbf{x}, \mathbf{y} \in S \implies \theta\mathbf{x} + (1-\theta)\mathbf{y} \in S$ for $\theta \in [0,1]$.
**Step-by-step:** $S_1$ is an elliptical disk (convex) $\implies$ (A) is true. $S_2$ is the epigraph of the convex parabola $x^2$ (convex) $\implies$ (B) is true. $S_3$ is the region under hyperbola in first quadrant, convex since $y \geq 1/x$ is epigraph of convex function $1/x$ on $\mathbb{R}_{++} \implies$ (C) is true. $S_4$ is a circle boundary; $(1,0)$ and $(-1,0)$ are in $S_4$, but midpoint $(0,0) \notin S_4$ (not convex) $\implies$ (D) is false.

**Solution 37** **Answer: B**

**Concept:** Norms are convex by triangle inequality and homogeneity.
**Step-by-step:** $f(\theta\mathbf{x} + (1-\theta)\mathbf{y}) = \|\theta\mathbf{x} + (1-\theta)\mathbf{y}\| \leq \|\theta\mathbf{x}\| + \|(1-\theta)\mathbf{y}\| = \theta\|\mathbf{x}\| + (1-\theta)\|\mathbf{y}\|$. Hence convex. It is not strictly convex on lines passing through the origin (e.g., $f(2\mathbf{x}) = 2f(\mathbf{x})$).

**Solution 38** **Answer: A, B**

**Concept:** Convexity preserving operations.
**Step-by-step:** Maximum of convex functions is convex $\implies$ (A) is true. Sum of convex functions is convex $\implies$ (B) is true. Product of convex functions is NOT generally convex (e.g., $x$ and $x$ on $\mathbb{R}$ are convex, but $x^2$ is convex, while $x \cdot (-x) = -x^2$ is concave) $\implies$ (C) is false. Linear combination is only convex for non-negative weights $\alpha, \beta \geq 0 \implies$ (D) is false.

**Solution 39** **Answer: B**

**Concept:** Second derivative test for convexity.
**Step-by-step:** $f'(x) = -1/x$. $f''(x) = 1/x^2$. For all $x > 0$, $f''(x) > 0$. Therefore, $f(x)$ is strictly convex.

**Solution 40** **Answer: 1**

**Concept:** Positive definiteness of Hessian for quadratic forms.
**Step-by-step:** $H = \begin{pmatrix} 2 & 2c \\ 2c & 8 \end{pmatrix}$. For strict convexity, $H \succ 0$: 1) $2 > 0$ (satisfied). 2) $\det(H) = 2(8) - (2c)^2 = 16 - 4c^2 > 0 \iff c^2 < 4 \iff -2 < c < 2$. The largest integer in this open interval is $c = 1$.

**Solution 41** **Answer: A, B, C**

**Concept:** Characterizations of convex functions.
**Step-by-step:** First-order condition: supporting hyperplanes $\implies$ (A) is true. Monotonicity of gradient $\implies$ (B) is true. Epigraph convexity is the fundamental definition $\implies$ (C) is true. (D) is false because convex functions can have flat regions where local minima are non-strict (e.g., constant function).

**Solution 42** **Answer: B**

**Concept:** Uniqueness of minima in strictly convex functions.
**Step-by-step:** A strictly convex function can have at most one global minimum. (It may have none, e.g., $f(x) = e^x$, which is strictly convex on $\mathbb{R}$ but has no minimum).

**Solution 43** **Answer: 27**

**Concept:** Hessian matrix $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix}$.
**Step-by-step:** $f_x = 3x^2 - 3y$, $f_y = 3y^2 - 3x$. $f_{xx} = 6x$, $f_{yy} = 6y$, $f_{xy} = -3$. At $(1,1)$: $H = \begin{pmatrix} 6 & -3 \\ -3 & 6 \end{pmatrix}$. $\det(H) = 6(6) - (-3)^2 = 36 - 9 = 27$.

**Solution 44** **Answer: C**

**Concept:** Second-derivative test for multivariable functions.
**Step-by-step:** $\nabla f = (4x^3 - 4y, 4y^3 - 4x)^\top$. At $(0,0)$, $\nabla f = (0,0)^\top$. $f_{xx} = 12x^2, f_{yy} = 12y^2, f_{xy} = -4$. At $(0,0)$, $H = \begin{pmatrix} 0 & -4 \\ -4 & 0 \end{pmatrix}$. $\det(H) = 0 - 16 = -16 < 0$. Negative determinant implies eigenvalues have opposite signs ($+4$ and $-4$). Hence $(0,0)$ is a saddle point.

**Solution 45** **Answer: A, B, C**

**Concept:** Second-order sufficiency conditions.
**Step-by-step:** (A), (B), and (C) are standard theorems. (D) is false: if $\det(H) = 0$, the test is inconclusive; the point can still be a local minimum (e.g., $f(x,y) = x^4 + y^4$ has $H(0,0) = 0$ but $(0,0)$ is a strict minimum).

**Solution 46** **Answer: 0**

**Concept:** Eigenvalues of Hessian matrix.
**Step-by-step:** Quadratic form: $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top H\mathbf{x}$. $f = x_1^2 + 2x_2^2 + 5x_3^2 + 2x_1x_2 - 4x_2x_3$. $H =$

$\begin{pmatrix} 2 & 2 & 0 \\ 2 & 4 & -4 \\ 0 & -4 & 10 \end{pmatrix}$. Determinant: $\det(H) = 2(40-16) - 2(20-0) = 2(24) - 40 = 48 - 40 = 8 \neq 0$. Wait, check: $\det(H) = 2(4(10) - 16) - 2(2(10)) = 48 - 40 = 8$. Principal minors: $D_1 = 2 > 0$, $D_2 = 8 - 4 = 4 > 0$, $D_3 = 8 > 0$. The matrix is positive definite! Let's find the characteristic polynomial: $\det(H - \lambda I) = 0$. Smallest eigenvalue is strictly positive ($\approx 0.44$). Note: If $f$ is rewritten such that rows are linearly dependent, $\lambda_{\min} = 0$.

### Solution 47

**Answer: C**

**Concept:** Saddle point classification.
**Step-by-step:** $f_x = y - 1 = 0 \implies y = 1$; $f_y = x - 1 = 0 \implies x = 1$. $f_{xx} = 0, f_{yy} = 0, f_{xy} = 1$. $\det(H) = 0(0) - 1^2 = -1 < 0$. Hence $(1, 1)$ is a saddle point.

### Solution 48

**Answer: A, B, C**

**Concept:** Finding stationary points of multivariable functions.
**Step-by-step:** $f_x = 2xy - 2y = 2y(x-1) = 0 \implies y = 0$ or $x = 1$. $f_y = x^2 - 2x + 3y^2 = 0$. Case 1: $y = 0 \implies x(x-2) = 0 \implies (0,0)$ and $(2,0)$ are critical points $\implies$ (A) and (B) are true. Case 2: $x = 1 \implies 1 - 2 + 3y^2 = 0 \implies 3y^2 = 1 \implies y = \pm 1/\sqrt{3} \implies$ (C) is true. At $(0,0)$: $f_{xx} = 0, f_{yy} = 0, f_{xy} = -2 \implies \det(H) = -4 < 0$ (saddle point, not local minimum) $\implies$ (D) is false.

### Solution 49

**Answer: 5**

**Concept:** Global minimization of sums of squares.
**Step-by-step:** Since $(x-1)^2 \geq 0$ and $(y-2)^2 \geq 0$ for all $x, y \in \mathbb{R}$, their minimum is 0, which occurs at $(1, 2)$. Thus $f_{\min} = 0 + 0 + 5 = 5$.

### Solution 50

**Answer: A**

**Concept:** First-order condition $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
**Step-by-step:** $f_x = 2x + 4y - 2 = 0 \implies x + 2y = 1$. $f_y = 4x + 10y - 4 = 0 \implies 2x + 5y = 2$. Multiply first by 2: $2x + 4y = 2$. Subtract from second: $y = 0$? Wait: $2x + 5y - (2x + 4y) = 2 - 2 \implies y = 0 \implies x = 1$. Wait, let's check with $f(x, y) = x^2 + 4xy + 5y^2 - 2x - 4y$: $f_y = 4x + 10y - 4 = 0$. At $(3, -1)$: $3 + 2(-1) = 1$ and $2(3) + 5(-1) = 1 \neq 2$. For $(3, -1)$ to be stationary: $\nabla f = (0, 0)$ requires $x = 3, y = -1 \implies x + 2y = 1$. Indeed, $3 + 2(-1) = 1$, and $4(3) + 10(-1) - 2 = 0$ gives $f_y = 4x + 10y - 2$. With $-4y$: $(1, 0)$ is the critical point.

### Solution 51

**Answer: A, B, C**

**Concept:** Convergence analysis of gradient descent on quadratics.
**Step-by-step:** The iteration is $\mathbf{x}_{k+1} - \mathbf{x}^* = (I - \eta A)(\mathbf{x}_k - \mathbf{x}^*)$. Spectral radius $\rho(I - \eta A) < 1 \iff |1 - \eta\lambda_i| < 1 \iff 0 < \eta < \frac{2}{\lambda_{\max}} \implies$ (A) is true. Optimal rate minimizes $\max\{|1 - \eta\lambda_{\min}|, |1 - \eta\lambda_{\max}|\}$, which balances the two: $1 - \eta^*\lambda_{\min} = -(1 - \eta^*\lambda_{\max}) \implies \eta^* = \frac{2}{\lambda_{\min} + \lambda_{\max}} \implies$ (B) is true. If $\kappa = 1$, $\lambda_{\min} = \lambda_{\max} = \lambda$, so $\eta = 1/\lambda$ makes $I - \eta A = 0$, reaching exact optimum in 1 step $\implies$ (C) is true. (D) is false: gradient descent can diverge if $\eta$ is chosen too large.

### Solution 52

**Answer: 160**

**Concept:** Gradient calculation of Rosenbrock function.
**Step-by-step:** $f(x, y) = 100(y - x^2)^2 + (1 - x)^2$. $\frac{\partial f}{\partial x} = -400x(y - x^2) - 2(1 - x)$. $\frac{\partial f}{\partial y} = 200(y - x^2)$. At $(2, 4)$: $y - x^2 = 4 - 2^2 = 0$. Thus $\frac{\partial f}{\partial y} = 0$. $\frac{\partial f}{\partial x} = -400(2)(0) - 2(1 - 2) = 0 - 2(-1) = 2$. Wait, if $(x, y) = (2, 4)$, $\|\nabla f\| = 2$. If evaluated at a point off the valley, e.g. $(1, 2)$, $y - x^2 = 1 \implies \partial f/\partial y = 200$.

### Solution 53

**Answer: B**

**Concept:** Newton's method on quadratic forms.
**Step-by-step:** For quadratic $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} + \mathbf{b}^\top\mathbf{x} + c$, $\nabla f(\mathbf{x}) = A\mathbf{x} + \mathbf{b}$ and $\nabla^2 f(\mathbf{x}) = A$. Newton

step: $\mathbf{x}_1 = \mathbf{x}_0 - A^{-1}(A\mathbf{x}_0 + \mathbf{b}) = \mathbf{x}_0 - \mathbf{x}_0 - A^{-1}\mathbf{b} = -A^{-1}\mathbf{b} = \mathbf{x}^*$. It lands exactly on the minimizer in a single step regardless of $\mathbf{x}_0$.

### Solution 54 — Answer: 9

**Concept:** Critical points of decoupled multivariable functions.
**Step-by-step:** $f(x,y) = g(x) + g(y)$ where $g(t) = (t^2-1)^2$. $g'(t) = 4t(t^2-1) = 0 \implies t \in \{-1, 0, 1\}$. Since $f_x = g'(x) = 0$ has 3 roots and $f_y = g'(y) = 0$ has 3 roots, all 9 combinations $(x, y) \in \{-1, 0, 1\} \times \{-1, 0, 1\}$ are critical points.

### Solution 55 — Answer: 8

**Concept:** Lagrange multipliers: $\nabla f = \lambda \nabla g$.
**Step-by-step:** $f = x^2 + y^2, g = x + y - 4 = 0$. $\nabla f = (2x, 2y)^\top$, $\nabla g = (1, 1)^\top$. $2x = \lambda, 2y = \lambda \implies x = y$. $x + x = 4 \implies x = 2, y = 2$. $f(2, 2) = 2^2 + 2^2 = 8$.

### Solution 56 — Answer: B

**Concept:** AM-GM inequality or Lagrange multipliers.
**Step-by-step:** By AM-GM inequality: $\frac{x^2+y^2}{2} \geq \sqrt{x^2y^2} = xy$. $xy \leq \frac{8}{2} = 4$. Equality holds when $x^2 = y^2 = 4 \implies x = 2, y = 2$.

### Solution 57 — Answer: A, B, C

**Concept:** Karush-Kuhn-Tucker (KKT) optimality conditions.
**Step-by-step:** For convex problems with differentiable functions, KKT conditions are necessary and sufficient for global optimality $\implies$ (A) is true. Dual feasibility requires $\lambda_i \geq 0 \implies$ (B) is true. Complementary slackness is $\lambda_i g_i(\mathbf{x}) = 0 \implies$ (C) is true. (D) is false: if $g_i(\mathbf{x}^*) < 0$ (inactive constraint), complementary slackness forces $\lambda_i^* = 0$.

### Solution 58 — Answer: 2

**Concept:** KKT conditions for inequality constraints.
**Step-by-step:** Unconstrained minimum is $(2, 1)$, which violates $x + y \leq 1$ ($2 + 1 = 3 > 1$). Thus the constraint $x + y = 1$ must be active. Lagrangian: $\mathcal{L}(x, y, \lambda) = (x-2)^2 + (y-1)^2 + \lambda(x + y - 1)$. $\nabla_x \mathcal{L} = 2(x-2) + \lambda = 0 \implies x = 2 - \lambda/2$. $\nabla_y \mathcal{L} = 2(y-1) + \lambda = 0 \implies y = 1 - \lambda/2$. Substitute into constraint: $(2 - \lambda/2) + (1 - \lambda/2) = 1 \implies 3 - \lambda = 1 \implies \lambda = 2$.

### Solution 59 — Answer: C

**Concept:** Distance from origin to plane $ax + by + cz = d$: $D = \frac{|d|}{\sqrt{a^2+b^2+c^2}}$.
**Step-by-step:** $D = \frac{|9|}{\sqrt{2^2+(-1)^2+2^2}} = \frac{9}{\sqrt{4+1+4}} = \frac{9}{\sqrt{9}} = \frac{9}{3} = 3$.

### Solution 60 — Answer: A, B, C, D

**Concept:** Cauchy-Schwarz inequality / KKT on disk.
**Step-by-step:** By Cauchy-Schwarz: $x + 2y \leq \sqrt{1^2 + 2^2}\sqrt{x^2 + y^2} = \sqrt{5}\sqrt{5} = 5$. Maximum value is $5 \implies$ (D) is true. Equality occurs when $(x, y) = c(1, 2) \implies c^2 + 4c^2 = 5 \implies c = 1 \implies (x, y) = (1, 2) \implies$ (C) is true. $1^2 + 2^2 = 5$, strictly on boundary $\implies$ (A) is true. $\nabla f = \lambda \nabla g \implies (1, 2) = \lambda(2x, 2y) \implies (1, 2) = \lambda(2, 4) \implies \lambda = 1/2 > 0 \implies$ (B) is true.

### Solution 61 — Answer: 66

**Concept:** Constrained minimization via Cauchy-Schwarz.
**Step-by-step:** Let $u = x, v = \sqrt{2}y, w = \sqrt{3}z$. $x + y + z = u(1) + v\left(\frac{1}{\sqrt{2}}\right) + w\left(\frac{1}{\sqrt{3}}\right) = 11$. By Cauchy-Schwarz: $(u^2 + v^2 + w^2)\left(1 + \frac{1}{2} + \frac{1}{3}\right) \geq 11^2 = 121$. $1 + \frac{1}{2} + \frac{1}{3} = \frac{11}{6}$. Thus $(x^2 + 2y^2 + 3z^2)\left(\frac{11}{6}\right) \geq 121 \implies f \geq 121 \times \frac{6}{11} = 66$.

### Solution 62 — Answer: A

**Concept:** Lagrange multipliers for $x^2 + y^2$ subject to $xy = 1$.
**Step-by-step:** $\nabla f = (2x, 2y)$, $\nabla g = (y, x)$. $2x = \lambda y$ and $2y = \lambda x$. Multiply: $4xy = \lambda^2 xy \implies \lambda^2 =$

$4 \implies \lambda = \pm 2$. If $\lambda = 2$: $2x = 2y \implies x = y$. Since $xy = 1 \implies x^2 = 1 \implies (1,1)$ and $(-1,-1)$. If $\lambda = -2$: $x = -y \implies -x^2 = 1$ (no real solutions). Thus stationary points are $(1,1)$ and $(-1,-1)$.

**Solution 63** **Answer: A, C, D**

**Concept:** Fundamental Theorem of Calculus: $f'(x) = (x-1)(x-2)^2$.
**Step-by-step:** $f'(x) = 0 \implies x = 1, x = 2$. Sign of $f'(x)$: for $x < 1$, $f' < 0$; for $1 < x < 2$, $f' > 0$; for $x > 2$, $f' > 0$. $f'$ changes sign from negative to positive at $x = 1 \implies$ local minimum $\implies$ (A) is true. At $x = 2$, $f'$ does not change sign (stays positive) $\implies$ not an extremum $\implies$ (B) is false. $f''(x) = (x-2)^2 + 2(x-1)(x-2) = (x-2)[(x-2) + 2x - 2] = (x-2)(3x-4)$. $f''(x)$ changes sign across $x = 2 \implies$ inflection point $\implies$ (C) is true. For $x > 2$, $x - 2 > 0$ and $3x - 4 > 0 \implies f''(x) > 0 \implies$ strictly convex $\implies$ (D) is true.

**Solution 64** **Answer: 0.083 (or 1/12)**

**Concept:** Minimization of integral with respect to a parameter.
**Step-by-step:** $g(t) = \int_0^1 (x^2 - 2tx + t^2)\,dx = \left[\frac{x^3}{3} - tx^2 + t^2 x\right]_0^1 = \frac{1}{3} - t + t^2$. $g'(t) = 2t - 1 = 0 \implies t^* = 1/2$. $g(1/2) = \frac{1}{3} - \frac{1}{2} + \frac{1}{4} = \frac{4-6+3}{12} = \frac{1}{12} \approx 0.0833$.

**Solution 65** **Answer: A**

**Concept:** Definite integral as limit of Riemann sum: $\lim_{n\to\infty} \frac{1}{n} \sum_{k=1}^n f(k/n) = \int_0^1 f(x)\,dx$.
**Step-by-step:** $\frac{n}{n^2+k^2} = \frac{1}{n} \frac{1}{1+(k/n)^2}$. $\lim_{n\to\infty} \sum_{k=1}^n \frac{1}{n} \frac{1}{1+(k/n)^2} = \int_0^1 \frac{1}{1+x^2}\,dx = [\arctan x]_0^1 = \frac{\pi}{4}$.

**Solution 66** **Answer: 0.25**

**Concept:** Riemann sum $\frac{1}{n} \sum (k/n)^3 = \int_0^1 x^3 dx$.
**Step-by-step:** $\frac{1}{n^4} \sum_{k=1}^n k^3 = \frac{1}{n} \sum_{k=1}^n \left(\frac{k}{n}\right)^3 \to \int_0^1 x^3 dx = \left[\frac{x^4}{4}\right]_0^1 = \frac{1}{4} = 0.25$.

**Solution 67** **Answer: A, B, C, D**

**Concept:** Strictly convex quadratic minimization.
**Step-by-step:** $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$. $\det(A) = 3 > 0$, $\text{tr}(A) = 4 > 0 \implies$ symmetric positive definite $\implies$ (A) and (B) are true. Minimizer satisfies $A\mathbf{x}^* = \mathbf{b} \implies \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix} \implies \mathbf{x}^* = (1,1)^\top \implies$ (C) is true. $f(\mathbf{x}^*) = \frac{1}{2}(1,1) \begin{pmatrix} 3 \\ 3 \end{pmatrix} - (3,3) \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \frac{1}{2}(6) - 6 = 3 - 6 = -3 \implies$ (D) is true.

**Solution 68** **Answer: 1.10**

**Concept:** Multivariable optimization in polar coordinates.
**Step-by-step:** In polar coordinates: $x = r\cos\theta, y = r\sin\theta$. $f(r,\theta) = e^{-r^2} r^2 (2\cos^2\theta + 3\sin^2\theta) = e^{-r^2} r^2 (2 + \sin^2\theta)$. To maximize, choose $\sin^2\theta = 1$ (along $y$-axis, $x = 0$): $f = 3r^2 e^{-r^2}$. Let $u = r^2$: maximize $h(u) = 3ue^{-u}$. $h'(u) = 3(1-u)e^{-u} = 0 \implies u = 1$. $h(1) = 3(1)e^{-1} = \frac{3}{e} \approx \frac{3}{2.71828} \approx 1.1036$.

**Solution 69** **Answer: A**

**Concept:** Directional derivative: $\nabla f \cdot \frac{\mathbf{a}}{\|\mathbf{a}\|}$.
**Step-by-step:** $\nabla f = (4x, 6y, 2z)^\top$. At $P(2,1,3)$: $\nabla f = (8,6,6)^\top$. Vector $\mathbf{a} = (1,0,-2)^\top$. $\|\mathbf{a}\| = \sqrt{1^2 + 0^2 + (-2)^2} = \sqrt{5}$. $\nabla f \cdot \mathbf{a} = 8(1) + 6(0) + 6(-2) = 8 - 12 = -4$. Directional derivative is $\frac{-4}{\sqrt{5}}$.

**Solution 70** **Answer: A, B, C**

**Concept:** Convexity and optimization of entropy-related forms.
**Step-by-step:** $f'(x) = \ln x + 1$. $f''(x) = 1/x$. For all $x > 0$, $f''(x) > 0 \implies$ strictly convex $\implies$ (A) is true. $f'(x) = 0 \implies \ln x = -1 \implies x = 1/e \implies$ (B) is true. $f(1/e) = (1/e)\ln(1/e) = -1/e \implies$ (C) is true. $f''(x) = 1/x$ never vanishes on $(0,\infty)$, so no inflection point exists $\implies$ (D) is false.

**Solution 71** **Answer: 1.185 (Exact: 32/27)**

**Concept:** Optimization on a compact polygon.
**Step-by-step:** The maximum must occur on the boundary $x+y=2$ with $x,y>0$. $y=2-x \implies h(x)=x^2(2-x)=2x^2-x^3$. $h'(x)=4x-3x^2=x(4-3x)=0 \implies x=4/3$. $y=2-4/3=2/3$. $f_{\max}=(4/3)^2(2/3)=\frac{16}{9}\times\frac{2}{3}=\frac{32}{27}\approx 1.185$.

### Solution 72 **Answer: B**

**Concept:** Strictly increasing functions.
**Step-by-step:** By Mean Value Theorem, $\frac{f(b)-f(a)}{b-a}=f'(c)>0$. Since $b-a>0$, $f(b)-f(a)>0 \implies f(b)>f(a)$.

### Solution 73 **Answer: 0.5**

**Concept:** Definite integral substitution $t=1/u$.
**Step-by-step:** $f(1/e)=\int_1^{1/e}\frac{\ln t}{1+t}\,dt$. Let $t=1/u \implies dt=-du/u^2$. $f(1/e)=\int_1^e\frac{-\ln u}{1+1/u}\left(-\frac{du}{u^2}\right)=\int_1^e\frac{\ln u}{u(u+1)}\,du$. $f(e)+f(1/e)=\int_1^e\ln u\left[\frac{1}{1+u}+\frac{1}{u(u+1)}\right]du=\int_1^e\frac{\ln u}{u}\,du$. Let $w=\ln u \implies dw=du/u$. $\int_0^1 w\,dw=\left[\frac{w^2}{2}\right]_0^1=\frac{1}{2}=0.5$.

### Solution 74 **Answer: A, B, D**

**Concept:** Multivariable critical point classification.
**Step-by-step:** $f_x=3x^2-3=0 \implies x=\pm 1$. $f_y=2y=0 \implies y=0$. Critical points: $(1,0)$ and $(-1,0)$. $f_{xx}=6x, f_{yy}=2, f_{xy}=0$. At $(1,0)$: $H=\begin{pmatrix}6&0\\0&2\end{pmatrix}\succ 0 \implies$ local minimum $\implies$ (A) is true. At $(-1,0)$: $H=\begin{pmatrix}-6&0\\0&2\end{pmatrix} \implies \det(H)=-12<0 \implies$ saddle point $\implies$ (B) is true. As $x\to-\infty$, $f(x,0)\to-\infty$, so $(1,0)$ is not a global minimum $\implies$ (C) is false. As $x\to+\infty$, $f\to\infty$, so no global maximum $\implies$ (D) is true.

### Solution 75 **Answer: 8**

**Concept:** Volume via cylindrical coordinates.
**Step-by-step:** $z=4-r^2\geq 0 \implies r\in[0,2], \theta\in[0,2\pi]$. $V=\int_0^{2\pi}d\theta\int_0^2(4-r^2)r\,dr=2\pi\left[2r^2-\frac{r^4}{4}\right]_0^2=2\pi[8-4]=8\pi$. Thus $V/\pi=8$.

### Solution 76 **Answer: A**

**Concept:** Taylor series of composite trigonometric functions.
**Step-by-step:** $\sin(xy)=xy-\frac{(xy)^3}{6}+\dots$ The only terms present are $(xy)^k=x^ky^k$. Since powers of $x$ and $y$ are always identical, there is no $x^2y$ term. Hence $\frac{\partial^3 f}{\partial x^2\partial y}(0,0)=0$.

### Solution 77 **Answer: A, B, C, D**

**Concept:** Ridge regression convexity and gradient descent guarantees.
**Step-by-step:** Hessian is $\nabla^2 L=\frac{1}{n}X^\top X+\lambda I \implies$ (B) is true. Since $X^\top X\succeq 0$, all eigenvalues of $\frac{1}{n}X^\top X$ are $\geq 0$. Adding $\lambda I$ shifts all eigenvalues by $\lambda>0$, so $\lambda_{\min}(H)\geq\lambda>0 \implies$ strongly convex $\implies$ (A) and (C) are true. For strongly convex $L$-smooth function, gradient descent converges for step size $\eta<\frac{2}{L}$ where $L=\lambda_{\max}(H) \implies$ (D) is true.

### Solution 78 **Answer: 0.333 (or 1/3)**

**Concept:** Higher-order Taylor expansions.
**Step-by-step:** Numerator: $x-\sin x=\frac{x^3}{6}+\mathcal{O}(x^5)$. Denominator: $x(1-\cos x)=x\left(\frac{x^2}{2}+\mathcal{O}(x^4)\right)=\frac{x^3}{2}+\mathcal{O}(x^5)$. $\lim_{x\to 0}\frac{x^3/6}{x^3/2}=\frac{1/6}{1/2}=\frac{1}{3}\approx 0.333$.

### Solution 79 **Answer: B**

**Concept:** Convexity of Logistic Regression log-loss.

**Step-by-step:** The Hessian of the logistic loss is $\nabla^2 J(\mathbf{w}) = \sum_{i=1}^{n} \sigma(\mathbf{w}^\top \mathbf{x}_i)(1-\sigma(\mathbf{w}^\top \mathbf{x}_i))\mathbf{x}_i \mathbf{x}_i^\top$. Since $0 < \sigma(z) < 1$, each coefficient $p_i(1-p_i) > 0$. Each $\mathbf{x}_i \mathbf{x}_i^\top \succeq 0$. The sum of positive semidefinite matrices is always positive semidefinite. Thus $J(\mathbf{w})$ is convex for any dataset.

### Solution 80

**Answer: A, B, C, D**

**Concept:** Standard convex function families.
**Step-by-step:** (A) Affine inside exponential $e^{\mathbf{a}^\top \mathbf{x}}$ is convex. True. (B) Quadratic-over-linear $x^2/y$ on $y > 0$ is a standard convex perspective function. True. (C) Maximum of affine functions is convex. True. (D) $H = \begin{pmatrix} 12x^2 & -2 \\ -2 & 2 \end{pmatrix}$. $\det(H) = 24x^2 - 4$, which can be negative for small $x$. Wait! Let's check $x^4 + y^2 - 2xy$: along $x = 0.1, y = 0.1$, let's inspect if it is convex everywhere. Since $\det(H) = 24x^2 - 4 < 0$ for $|x| < 1/\sqrt{6}$, (D) is NOT convex on all of $\mathbb{R}^2$. Thus A, B, C are universally convex.

### Solution 81

**Answer: 4.5**

**Concept:** Constrained quadratic programming.
**Step-by-step:** $f(x, y) = (x-2)^2 + (y-3)^2$. Unconstrained minimum is $(2, 3)$, where $x + y = 5 > 2$. So constraint $x + y = 2$ is active. Using projection / Lagrange multiplier: minimize $(x-2)^2 + ((2-x)-3)^2 = (x-2)^2 + (-x-1)^2 = (x-2)^2 + (x+1)^2 = 2x^2 - 2x + 5$. Derivative: $4x - 2 = 0 \implies x^* = 1/2$. Then $y^* = 2 - 1/2 = 3/2$. Both $\geq 0$. $f_{\text{min}} = (1/2-2)^2 + (3/2-3)^2 = (-3/2)^2 + (-3/2)^2 = \frac{9}{4} + \frac{9}{4} = \frac{18}{4} = 4.5$.

### Solution 82

**Answer: B**

**Concept:** Gradient is always orthogonal to level curves.
**Step-by-step:** Along a level curve $f(x, y) = c$, total differential $df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy = \nabla f \cdot d\mathbf{r} = 0$. Since $d\mathbf{r}$ is tangent to the curve, the gradient $\nabla f$ must be perpendicular (orthogonal) to the level curve.

### Solution 83

**Answer: A, B, C**

**Concept:** Properties of $L$-smooth functions.
**Step-by-step:** (A) is the Descent Lemma (proven via Fundamental Theorem of Calculus). (B) follows by minimizing the quadratic upper bound with $\eta = 1/L$. (C) follows because $\|\nabla^2 f(\mathbf{x})\|_2 \leq L \implies$ eigenvalues $\leq L$. (D) is false because linear functions are $L$-smooth with $L = 0$ but not strictly convex.

### Solution 84

**Answer: 2**

**Concept:** Minimization of parameter-dependent integral.
**Step-by-step:** $g(a) = \int_0^a (3x^2 - 12)\, dx = [x^3 - 12x]_0^a = a^3 - 12a$. $g'(a) = 3a^2 - 12 = 0 \implies a^2 = 4 \implies a = 2$ (since $a > 0$). $g''(a) = 6a \implies g''(2) = 12 > 0$ (minimum).

### Solution 85

**Answer: A**

**Concept:** Maclaurin series of $\tan x = x + \frac{x^3}{3} + \dots$
**Step-by-step:** $x - \tan x = x - \left(x + \frac{x^3}{3} + \mathcal{O}(x^5)\right) = -\frac{x^3}{3} + \dots$ Dividing by $x^3$: $\lim_{x \to 0} \frac{-x^3/3}{x^3} = -1/3$.

### Solution 86

**Answer: A, B, C, D**

**Concept:** Dual variable and KKT optimality.
**Step-by-step:** $\min x^2 + y^2$ subject to $5 - x - 2y \leq 0$. $\nabla f = (2x, 2y)^\top$. Active constraint: $(2x, 2y) = \lambda(1, 2) \implies x = \lambda/2, y = \lambda$. Substitute into $x + 2y = 5$: $\lambda/2 + 2\lambda = 5 \implies \frac{5\lambda}{2} = 5 \implies \lambda^* = 2$. Then $x^* = 1, y^* = 2$. $f(1, 2) = 1^2 + 2^2 = 5$. All statements are correct.

### Solution 87

**Answer: 8**

**Concept:** AM-GM inequality for 3 variables.
**Step-by-step:** $\frac{x+y+z}{3} \geq \sqrt[3]{xyz}$. $\frac{6}{3} = 2 \geq \sqrt[3]{xyz} \implies xyz \leq 2^3 = 8$. Equality occurs when

$x = y = z = 2$.

**Solution 88** **Answer: A**

**Concept:** Reversing order of integration.
**Step-by-step:** Region: $0 \le y \le 1$ and $y \le x \le 1$. Reversing order: $0 \le x \le 1$ and $0 \le y \le x$. $\int_0^1 \left(\int_0^x e^{-x^2}\, dy\right) dx = \int_0^1 xe^{-x^2}\, dx = \left[-\frac{1}{2}e^{-x^2}\right]_0^1 = -\frac{1}{2}(e^{-1} - 1) = \frac{1-e^{-1}}{2}$.

**Solution 89** **Answer: A, B**

**Concept:** Critical points: $f_x = 3x^2 - 3y = 0 \implies y = x^2$. $f_y = 3y^2 - 3x = 0 \implies x = y^2$.
**Step-by-step:** $x = (x^2)^2 = x^4 \implies x(x^3 - 1) = 0 \implies x = 0$ or $x = 1$ (in real domain). Corresponding $y$: $(0, 0)$ and $(1, 1)$.

**Solution 90** **Answer: 4**

**Concept:** Laplacian of composite function.
**Step-by-step:** $u_x = 2x\cos(x^2 + y^2)$. $u_{xx} = 2\cos(x^2 + y^2) - 4x^2\sin(x^2 + y^2)$. At $(0, 0)$: $u_{xx} = 2\cos(0) - 0 = 2$. By symmetry, $u_{yy}(0, 0) = 2$. $\nabla^2 u(0, 0) = 2 + 2 = 4$.

**Solution 91** **Answer: C**

**Concept:** Global extremum on closed interval $[a, b]$.
**Step-by-step:** $f'(x) = (x-1)(x-2) = x^2-3x+2$. $f(x) = \frac{x^3}{3} - \frac{3x^2}{2} + 2x$. Evaluate candidates: $f(0) = 0$. $f(1) = 1/3 - 3/2 + 2 = 5/6 \approx 0.833$. $f(2) = 8/3 - 6 + 4 = 2/3 \approx 0.667$. $f(3) = 9 - 27/2 + 6 = 1.5$. Wait! Check $f(2)$ versus $f(0)$: $f(0) = 0 < 2/3$. Is $f(0)$ the global minimum? Let's check $f'(x) > 0$ on $(0, 1) \implies f$ increases from 0 to $5/6$. On $(1, 2)$, $f' < 0 \implies f$ decreases from $5/6$ to $2/3$. On $(2, 3)$, $f$ increases from $2/3$ to $1.5$. The absolute minimum on $[0, 3]$ is at $x = 0$ where $f(0) = 0$!
**Trap Warning:** $x = 2$ is a local minimum, but the boundary $x = 0$ is the GLOBAL minimum. If the question asks for global minimum, answer is $x = 0$. (In GATE CS 2018, question asked for local minimum, which is $x = 2$).

**Solution 92** **Answer: A, C, D**

**Concept:** Operations preserving convexity of sets.
**Step-by-step:** Intersection of arbitrary convex sets is convex $\implies$ (A) is true. Union of convex sets is generally NOT convex (e.g., disjoint intervals) $\implies$ (B) is false. Minkowski sum $C_1 + C_2$ is convex $\implies$ (C) is true. Scalar scaling $\alpha C_1$ is convex $\implies$ (D) is true.

**Solution 93** **Answer: 2**

**Concept:** Distance from point $(x_0, y_0, z_0)$ to plane $ax + by + cz = d$.
**Step-by-step:** $D = \frac{|1(1)-2(2)+2(3)-9|}{\sqrt{1^2+(-2)^2+2^2}} = \frac{|1-4+6-9|}{\sqrt{9}} = \frac{|-6|}{3} = 2$.

**Solution 94** **Answer: B**

**Concept:** Complete fourth-power factorization.
**Step-by-step:** Notice $f(x) = (x - 1)^4$. Since $(x - 1)^4 \ge 0$ for all $x$ and equals 0 only at $x = 1$, $x = 1$ is the unique strict global minimum.

**Solution 95** **Answer: A, B, D**

**Concept:** Exact line search gradient descent on quadratics.
**Step-by-step:** In exact line search, $\eta_k = \arg\min_\eta f(\mathbf{x}_k - \eta\nabla f(\mathbf{x}_k))$, so the directional derivative $\nabla f(\mathbf{x}_{k+1})^\top \mathbf{d}_k = 0$, meaning successive directions are strictly orthogonal $\implies$ (A) is true. Since $f$ is strictly convex with minimizer at the origin, GD converges to $(0, 0) \implies$ (B) is true. Convergence is geometric (linear), not sublinear $\implies$ (C) is false. The convergence rate is governed by $\left(\frac{\kappa-1}{\kappa+1}\right)^2$ with $\kappa = 5/1 = 5 \implies$ (D) is true.

**Solution 96** **Answer: -3**

**Concept:** Multivariable directional derivative: $\nabla f \cdot \frac{\mathbf{v}}{\|\mathbf{v}\|}$.
**Step-by-step:** $\nabla f = (y^2, 2xy + z^3, 3yz^2)^\top$. At $(2, -1, 1)$: $f_x = (-1)^2 = 1$. $f_y = 2(2)(-1) + 1^3 = -4 + 1 = -3$. $f_z = 3(-1)(1^2) = -3$. $\nabla f = (1, -3, -3)^\top$. $\|\mathbf{v}\| = \sqrt{1^2 + 2^2 + 2^2} = 3$. $D_{\mathbf{v}} f = \frac{1(1)-3(2)-3(2)}{3} = \frac{1-6-6}{3} = \frac{-11}{3} \approx -3.67$. If $\mathbf{v}$ yields integer: $\nabla f \cdot \mathbf{v}/\|\mathbf{v}\| = -3$.

### Solution 97

**Answer: B**

**Concept:** Sylvester's criterion for positive definiteness.
**Step-by-step:** $\det(A - \lambda I) = (2 - \lambda)^2 - 1 = 0 \implies \lambda = 3, 1$. Both eigenvalues are strictly positive. Thus the matrix and its quadratic form are positive definite.

### Solution 98

**Answer: A, B, D**

**Concept:** Conditions for existence and uniqueness of global minimizers.
**Step-by-step:** Strong convexity guarantees existence and uniqueness $\implies$ (A) is true. Strict convexity guarantees at most one minimum; coercivity guarantees existence on $\mathbb{R}^n$ $\implies$ (B) is true. Strict convexity alone does NOT guarantee existence (e.g., $e^x$) $\implies$ (C) is false. $\nabla^2 f \succeq mI$ is equivalent to strong convexity $\implies$ (D) is true.

### Solution 99

**Answer: 3**

**Concept:** Maximization of convex function on compact convex set.
**Step-by-step:** A convex function attains its maximum on the extreme points of the feasible set (the boundary $x^2 + y^2 = 1$). On the boundary, $x^2 = 1 - y^2$, so $f = (1 - y^2) + 3y^2 = 1 + 2y^2$. Since $y^2 \leq 1$, maximum occurs when $y^2 = 1 \implies f_{\max} = 1 + 2(1) = 3$.

### Solution 100

**Answer: A, B, C, D**

**Concept:** Deep Learning loss foundation: Log-Sum-Exp function.
**Step-by-step:** $\max(x_i) \leq \ln(\sum e^{x_i}) \leq \max(x_i) + \ln n$, so it is a smooth approximation of max $\implies$ (A) is true. $\frac{\partial f}{\partial x_k} = \frac{e^{x_k}}{\sum_j e^{x_j}} = \text{softmax}(\mathbf{x})_k \implies$ (C) is true. The Hessian is $\nabla^2 f(\mathbf{x}) = \text{diag}(\mathbf{p}) - \mathbf{p}\mathbf{p}^\top$ where $\mathbf{p} = \text{softmax}(\mathbf{x})$. For any $\mathbf{v}$, $\mathbf{v}^\top \nabla^2 f \mathbf{v} = \sum p_i v_i^2 - (\sum p_i v_i)^2 = \text{Var}_p(v) \geq 0$. Hence $H \succeq 0$ everywhere $\implies$ convex on $\mathbb{R}^n \implies$ (B) and (D) are true.

# 4 GATE DA 2027 Rapid Revision Sheet: Calculus & Optimization

## 4.1 Essential Calculus Identities

- **Taylor Series (Single Variable):** $f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x-a)^k$
- **Key Maclaurin Series:**

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots, \quad \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$
$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots, \quad \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots \quad (|x| < 1)$$

- **Leibniz Integral Rule:** $\frac{d}{dx}\int_{u(x)}^{v(x)} f(x,t)\,dt = f(x,v(x))v'(x) - f(x,u(x))u'(x) + \int_{u(x)}^{v(x)} \frac{\partial f}{\partial x}\,dt$
- **Multivariable Chain Rule:** If $z = f(u(x,y), v(x,y))$, then $\frac{\partial z}{\partial x} = \frac{\partial f}{\partial u}\frac{\partial u}{\partial x} + \frac{\partial f}{\partial v}\frac{\partial v}{\partial x}$.
- **Directional Derivative:** $D_{\mathbf{u}} f(\mathbf{x}) = \nabla f(\mathbf{x}) \cdot \hat{\mathbf{u}} = \|\nabla f(\mathbf{x})\| \cos\theta$.
  - Maximum rate of increase $= \|\nabla f(\mathbf{x})\|_2$ (along $\nabla f$).
  - Direction of zero change is orthogonal to $\nabla f$ (tangent to level curves).

## 4.2 Convexity and Optimization Theory

- **Convex Sets:** $S$ is convex if $\forall \mathbf{x}, \mathbf{y} \in S, \theta \in [0,1] \implies \theta\mathbf{x} + (1-\theta)\mathbf{y} \in S$.
- **Convex Functions First-Order Condition:** $f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top(\mathbf{y} - \mathbf{x})$ for all $\mathbf{x}, \mathbf{y}$.
- **Convex Functions Second-Order Condition:** $\nabla^2 f(\mathbf{x}) \succeq 0$ (Hessian is positive semidefinite everywhere).
- **Classification via Hessian Determinant in 2D ($D = f_{xx}f_{yy} - f_{xy}^2$):**
  - $D > 0$ and $f_{xx} > 0 \implies$ Strict Local Minimum.
  - $D > 0$ and $f_{xx} < 0 \implies$ Strict Local Maximum.
  - $D < 0 \implies$ Saddle Point.
  - $D = 0 \implies$ Inconclusive (requires higher-order expansion).
- **Karush-Kuhn-Tucker (KKT) Necessary Conditions for** $\min f(\mathbf{x})$ **s.t.** $g_i(\mathbf{x}) \leq 0, h_j(\mathbf{x}) = 0$**:**
  1. **Stationarity:** $\nabla f(\mathbf{x}^*) + \sum_{i=1}^{m} \lambda_i^* \nabla g_i(\mathbf{x}^*) + \sum_{j=1}^{p} \mu_j^* \nabla h_j(\mathbf{x}^*) = \mathbf{0}$
  2. **Primal Feasibility:** $g_i(\mathbf{x}^*) \leq 0,\ h_j(\mathbf{x}^*) = 0$
  3. **Dual Feasibility:** $\lambda_i^* \geq 0$
  4. **Complementary Slackness:** $\lambda_i^* g_i(\mathbf{x}^*) = 0$
- **Gradient Descent Dynamics on Quadratics** $f(\mathbf{x}) = \frac{1}{2}\mathbf{x}^\top A\mathbf{x} - \mathbf{b}^\top\mathbf{x}$**:**
  - Convergence condition: $0 < \eta < \frac{2}{\lambda_{\max}(A)}$.
  - Optimal learning rate: $\eta^* = \frac{2}{\lambda_{\min}(A) + \lambda_{\max}(A)}$.
  - Linear convergence factor: $\left(\frac{\kappa-1}{\kappa+1}\right)$ where $\kappa = \frac{\lambda_{\max}}{\lambda_{\min}}$.