

THE UNIVERSITY OF NEW SOUTH WALES

DEPARTMENT OF STATISTICS

PRACTICE MID SESSION TEST - 2020 - Thursday, 26th March (Week 6)  
Solutions

MATH5905

Time allowed: 135 minutes

1. Let  $\mathbf{X} = (X_1, X_2, \dots, X_n)$  be i.i.d.  $\text{Poisson}(\theta)$  random variables with density function

$$f(x, \theta) = \frac{e^{-\theta} \theta^x}{x!}, \quad x = 0, 1, 2, \dots, \quad \text{and} \quad \theta > 0.$$

- a) The statistic  $T(X) = \sum_{i=1}^n X_i$  is complete and sufficient for  $\theta$ . Provide justification for why this statement is true.

**Using the property of the one-parameter exponential family, we observe that**

$$f(x, \theta) = \frac{e^{-\theta} \theta^x}{x!} = e^{-\theta} \frac{1}{x!} \exp(x \log \theta)$$

**Thus, the Poisson distribution belongs to the one-parameter exponential family. Then this implies that  $T(X) = \sum_{i=1}^n X_i$  is complete and (minimal) sufficient for  $\theta$ .**

- b) Derive the UMVUE of  $h(\theta) = e^{-k\theta}$  where  $k = 1, 2, \dots, n$  is a known integer. You must justify each step in your answer. Hint: Use the interpretation that  $P(X_1 = 0) = e^{-\theta}$  and therefore  $P(X_1 = 0, \dots, X_k = 0) = P(X_1 = 0)^k = e^{-k\theta}$ .

**Since  $T$  is sufficient and complete for  $\theta$ , we first need to find an unbiased estimator of  $h$ . Let  $W = I_{\{X_1=0, \dots, X_k=0\}}(X)$  and then using the fact that  $P(X_i = 0) = e^{-\theta}$  we see that**

$$E(W) = P(X_1 = 0, \dots, X_k = 0) = P(X_1 = 0)^k = [e^{-\theta}]^k = e^{-k\theta},$$

**which is unbiased for  $h(\theta)$ . Now, we apply the Theorem of Lehmann-Scheffe, and obtain the following**

$$\begin{aligned} \hat{\tau}(T) &= E(W|T = t) = E\left(I_{\{X_1=0, \dots, X_k=0\}} \mid \sum_{i=1}^n X_i = t\right) \\ &= P\left(X_1 = 0, \dots, X_k = 0 \mid \sum_{i=1}^n X_i = t\right) \\ &= \frac{P\left(X_1 = 0, \dots, X_k = 0, \sum_{i=1}^n X_i = t\right)}{P\left(\sum_{i=1}^n X_i = t\right)} \end{aligned}$$

Now we observe that the events in the numerator must be satisfied simultaneously to have a non-zero probability and hence this reduces to

$$\begin{aligned}
\hat{\tau}(T) &= \frac{P\left(X_1 = 0, \dots, X_k = 0, \sum_{i=k+1}^n X_i = t\right)}{P\left(\sum_{i=1}^n X_i = t\right)} \\
&= e^{-k\theta} \frac{P\left(\sum_{i=k+1}^n X_i = t\right)}{P\left(\sum_{i=1}^n X_i = t\right)} \\
&= e^{-k\theta} \frac{e^{-\theta(n-k)} (\theta(n-k))^t}{t!} \cdot \frac{t!}{e^{-n\theta} (n\theta)^t} \\
&= \left(\frac{n-k}{n}\right)^t \\
&= \left(1 - \frac{k}{n}\right)^{n\bar{X}}
\end{aligned}$$

which is the UMVUE for  $h(\theta) = e^{-k\theta}$ .

- c) Calculate the Cramer-Rao lower bound for the minimal variance of an unbiased estimator of  $h(\theta) = e^{-k\theta}$ .

**First we calculate the fisher information as follows. We have that**

$$\begin{aligned}
\log f(x) &= -\theta + x \log \theta - \log x! \\
\frac{\partial}{\partial \theta} \log f(x) &= -1 + \frac{x}{\theta} \\
\frac{\partial^2}{\partial \theta^2} \log f(x) &= -\frac{x}{\theta^2}
\end{aligned}$$

and so the Fisher information in a single sample is

$$I_{X_1}(\theta) = -E\left[\frac{\partial^2}{\partial \theta^2} \log f(x)\right] = \frac{E(X)}{\theta^2} = \frac{\theta}{\theta^2} = \frac{1}{\theta}$$

Hence the Fisher information for the whole sample is

$$I_{\mathbf{X}} = nI_{X_1}(\theta) = \frac{n}{\theta}.$$

Then notice that

$$\frac{\partial}{\partial \theta} h(\theta) = -ke^{-k\theta}.$$

Hence the Cramer-Rao lower bound is

$$\frac{\left(\frac{\partial}{\partial \theta} h(\theta)\right)^2}{I_{\mathbf{X}}(\theta)} = \frac{\theta}{n} k^2 e^{-2k\theta}$$

- d) Show that there does not exist an integer  $k$  for which the variance of the UMVUE of  $h(\theta)$  attains this bound.

**To show that the Cramer-Rao bound is attainable we will look at the score function:**

$$\begin{aligned} V(\mathbf{X}, \theta) &= -n + \frac{1}{\theta} \sum_{i=1}^n X_i \\ &= -n + \frac{n\bar{X}}{\theta} \\ &= \frac{n}{e^{-k\theta}} \left( \frac{\bar{X} e^{-k\theta}}{\theta} - e^{-k\theta} \right) \end{aligned}$$

**Therefore, no matter what integer  $k$  we choose the term  $\frac{\bar{X} e^{-k\theta}}{\theta}$  cannot be a statistic as it will still depend on  $\theta$ .**

- e) Determine the MLE  $\hat{h}$  of  $h(\theta)$ .

**The MLE for  $\theta$  is simply  $\hat{\theta} = \bar{X}$  and hence the**

$$\widehat{h(\theta)}_{\text{mle}} = h(\hat{\theta}_{\text{mle}}) = e^{-k\bar{X}}.$$

- f) Suppose that  $n = 5$ ,  $T = 10$  and  $k = 1$  compute the numerical values of the UMVUE in part (b) and the MLE in part (e). Comment on these values.

**The UMVUE is**

$$\widehat{h(\theta)}_{\text{umvue}} = \left(1 - \frac{1}{5}\right)^{5 \times 2} = 0.107$$

**The MLE is**

$$\widehat{h(\theta)}_{\text{mle}} = e^{-2} = 0.135$$

**As  $n \rightarrow \infty$  the UMVUE would converge to the MLE while for finite sample size these values are slightly different.**

- g) Consider testing  $H_0 : \theta \leq 2$  versus  $H_1 : \theta > 2$  with a 0-1 loss in Bayesian setting with the prior  $\tau(\theta) = 4\theta^2 e^{-2\theta}$ . What is your decision when  $n = 5$  and  $T = 10$ . You may use:

$$\int_0^2 x^{12} e^{-7x} dx = 0.00317$$

**Note:** The continuous random variable  $X$  has a gamma density  $f$  with parameters  $\alpha > 0$  and  $\beta > 0$  if

$$f(x; \alpha, \beta) = \frac{1}{\Gamma(\alpha)\beta^\alpha} x^{\alpha-1} e^{-x/\beta}$$

and

$$\Gamma(\alpha + 1) = \alpha\Gamma(\alpha) = \alpha!$$

First we need to compute the posterior by observing that it proportional to the likelihood times the prior. The likelihood for this same is

$$L(X, \theta) = \frac{e^{-n\theta} \theta^{\sum_{i=1}^n X_i}}{\prod_{i=2}^n X_i!} \propto e^{-5\theta} \theta^{10}$$

therefore

$$h(\theta|x) \propto 4\theta^2 e^{-2\theta} \times e^{-5\theta} \theta^{10} = \theta^{12} e^{-7\theta}$$

which implies that

$$\theta|X \sim \text{gamma}(13, \frac{1}{7}).$$

Hence we are interested in computing the posterior probability

$$\begin{aligned} P(\theta < 2|X) &= \int_0^2 \frac{1}{\Gamma(13) \left(\frac{1}{7}\right)^{13}} \theta^{12} e^{-7\theta} d\theta \\ &= 202.27 \times 0.00317 \\ &= 0.64 \end{aligned}$$

We compare this posterior probability with 0.5 since we are dealing with a 0-1 loss. Since this probability is greater than 0.5 we must NOT reject  $H_0$ .

2. Let  $X_1, X_2, \dots, X_n$  be independent random variables, with a density

$$f(x; \theta) = \begin{cases} e^{-(x-\theta)}, & x > \theta, \\ 0 & \text{else} \end{cases}$$

where  $\theta \in \mathbb{R}^1$  is an unknown parameter. Let  $T = \min\{X_1, \dots, X_n\} = X_{(1)}$  be the minimal of the  $n$  observations.

a) Show that  $T$  is a sufficient statistic for the parameter  $\theta$ .

**First calculate the likelihood as follows**

$$L(\mathbf{X}, \theta) = \prod_{i=1}^n e^{-(X_i - \theta)} I_{(-\infty, X_i)}(\theta) = \exp\left(-\sum_{i=1}^n X_i + n\theta\right) I_{(-\infty, X_{(1)})}(\theta)$$

**Therefore the likelihood can be written as  $L(X, \theta) = g(T, \theta)h(X)$  where**

$$g(T, \theta) = \exp(n\theta) I_{(-\infty, T)}(\theta) \quad \text{and} \quad h(X) = \exp\left(-\sum_{i=1}^n X_i\right)$$

**Hence,  $T = X_{(1)}$  is sufficient by the Neyman Fisher Factorization Criterion.**

b) Show that the density of  $T$  is

$$f_T(t) = \begin{cases} ne^{-n(t-\theta)}, & t > \theta, \\ 0 & \text{else} \end{cases}$$

**Hint:** You may find the CDF first by using

$$P(X_{(1)} < x) = 1 - P(X_1 > x \cap X_2 > x \cdots \cap X_n > x).$$

**First note that for  $x < \theta$  we have  $P(X_1 \geq x) = 1$  and for  $x \geq \theta$  we have,**

$$P(X_1 \geq x) = \int_x^\infty e^{-(y-\theta)} dy = \left[ -e^{-(y-\theta)} \right]_{y=x}^{y=\infty} = e^{-(x-\theta)}$$

**Hence,**

$$\begin{aligned} F_T(x, \theta) &= P(T \leq t) \\ &= 1 - P(X_1 \geq t, X_2 \geq t, \dots, X_n \geq t) \\ &= 1 - P(X_1 \geq t)^n \\ &= \begin{cases} 1 - e^{-n(t-\theta)} & \text{if } t \geq \theta \\ 0 & \text{if } t < \theta. \end{cases} \end{aligned}$$

**Then by differentiation**

$$f_T(t, \theta) = ne^{-n(t-\theta)}, \quad t \geq \theta,$$

**otherwise zero.**

- c) Find the maximum likelihood estimator of  $\theta$  and provide justification.

**The MLE for  $\theta$  is calculated by maximizing over all  $\theta$  values**

$$L(\mathbf{X}, \theta) = \exp \left( - \sum_{i=1}^n X_i + n\theta \right) I_{(-\infty, X_{(1)})}(\theta)$$

**The graph of the Likelihood function is at zero at minus infinity and then continues to increase until the value  $X_{(1)}$  and then from here stays at zero. Hence  $X_{(1)}$  must be the MLE.**

- d) Show that the MLE is a biased estimator. Hint: You might want to consider the transformation  $Y = T - \theta$  when performing the integral and then utilize the density of an exponential distribution.

**By calculating**

$$E(X_{(1)}) = \int_{\theta}^{\infty} t \cdot n e^{-n(t-\theta)} dt$$

**Now let  $y = t - \theta$  then we get**

$$\begin{aligned} E(X_{(1)}) &= n \int_0^{\infty} (y + \theta) e^{-ny} dy \\ &= \int_0^{\infty} y \cdot n e^{-ny} dy + \theta \int_0^{\infty} n e^{-ny} dy \\ &= \frac{1}{n} + \theta \\ &\neq \theta \end{aligned}$$

**and hence the MLE is a biased estimator.**

- e) Show that  $T = X_{(1)}$  is complete for  $\theta$ .

**Take any function  $g$  such that  $E_{\theta}g(T) = 0 \quad \forall \theta$ . Then**

$$\begin{aligned} 0 = E_{\theta}g(T) &= \int_{-\infty}^{\infty} g(t) f_T(t) dt \\ &= \int_{\theta}^{\infty} g(t) \cdot n e^{-n(t-\theta)} dt \\ &= n e^{n\theta} \int_{\theta}^{\infty} g(t) e^{-nt} dt \end{aligned}$$

**This implies that**

$$\int_{\theta}^{\infty} g(t) e^{-nt} dt = 0 \quad \forall \theta$$

**By taking the derivative with respect to  $\theta$  to both sides we get**

$$-g(\theta) e^{-n\theta} = 0 \quad \forall \theta$$

**but since  $e^{-n\theta} \neq 0$  we must have  $g(\theta) = 0$  for all  $\theta$ . Since  $\theta$  is just acting as a variable we have  $g(t) = 0$  for all  $t$ . Hence  $P(g(T) = 0) = 1$  for all  $\theta$  and hence  $T = X_{(1)}$  is complete for  $\theta$ .**

f) Hence determine the UMVUE of  $\theta$ .

**From part d) we know that  $W = X_{(1)} - \frac{1}{n}$  is an unbiased estimator for  $\theta$ . Thus by Lehmann-Scheffe theorem,**

$$\hat{\theta} = E(W|X_{(1)}) = X_{(1)} - \frac{1}{n}$$

**is the unique UMVUE of  $\theta$ .**