



LINEAR ALGEBRA AND ITS APPLICATIONS

UE19MA251

Unit 3. Linear Transformations and Orthogonality

Projection Matrices



The matrix P that projects onto $C(A)$ is given by

Projection matrix $P = A(A^T A)^{-1} A^T.$

Also , if P and Q are the matrices that project onto orthogonal subspaces then it is always true that $PQ = 0$ and $P + Q = I$

Unit 3. Linear Transformations and Orthogonality

Least Squares Fitting Of Data



Suppose we do a series of experiments and expect the output b to be a linear function of the input t . We look for a straight line

$$b = C + Dt$$

If there is no experimental error then two measurements of b will determine the line. But, if there is error, we minimize it by the method of least squares and find the optimal straight line.

Unit 3. Linear Transformations and Orthogonality

Least Squares Fitting Of Data



Consider the following system of equations:

$$C + Dt_1 = b_1$$

$$C + Dt_2 = b_2 \dots$$

$$C + Dt_m = b_m$$

In matrix form, $\begin{bmatrix} 1 & t_1 \\ 1 & t_2 \\ \vdots & \vdots \\ 1 & t_m \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$ or $Ax = b$

The best solution \hat{x} can be obtained by solving the normal equations.

Unit 3. Linear Transformations and Orthogonality

Problems on Least Squares Fitting Of Data



Use the method of least squares to fit the best line to the data $b = 4, 3, 1, 0$ at $t = -2, -1, 0, 2$ respectively. Find the projection of $b = (4, 3, 1, 0)$ onto the column space of A . Calculate the error vector 'e' and check that 'e' is orthogonal to the columns of A .

Solution: Let $C + Dt = b$ be the best fit straight line for the given data.

Unit 3. Linear Transformations and Orthogonality

Problems on Least Squares Fitting Of Data



Given that $b = 4, 3, 1, 0$ at $t = -2, -1, 0, 2$

$$\Rightarrow \left. \begin{array}{l} C + D(-2) = 4 \\ C + D(-1) = 3 \\ C + D(0) = 1 \\ C + D(2) = 0 \end{array} \right\} \Rightarrow \begin{bmatrix} 1 & -2 \\ 1 & -1 \\ 1 & 0 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ 1 \\ 0 \end{bmatrix}$$

$$\Rightarrow Ax = b$$

The system is inconsistent. To find least square solution \hat{x} , we have to solve normal equation

$$\text{ie } A^T A \hat{x} = A^T b$$

Unit 3. Linear Transformations and Orthogonality

Problems on Least Squares Fitting Of Data



$$\Rightarrow \begin{bmatrix} 4 & -1 \\ -1 & 9 \end{bmatrix} \hat{x} = \begin{bmatrix} 8 \\ -11 \end{bmatrix}$$

$$\hat{x} = \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 61/35 \\ -36/35 \end{pmatrix}$$

\therefore The best straight line fit for the given data is

$$b = \frac{61}{35} - \frac{36}{35}t$$

Let p be the projection of $b = (4, 3, 1, 0)$ onto $C(A)$

$$\therefore p = A\hat{x} = \frac{1}{35} \begin{pmatrix} 133 \\ 97 \\ 61 \\ -11 \end{pmatrix}$$

Unit 3. Linear Transformations and Orthogonality

Problems on Least Squares Fitting Of Data



The error vector

$$e = b - \hat{p} = \begin{pmatrix} 1/5 \\ 8/35 \\ -26/35 \\ 11/35 \end{pmatrix}$$

The error vector e is orthogonal to both the columns of A

$$\text{i.e. } e^T \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} = 0 \quad \text{and} \quad e^T \begin{pmatrix} -2 \\ -1 \\ 0 \\ 2 \end{pmatrix} = 0$$

Therefore the vector ' e ' is orthogonal to column space of A .



THANK YOU
