

Design-Connectivity-Maher-2019

Maher Yann Alain

April 2019

1 Introduction

A faire par Maher

Definitions

Let $n \in \mathbb{N}^*$.

Let Σ be an alphabet, $|\Sigma| \geq 2$. (We are especially interested in the case $\Sigma = \{A, U, G, C\}$).

Let \mathcal{F} be the set of forbidden motifs.

Let $\mathcal{L}_{\mathcal{F}}$ be the set of words in Σ^n that do not contain any motif in \mathcal{F} .

Let $H(w, w')$ be the Hamming distance between two words w, w' in Σ^n .

General problem

Input: \mathcal{F} a set of forbidden motifs, $\delta : \mathcal{L}_{\mathcal{F}} \rightarrow \mathcal{L}_{\mathcal{F}}$ a neighborhood function on $\mathcal{L}_{\mathcal{F}}$

Question: The graph $G = (\mathcal{L}_{\mathcal{F}}, \delta)$ is strongly connected.

2 Results

2.1 With the k -Hamming neighborhood

Given $k \in \mathbb{N}^*$, we define δ_k the k -Hamming neighborhood as follows:

$$\forall w \in \mathcal{L}_{\mathcal{F}}, \delta_k(w) = \{w' \in \mathcal{L}_{\mathcal{F}} \mid H(w, w') \leq k\}.$$

With $k = n$, any $w \in \mathcal{L}_{\mathcal{F}}$ can be changed into any other $w' \in \mathcal{L}_{\mathcal{F}}$ in one step. Hence $G = (\mathcal{L}_{\mathcal{F}}, \delta_n)$ is always strongly connected.

Hence with the k -Hamming neighborhood a variant of the general problem can be considered:

Input: \mathcal{F} a set of forbidden motifs

Question: minimal $k \in \mathbb{N}^*$ such that the graph $G = (\mathcal{L}_{\mathcal{F}}, \delta_k)$ is strongly connected.

2.1.1 One motif

Consider the case where \mathcal{F} contains a single motif: $\mathcal{F} = \{f\}$.

Then $k = 1$ is sufficient to guarantee strong connectivity.

Result 1. $\forall f \in \Sigma^+, G = (\mathcal{L}_{\{f\}}, \delta_1)$ is strongly connected.

Proof. Let w and w' be two words in $\mathcal{L}_{\{f\}}$ (of length n).

As $f \neq \epsilon$, it can be written letter by letter as follows: $f = f_1 \dots f_{|f|}$.

Since $|\Sigma| \geq 2$, let $a \in \Sigma$ such that $a \neq f_1$.

We show that there is a path from w to a^n .

To do so, from left to right we will replace each letter in w by a (or keep it the same if already an a).

Formally, from $w = w_1 \dots w_n$ we define the sequence $(u_i)_{0 \leq i \leq n}$ of intermediate words:

$$\forall 0 \leq i \leq n, u_i = a^i w_{i+1} \dots w_n.$$

Then:

- $\forall 0 \leq i \leq n-1, H(u_i, u_{i+1}) \leq 1$,
- for every i in $[1..n]$ we must prove that u_i is in $\mathcal{L}_{\{f\}}$. By contradiction, suppose that f appears in a u_i . Let j the position in u_i of the leftmost letter of this occurrence of f .
 - if $j \leq i$: then the leftmost letter of f would be a , which is not by definition of a .
 - if $j > i$: then this occurrence of f would be a factor of w , which it cannot be since $w \in \mathcal{L}_{\{f\}}$.

Contradiction. Hence every u_i is in $\mathcal{L}_{\{f\}}$.

This proves that there is a path from w to a^n with δ_1 as the neighborhood function.

The same can be done to obtain a path from w' to a^n .

Finally, since δ_1 is symmetric this gives a path from w to w' and vice-versa. □