

# Discrete Structures

Set Theory

# Introduction to Set Theory

- ◆ A *set* is a structure, representing an unordered collection (group, plurality) of zero or more distinct (different) objects.
- ◆ Set theory deals with operations between, relations among, and statements about sets.

# Basic notations for sets

- ◆ For sets, we'll use variables  $S, T, U, \dots$
- ◆ We can denote a set  $S$  in writing by listing all of its elements in curly braces:
  - $\{a, b, c\}$  is the set of whatever 3 objects are denoted by  $a, b, c$ .
- ◆ *Set builder notation*: For any proposition  $P(x)$  over any universe of discourse,  $\{x | P(x)\}$  is *the set of all  $x$  such that  $P(x)$* .  
e.g.,  $\{x | x \text{ is an integer where } x > 0 \text{ and } x < 5\}$

# Basic properties of sets

- ◆ Sets are inherently unordered:
  - No matter what objects  $a$ ,  $b$ , and  $c$  denote,  
 $\{a, b, c\} = \{a, c, b\} = \{b, a, c\} =$   
 $\{b, c, a\} = \{c, a, b\} = \{c, b, a\}.$
- ◆ All elements are distinct (unequal);  
multiple listings make no difference!
  - $\{a, b, c\} = \{a, a, b, a, b, c, c, c, c\}.$
  - This set contains at most 3 elements!


# Definition of Set Equality

- ◆ Two sets are declared to be equal *if and only if* they contain exactly the same elements.
- ◆ In particular, it does not matter *how the set is defined or denoted*.
- ◆ For example: The set  $\{1, 2, 3, 4\} =$   
 $\{x \mid x \text{ is an integer where } x > 0 \text{ and } x < 5\} =$   
 $\{x \mid x \text{ is a positive integer whose square}$   
 $\text{is } > 0 \text{ and } < 25\}$

# Infinite Sets

- ◆ Conceptually, sets may be *infinite* (i.e., not *finite*, without end, unending).
- ◆ Symbols for some special infinite sets:  
 $\mathbb{N} = \{0, 1, 2, \dots\}$  The natural numbers.  
 $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  The integers.  
 $\mathbb{R}$  = The "real" numbers, such as  
374.1828471929498181917281943125...
- ◆ Infinite sets come in different sizes!

# Venn Diagrams



# Basic Set Relations: Member of

- ◆  $x \in S$  ("x is in S") is the proposition that object  $x$  is an *element* or *member* of set  $S$ .
  - e.g.  $3 \in \mathbb{N}$ , "a"  $\in \{x \mid x \text{ is a letter of the alphabet}\}$
- ◆ Can define set equality in terms of  $\in$  relation:  
$$\forall S, T: S = T \leftrightarrow (\forall x: x \in S \leftrightarrow x \in T)$$

"Two sets are equal iff they have all the same members."
- ◆  $x \notin S \equiv \neg(x \in S)$       "x is not in S"



# The Empty Set

- ◆  $\emptyset$  ("null", "the empty set") is the unique set that contains no elements whatsoever.
- ◆  $\emptyset = \{\} = \{x/\text{False}\}$
- ◆ No matter the domain of discourse, we have the axiom


$$\neg \exists x: x \in \emptyset.$$

# Subset and Superset Relations

- ◆  $S \subseteq T$  (" $S$  is a subset of  $T$ ") means that every element of  $S$  is also an element of  $T$ .
- ◆  $S \subseteq T \Leftrightarrow \forall x (x \in S \rightarrow x \in T)$
- ◆  $\emptyset \subseteq S, S \subseteq S$ .
- ◆  $S \supseteq T$  (" $S$  is a superset of  $T$ ") means  $T \subseteq S$ .
- ◆ Note  $S = T \Leftrightarrow S \subseteq T \wedge S \supseteq T$ .
- ◆  $S \not\subseteq T$  means  $\neg(S \subseteq T)$ , i.e.  $\exists x (x \in S \wedge x \notin T)$

# Proper (Strict) Subsets & Supersets

- ◆  $S \subset T$  (" $S$  is a proper subset of  $T$ ") means that  $S \subseteq T$  but  $T \not\subseteq S$ . Similar for  $S \supset T$ .



Example:

$$\{1,2\} \subset \{1,2,3\}$$

Venn Diagram equivalent of  $S \subset T$

# Sets Are Objects, Too!

- ◆ The objects that are elements of a set may *themselves* be sets.
- ◆ E.g. let  $S = \{x \mid x \subseteq \{1, 2, 3\}\}$   
then  $S = \{\emptyset,$   
           $\{1\}, \{2\}, \{3\},$   
           $\{1, 2\}, \{1, 3\}, \{2, 3\},$   
           $\{1, 2, 3\}\}$
- ◆ Note that  $1 \neq \{1\} \neq \{\{1\}\} !!!!$

# Cardinality and Finiteness

- ◆  $|S|$  (read “the *cardinality* of  $S$ ”) is a measure of how many different elements  $S$  has.
- ◆ E.g.,  $|\emptyset|=0$ ,  $|\{1,2,3\}| = 3$ ,  $|\{a,b\}| = 2$ ,  
 $|\{\{1,2,3\},\{4,5\}\}| = \underline{2}$
- ◆ We say  $S$  is *infinite* if it is not *finite*.
- ◆ What are some infinite sets we've seen?

**N Z R**

# The *Power Set* Operation

- ◆ The *power set*  $P(S)$  of a set  $S$  is the set of all subsets of  $S$ .  $P(S) = \{x \mid x \subseteq S\}$ .
- ◆ E.g.  $P(\{a,b\}) = \{\emptyset, \{a\}, \{b\}, \{a,b\}\}$ .
- ◆ Sometimes  $P(S)$  is written  $2^S$ .  
Note that for finite  $S$ ,  $|P(S)| = 2^{|S|}$ .
- ◆ It turns out that  $|P(\mathbb{N})| > |\mathbb{N}|$ .  
*There are different sizes of infinite sets!*

# Ordered $n$ -tuples

- ◆ For  $n \in \mathbf{N}$ , an *ordered  $n$ -tuple* or a sequence of length  $n$  is written  $(a_1, a_2, \dots, a_n)$ . The *first* element is  $a_1$ , *etc.*
- ◆ These are like sets, except that duplicates matter, and the order makes a difference.
- ◆ Note  $(1, 2) \neq (2, 1) \neq (2, 1, 1)$ .
- ◆ Empty sequence, singlets, pairs, triples, quadruples, quintuples, ...,  $n$ -tuples.

# Cartesian Products of Sets

- ◆ For sets  $A, B$ , their *Cartesian product*  $A \times B := \{(a, b) \mid a \in A \wedge b \in B\}$ .
- ◆ E.g.  $\{a, b\} \times \{1, 2\} = \{(a, 1), (a, 2), (b, 1), (b, 2)\}$
- ◆ Note that for finite  $A, B$ ,  $|A \times B| = |A| |B|$ .
- ◆ Note that the Cartesian product is *not* commutative:  $\neg \forall A, B: A \times B = B \times A$ .
- ◆ Extends to  $A_1 \times A_2 \times \dots \times A_n \dots$




# The Union Operator

- ◆ For sets  $A, B$ , their *union*  $A \cup B$  is the set containing all elements that are either in  $A$ , or (" $\vee$ ") in  $B$  (or, of course, in both).
- ◆ Formally,  $\forall A, B: A \cup B = \{x \mid x \in A \vee x \in B\}$ .
- ◆ Note that  $A \cup B$  contains all the elements of  $A$  and it contains all the elements of  $B$ :  
$$\forall A, B: (A \cup B \supseteq A) \wedge (A \cup B \supseteq B)$$

# Union Examples

◆  $\{a,b,c\} \cup \{2,3\} = \{a,b,c,2,3\}$

◆  $\{2,3,5\} \cup \{3,5,7\} = \{2,3,5,3,5,7\} = \{2,3,5,7\}$




# The Intersection Operator

- ◆ For sets  $A, B$ , their *intersection*  $A \cap B$  is the set containing all elements that are simultaneously in  $A$  and (" $\wedge$ ") in  $B$ .
- ◆ Formally,  $\forall A, B: A \cap B \equiv \{x \mid x \in A \wedge x \in B\}$ .
- ◆ Note that  $A \cap B$  is a subset of  $A$  and it is a subset of  $B$ :  
 $\forall A, B: (A \cap B \subseteq A) \wedge (A \cap B \subseteq B)$

# Intersection Examples

◆  $\{a,b,c\} \cap \{2,3\} = \underline{\emptyset}$

◆  $\{2,4,6\} \cap \{3,4,5\} = \underline{\{4\}}$



# Disjointedness

- ◆ Two sets  $A, B$  are called *disjoint* (i.e., unjoined) iff their intersection is empty. ( $A \cap B = \emptyset$ )
- ◆ Example: the set of even integers is disjoint with the set of odd integers.

# Inclusion-Exclusion Principle

◆ How many elements are in  $A \cup B$ ?

$$|A \cup B| = |A| + |B| - |A \cap B|$$

◆ Example:


$$\{2,3,5\} \cup \{3,5,7\} = \{2,3,5,3,5,7\} = \{2,3,5,7\}$$

# Set Difference

- ◆ For sets  $A, B$ , the *difference of  $A$  and  $B$* , written  $A - B$ , is the set of all elements that are in  $A$  but not  $B$ .
- ◆  $A - B \equiv \{x \mid x \in A \wedge x \notin B\}$   
 $= \{x \mid \neg(x \in A \rightarrow x \in B)\}$
- ◆ Also called:  
The complement of  $B$  with respect to  $A$ .

# Set Difference - Venn Diagram

◆  $A - B$  is what's left after  $B$  "takes a bite out of  $A$ "





# Set Complements


- ◆ The *universe of discourse* can itself be considered a set, call it  $U$ .
- ◆ The *complement* of  $A$ , written  $\overline{A}$ , is the complement of  $A$  w.r.t.  $U$ , i.e., it is  $U - A$ .
- ◆ E.g., If  $U = \mathbb{N}$ ,

$$\overline{\{3,5\}} = \{0,1,2,4,6,7,\dots\}$$

# More on Set Complements

- ◆ An equivalent definition, when  $U$  is clear:

$$\bar{A} = \{x \mid x \notin A\}$$



# Set Identities

- ◆ Identity:  $A \cup \emptyset = A$   $A \cap U = A$
- ◆ Domination:  $A \cup U = U$   $A \cap \emptyset = \emptyset$
- ◆ Idempotent:  $A \cup A = A = A \cap A$
- ◆ Double complement:
- ◆ Commutative:  $A \cup B = B \cup A$   $A \cap B = B \cap A$
- ◆ Associative:  $A \cup (B \cup C) = (A \cup B) \cup C$   
 $A \cap (B \cap C) = \underline{\underline{(A \cap B)}} \cap C$   
 $\overline{(\overline{A})} = A$

# DeMorgan's Law for Sets

- ◆ Exactly analogous to (and derivable from) DeMorgan's Law for propositions.

$$\overline{A \cup B} = \bar{A} \cap \bar{B}$$

$$\overline{A \cap B} = \bar{A} \cup \bar{B}$$

# Proving Set Identities

To prove statements about sets, of the form  $E_1 = E_2$  (where  $E$ s are set expressions), here are three useful techniques:

- ◆ Prove  $E_1 \subseteq E_2$  and  $E_2 \subseteq E_1$  separately.
- ◆ Use logical equivalences.
- ◆ Use a *membership table*.

# Method 1: Mutual subsets

Example: Show  $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$ .

◆ Show  $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$ .

- Assume  $x \in A \cap (B \cup C)$ , & show  $x \in (A \cap B) \cup (A \cap C)$ .
- We know that  $x \in A$ , and either  $x \in B$  or  $x \in C$ .

- ◆ Case 1:  $x \in B$ . Then  $x \in A \cap B$ , so  $x \in (A \cap B) \cup (A \cap C)$ .

- ◆ Case 2:  $x \in C$ . Then  $x \in A \cap C$ , so  $x \in (A \cap B) \cup (A \cap C)$ .

- Therefore,  $x \in (A \cap B) \cup (A \cap C)$ .

- Therefore,  $A \cap (B \cup C) \subseteq (A \cap B) \cup (A \cap C)$ .

◆ Show  $(A \cap B) \cup (A \cap C) \subseteq A \cap (B \cup C)$ . ...

# Method 3: Membership Tables

- ◆ Just like truth tables for propositional logic.
- ◆ Columns for different set expressions.
- ◆ Rows for all combinations of memberships in constituent sets.
- ◆ Use "1" to indicate membership in the derived set, "0" for non-membership.
- ◆ Prove equivalence with identical columns.

# Membership Table Example

Prove  $(A \cup B) - B = A - B$ .

$A$	$B$	$A \cup B$	$(A \cup B) - B$	$A - B$
0	0	0	0	0
0	1	1	0	0
1	0	1	1	1
1	1	1	0	0



# Membership Table Exercise

Prove  $(A \cup B) - C = (A - C) \cup (B - C)$ .

$A$	$B$	$C$	$A \cup B$	$(A \cup B) - C$	$A - C$	$B - C$	$(A - C) \cup (B - C)$
0	0	0					
0	0	1					
0	1	0					
0	1	1					
1	0	0					
1	0	1					
1	1	0					
1	1	1					

# Generalized Union

◆ Binary union operator:  $A \cup B$

◆  $n$ -ary union:

$A \cup A_2 \cup \dots \cup A_n \equiv (((A_1 \cup A_2) \cup \dots) \cup A_n)$   
(grouping & order is irrelevant)

◆ "Big U" notation:  $\bigcup_{i=1}^n A_i$

◆ Or for infinite sets of sets:  $\bigcup_{A \in X} A$

# Generalized Intersection

◆ Binary intersection operator:  $A \cap B$

◆  $n$ -ary intersection:

$A_1 \cap A_2 \cap \dots \cap A_n \equiv ((\dots((A_1 \cap A_2) \cap \dots) \cap A_n)$   
(grouping & order is irrelevant)

◆ "Big Arch" notation:  $\bigcap_{i=1}^n A_i$

◆ Or for infinite sets of sets:  $\bigcap_{A \in X} A$

# References

- *Sections 2.1 and 2.2 of the text book "Discrete Mathematics and its Applications" by Rosen, 6<sup>th</sup> edition.*
- The original slides were prepared by Bebis