
Status and plans towards the government of the state of t





Cristina Botta

for the ttH leptonic working group





Target

- ttH->leptons analysis PAS with 2015 data for Moriond 2016
 - from latest LHC news: we expect between 2.3 and 3.4 fb⁻¹
 - → if similar S/B, σ x 4, L/10 : 1.6 x (8 TeV analysis sensitivity)
 - \rightarrow expected uncertainty on $\mu(ttH)\sim1.9$ (we were at 1.2 @ 8 TeV)
 - ⇒given that we were measuring µ=4 @ 8 TeV a measurement with a 1.9 uncertainty may be of interest
- First important milestone:
 - Full analysis review at the HWW MiniWorkshop 23/24 November
 - AtoZ MC strategy
 - Control of object performances in data and scale factors
 - Data driven bkg estimates in place



Samples

- 25ns samples from the 1B Spring15 campaign (waiting for Spring15MiniAODv2)
 - for final signal extraction
 - ttHJetToNonbb_M125_13TeV_amcatnloFXFX_madspin_pythia8_mWCutfix/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v1/MINIA0DSIM
 - ttHJetToNonbb_M125_13TeV_amcatnloFXFX_madspin_pythia8_mWCutfix/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9_ext1-v1/MINIA0DSIM
 - TTZToLLNuNu_M-10_TuneCUETP8M1_13TeV-amcatnlo-pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v1/MINIA0DSIM
 - the associated M-1to10 is in preparation
 - TTWJetsToLNu_TuneCUETP8M1_13TeV-amcatnloFXFX-madspin-pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v1/MINIA0DSIM
 - TTGJets_TuneCUETP8M1_13TeV-amcatnloFXFX-madspin-pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v1/MINIA0DSIM
 - WZTo3LNu TuneCUETP8M1 13TeV-powheg-pythia8/RunIISpring15DR74-Asympt25ns MCRUN2 74 V9-v1/MINIAODSIM
 - or the Madgraph one when it will become available
 - ZZTo4L_13TeV_powheg_pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v1/MINIA0DSIM
 - or a new Madgraph one (the current version is bugged)
 - rares: WWsamesign, Tribosons
 - for lepton MVA and final BDT training
 - ttHToNonbb_M125_13TeV_powheg_pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v2/MINIA0DSIM
 - TT_TuneCUETP8M1_13TeV-powheg-pythia8/RunIISpring15DR74-Asympt25ns_MCRUN2_74_V9-v2/MINIA0DSIM
 - or the semileptonic Madgraph one
 - ttZ, ttW madgraph LO in preparation
 - for bkg data driven estimates
 - all QCD samples MuEnriched and EMEnriched for fake-rate estimate
 - TTJets (maybe the semileptonic Madgraph one), Wjets for fake-rate estimate
 - DYJets for Charge-Flip probability estimate





Triggers

- As for in Frozen 25ns Menu: /frozen/2015/25ns14e33/v4.2/HLT/V1
 - Double Electron Iso: 17/12 instead of 23/12 (available in MC)
 - Single Ele Iso: new Ele23_WPLoose_Gsf instead of 32eta2p1 (but not available in MC)
 - lowest threshold un-prescaled present both in data and MC: Ele27_WP85
 - Double Mu Iso: 17/8 (available in MC)
 - Single Mu Iso: new IsoMu18 instead of 24eta2p1 (but not available in MC)
 - lowest threshold un-prescaled present both in data and MC: IsoMu20





Lepton flavour & p_T categories

8 TeV analysis

- ee (20,20) from DoubleEle triggers
- μμ (20,20) from DoubleMu triggers
- eµ (20,20) from MuEG triggers
- 3I (20,10,5/7) from Double/Triple Lepton triggers
- 41 (20,10,5/7) from Double/Triple Lepton triggers



24.09.15



Lepton flavour & p_T categories

- ee (20,15) from Double and Single Ele triggers + (30,7-15) from Single Ele
 - to be splitted in 0 or >= 1 hadronic taus
- μμ (20,15) from Double and Single Mu triggers + (30,5-15) from Single Mu
 - to be splitted in 0 or >= 1 hadronic taus
- eμ (20,15) from MuEG and SingleMu/Ele triggers + (30,5/7-15) from Single Mu/Ele
 - to be splitted in 0 or >= 1 hadronic taus
- 31 (20,10,5/7) from Double/Triple Lepton triggers
- 41 (20,10,5/7) from Double/Triple Lepton triggers

Comments

- additional low sub-leading p_T category bring ~10% signal efficiency in each channel but need to check S/B (Geoff, Charlie to finalize the study with new lepton mva)
- is it worth it to add the tau categories with 2/3 fb⁻¹? (Jan to answer this question)

13 TeV analysis





Recall: Lepton selection

- Loose: for vetoing additional leptons in each categories
 - Common: |dxy| < 500um, |dz| < 1 mm, MiniRellso (dR=0.3, rho*EA) < 0.4, SIP_{3D} < 8
 - Muons: Loose ID
 - Electrons: VLoose WP of Phys14 non-triggering MVA ID, lost hits ≤ 1
- Tight: for signal region selections
 - Common: lepMVAttH < 0.6-0.8
 - Muons: Medium ID
 - for ss-dilepton only: track p_T / track p_T error < 0.2
 - Electrons: passConversionsVeto, lost hits = 0
 - for ss-dilepton only: isGsfCtfScPixChargeConsistent
- Fakable-object: for fake-rate denominator
 - Same as tight but without lepMVAttH requirement and without Medium Muon ID



Jet (Lepton-subtracted)

pTrel v1

Recall: lepMVAttH

input variables will be using in Runll

Isolation

- PF miniRellso, charged had. (R=0.3)
- **PF miniRellso**, **neutral** had. & photon (R=0.3, with EA*rho PU corrections, scaled with (0.3/R)^2)

Vertexing

- 3D IP significance (SIP_{3D})
- 2D IP | dxy | and | dz |
- Lepton's closest jet (leptons are not removed from jet collection)
 - p_T(l)/p_T(jet)
 - Lepton's p_T^{rel} wrt jet
 - jet CSV b-tag



- (µ)Segment compatibility/
- (e)POG non-triggering MVA Electron Id discriminator





Lepton ID on going studies

Effective Ares for minilso

- for the moment still using those extracted from PHYS14 MC
- we are deriving those for Spring15 25ns samples, in collaboration with POGs
 - Marco will send final results next week

Electron MVA ID

- for the moment still using PHYS14 non-triggering training
- will have to move to EGamma Spring15 25ns non-triggering training
 - available in miniAODv2 (~1/2 weeks)
- VLoose WP being extracted

Commissioning of input variables

- a lot of results with 50ns and 25ns data already reported at the HWW meeting
- new interesting studies on going on "how to apply JEC to the jet associated to the lepton" while computing the lepton-jet associated variables
 - JEC lep-aware proposal being discussed with JET/MET, SUSY PAG etc...
 - will be reported next week in our meeting by Giovanni/Marco





Lepton ID on going studies

Lepton MVA new training

- with new electron MVA preselection, new electron MVA training as input variable (when available)
- with new lep-aware JEC for lepton-jet related variables
- with weighted-lepton p_T as input variable (to partially smooth the difference between the signal and background leptons p_T spectrum)
- new working point being derived taking into account:
 - that we might want to be looser on low p_T leptons
 - the additional separation power between ttH and tt we can achieve with the final BDT
- Giovanni and Francesco are working on it





Data driven backgrounds

- Fakes: tight-to-loose method as last year (Marco, Giovanni)
 - with cone-corrected p_T improvement to reduce differences in fake-rate estimate between QCD and ttbar events
 - to be checked if current trigger strategy (~lepton p_T cuts) allows for enough sideband in the fake rate application
 - to be checked with 13 TeV data and MC the tight-to-loose ratio flavour dependence to decide how to approach the b-tagging categorisation for the signal-extraction
 - two fake-rate from two different measurement region are needed?
- Charge flip: same method as last year (who?)
 - or possible improvement:
 - extract data/MC SF for charge-flip probabilty in coarse lepton p_T bins
 - then interpolate and extrapolate the p_T dependence using MC
- WZ: same method as last year (who?)





Signal extraction

- BDT as in 8 TeV analysis with following improvements
 - might add new input variables
 - angular variables proposed by Milos (to be studied also with Spring15)
 - reconstruction of hadronic/leptonic top mass (Giovanni is working on this)
 - kinematic fits (Benjamin interested in this)
 - new categorisation
 - ⇒ b-tight and b-loose
 - ⇒ split 31 category in 2-3/4+ jets to better separate ttZ/ttW from ttH
 - add 3j category in the 2lss final state to control ttW from data
 - add the 3I category with inverted Z veto to control ttZ from data
 - 2D approach (against tt, against ttW/Z) as tested with Phys14 samples by Predrag and Milos

MEM

- what is the current status?
- what is expected for mid November?





Additional topics/proposals to discuss?