

# STAT 111

## Recitation 1

Mo Huang

Email: [mohuang@wharton.upenn.edu](mailto:mohuang@wharton.upenn.edu)

Office Hours: Wednesdays 3:00 - 4:00 pm, JMHH F96

Slides (adapted from Gemma Moran): [github.com/mohuangx/STAT111-Fall2018](https://github.com/mohuangx/STAT111-Fall2018)

September 7, 2018

# Logistics

- ▶ Section 201: 11:00 am - 11:50 am
- ▶ Section 202: 12:00 pm - 12:50 pm
  
- ▶ Every Friday, I will:
  1. Collect the homework for that week;
  2. Give you your graded homework; and
  3. Give you the homework for the week after (also posted on Canvas).
  
- ▶ If you do not collect your homework, it will be available in the STAT 111 box in the Statistics Department (4th Floor, JMHH).
  
- ▶ If you know in advance that you will not be able to attend a recitation, give your homework to Professor Ewens on Thursdays or put it in my mailbox (at entrance to the Statistics Department)
  
- ▶ Questions about course materials should be asked during office hours or in an email to Professor Ewens.

# Statistics and Probability


- ▶ **Probability theory** is *deductive* (or “top-down logic”). We start with a theory and then consider the implications resulting from that theory.
- ▶ **Statistics** is *inductive* (or “bottom-up logic”). We start with finite observations and then attempt to make objective statements about the world from a sample.
- ▶ Statistical methods use probability theory to draw conclusions from observed data.

# Statistics and Probability: Example

- ▶ We currently cure an illness with some medicine (“current medicine”) which cures 80% of patients.
- ▶ A new medicine is proposed (“new medicine”) which cures 1,643 people out of 2,000.

**Is the new medicine more effective?**

Probability theory




Statistics


# Probability: Axioms

- ▶  $0 \leq P(A) \leq 1$ , where  $A$  is any event.
- ▶  $P(A^C) = 1 - P(A)$  where  $A^C$  is the complement of  $A$ .
- ▶  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

**Intersection**      $P(A \cap B)$



**Union**      $P(A \cup B)$



# Probability: Independence

- ▶  $A$  and  $B$  are independent if and only if  $P(A \cap B) = P(A)P(B)$
- ▶ Conditional probability:  $P(A|B) = \frac{P(A \cap B)}{P(B)}$

Suppose  $A$  and  $B$  are independent. Then

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

So we have two definitions of independence:

1.  $P(A \cap B) = P(A)P(B)$
2.  $P(A|B) = P(A)$

**Note:** This year, conditional probability is not on the syllabus but it's still an interesting and useful concept!

## Probability: Questions

Q1: A smoke alarm consists of two parts,  $A$  and  $B$ . If there is smoke, part  $A$  will detect it with probability 0.96 and part  $B$  will detect it with probability 0.98. The probability that they both detect it is 0.95. What is the probability that smoke will not be detected?

A1: Let  $A$  be the event that part  $A$  detects smoke and let  $B$  be the event that part  $B$  detects smoke. What is the event that smoke is not detected?

$A \cup B = \text{Smoke is detected} \rightarrow [A \cup B]^C = \text{Smoke is not detected}$

$$\begin{aligned}P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\&= 0.96 + 0.98 - 0.95 \\&= 0.99\end{aligned}$$

$$\begin{aligned}P([A \cup B]^C) &= 1 - P(A \cup B) \\&= 0.01\end{aligned}$$

## Probability: Questions

Q2: A six-sided die is biased such that an odd number is twice as likely to occur as an even number. That is, the probability of an odd number is  $2/9$  and an even number is  $1/9$ . Let  $A$  be the event that an even number occurs. Let  $B$  be the event that a number greater than or equal to 4 occurs. Find  $P(A)$ ,  $P(B)$ ,  $P(A \cup B)$  and  $P(A \cap B)$ .

A2:

$$\begin{aligned}P(A) &= P(2, 4 \text{ or } 6) \\&= 1/9 + 1/9 + 1/9 \\&= 3/9\end{aligned}$$

$$\begin{aligned}P(A \cap B) &= P(4 \text{ or } 6) \\&= 1/9 + 1/9 \\&= 2/9\end{aligned}$$

$$\begin{aligned}P(B) &= P(4, 5 \text{ or } 6) \\&= 1/9 + 2/9 + 1/9 \\&= 4/9\end{aligned}$$

$$\begin{aligned}P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\&= 3/9 + 4/9 - 2/9 \\&= 5/9\end{aligned}$$



## Probability: Questions

A2:  $P(A) = 3/9$ ,  $P(B) = 4/9$ ,  $P(A \cap B) = 2/9$ ,  $P(A \cup B) = 5/9$

Q3: Following Q2, find  $P(A|B)$ .

A3:

$$\begin{aligned} P(A|B) &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{2/9}{4/9} \\ &= 1/2 \end{aligned}$$

Q4: Are  $A$  and  $B$  independent?

A4: No, because  $P(A|B) \neq P(A)$ . (Recall  $P(A) = 1/3$ ).

## Probability: Questions

Q5: An unfair coin ( $P(H) = 0.4$ ) was flipped twice. Given that on at least one flip a head occurred, what is the probability that a head occurred on both flips?

A5: We want to find  $P(HH|\text{at least one } H)$ .

$$\begin{aligned}P(\text{at least one } H) &= P(HH) + P(HT) + P(TH) \\&= (0.4)^2 + (0.4)(0.6) + (0.6)(0.4) \\&= 0.64.\end{aligned}$$

$$\begin{aligned}P(HH|\text{at least one } H) &= \frac{P(HH \cap \text{at least one } H)}{P(\text{at least one } H)} \\&= \frac{P(HH)}{P(\text{at least one } H)} \\&= \frac{(0.4)^2}{0.64} \\&= 0.25\end{aligned}$$

# Parameters ( $\theta$ ), Random Variables ( $X$ ), and Data ( $x$ )

- ▶ A **parameter** represents some underlying numerical constant of a phenomenon. Represented by Greek letters.
- ▶ A **random variable** is a numerical outcome of interest in a future experiment. Can be modeled by a probability **distribution** which depends on the parameter. Represented by capital letters.
- ▶ **Data** is the realization or observed value of a random variable after performing the experiment. Represented by lower-case letters.

Fair coin-flipping example:

- ▶ A **parameter**  $\theta$  is the underlying probability of obtaining a head.  $\theta = 0.5$ .
- ▶ A **random variable**  $X$  is the number of heads obtained in 10 tosses. Can be modeled by a binomial distribution dependent on  $\theta$ .
- ▶ **Data**  $x = 6$  is observing 6 heads in 10 tosses, a realization of  $X$ .

# Random Variables

- ▶ Two types of random variables: **discrete** and continuous.
- ▶ A **discrete random variable** is a random variable that can only take on a countable set of numbers.
- ▶ The **probability distribution** of a discrete random variable is the range of values it can take *and* the probabilities of these values.
- ▶ For example:
  - ▶ Let  $X$  be the number of heads I toss if I toss a fair coin 3 times.

$x$	0	1	2	3
$P(X = x)$	0.125	0.375	0.375	0.125

**Table:** Probability distribution of  $X$ .

# Random Variables: Mean

- ▶ The **mean of a random variable** (expected value) is the long-run average of realizations of the random variable over repeated experiments.
- ▶ This mean of random variable  $X$  is different from the sample mean, which is the average of a *finite* number of observations ( $x$ ).
- ▶ Let  $X$  be a discrete random variable that can take values  $\{v_1, v_2, \dots, v_k\}$ . Then the mean of  $X$  is given by:

$$\begin{aligned}\mu &= \sum_{i=1}^k v_i P(X = v_i) \\ &= v_1 P(X = v_1) + v_2 P(X = v_2) + \dots + v_k P(X = v_k).\end{aligned}$$

**Note:** We use  $\mu$  to denote the mean of a random variable.

**Note:** Can think of it as a weighted average, weighted by probability.

# The Binomial Distribution

- ▶ The binomial distribution arises if:
  1. We plan to conduct a fixed number of experiments. We denote the number of experiments as  $n$ .
  2. In each experiment, there are two outcomes: “success” or “failure”.
  3. The experiments are independent.
  4. The probability of a success is the same for each experiment.
  
- ▶ For example:
  1. I plan to toss a coin  $n$  times.
  2. I can toss either a head or a tail.
  3. Each coin toss is independent of the next - it doesn't matter whether I get a head or a tail on the previous toss.
  4. The probability of getting a head is the same for each toss.

# The Binomial Distribution

- ▶ Let  $X$  be a binomial random variable where  $\theta = P(\text{success})$  and there are  $n$  experiments. Then the probability distribution of  $X$  is given by:

$$P(X = x) = \binom{n}{x} \theta^x (1 - \theta)^{n-x}, \quad \text{for } x = 0, 1, 2, \dots, n.$$

- ▶  $\theta$  is called a **parameter**: a constant whose value may be known or unknown.
- ▶  $\binom{n}{x}$  is said as “ $n$  choose  $x$ ”: it is the number of ways  $x$  successes can occur in  $n$  experiments.

**Note:**  $X \sim \mathcal{B}(n, \theta)$  means “ $X$  is a binomial random variable with  $n$  experiments and probability of success  $\theta$ .”

# The Binomial Distribution: Mean

- ▶ For the binomial distribution, we have a simpler formula for the mean: if  $X \sim \mathcal{B}(n, \theta)$ , then the mean of  $X$  is

$$\mu = n\theta.$$

- ▶ For example:
  - ▶ On the previous slide, we calculated the mean of a binomial random variable with  $n = 3$  and  $\theta = 0.5$  to be 1.5. This is exactly  $n\theta$ .

**Note:** This formula is *only* for the binomial distribution.