

# STAT 111

## Recitation 5

Mo Huang

Email: [mohuang@wharton.upenn.edu](mailto:mohuang@wharton.upenn.edu)

Office Hours: Wednesdays 3:00 - 4:00 pm, JMHH F96

Slides: [github.com/mohuangx/STAT111-Spring2019](https://github.com/mohuangx/STAT111-Spring2019)

February 22, 2019

## Two-Standard-Deviation Rule

- ▶ From the chart:

$$P(Z < -1.96) = 0.025, \quad P(Z > 1.96) = 0.025.$$

# Two-Standard-Deviation Rule

- ▶ From the chart:

$$P(Z < -1.96) = 0.025, \quad P(Z > 1.96) = 0.025.$$

- ▶ Then:

$$P(-1.96 < Z < 1.96) = 0.95.$$

# Two-Standard-Deviation Rule

- ▶ From the chart:

$$P(Z < -1.96) = 0.025, \quad P(Z > 1.96) = 0.025.$$

- ▶ Then:

$$P(-1.96 < Z < 1.96) = 0.95.$$

- ▶ Approximate  $1.96 \approx 2$  and “unstandardize”:

$$P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = 0.95$$

# Two-Standard-Deviation Rule

- ▶ From the chart:

$$P(Z < -1.96) = 0.025, \quad P(Z > 1.96) = 0.025.$$

- ▶ Then:

$$P(-1.96 < Z < 1.96) = 0.95.$$

- ▶ Approximate  $1.96 \approx 2$  and “unstandardize”:

$$P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = 0.95$$

$\Rightarrow$

$$P(\mu - 2\sigma < X < \mu + 2\sigma) = 0.95.$$

# Two-Standard-Deviation Rule

- ▶ From the chart:

$$P(Z < -1.96) = 0.025, \quad P(Z > 1.96) = 0.025.$$

- ▶ Then:

$$P(-1.96 < Z < 1.96) = 0.95.$$

- ▶ Approximate  $1.96 \approx 2$  and “unstandardize”:

$$P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = 0.95$$

⇒

$$P(\mu - 2\sigma < X < \mu + 2\sigma) = 0.95.$$

- ▶ The probability that a normal random variable is within 2 standard deviations of the mean is 95%.

## Normal Distribution: Sums and Averages

- Let  $X_1, \dots, X_n$  be independent and *normally distributed*. Let

$$T_n = X_1 + \dots + X_n, \quad \bar{X} = \frac{X_1 + \dots + X_n}{n}, \quad D = X_2 - X_1.$$

## Normal Distribution: Sums and Averages

- ▶ Let  $X_1, \dots, X_n$  be independent and *normally distributed*. Let

$$T_n = X_1 + \dots + X_n, \quad \bar{X} = \frac{X_1 + \dots + X_n}{n}, \quad D = X_2 - X_1.$$

- ▶ Then  $T_n$ ,  $\bar{X}$  and  $D$  are **also normal random variables**.



# Normal Distribution: Sums and Averages

- ▶ Let  $X_1, \dots, X_n$  be independent and *normally distributed*. Let

$$T_n = X_1 + \dots + X_n, \quad \bar{X} = \frac{X_1 + \dots + X_n}{n}, \quad D = X_2 - X_1.$$

- ▶ Then  $T_n$ ,  $\bar{X}$  and  $D$  are **also normal random variables**.
- ▶ Let  $X_1, \dots, X_n \stackrel{i.i.d}{\sim} N(\mu, \sigma^2)$ . Then:

$$T_n \sim N(n\mu, n\sigma^2)$$

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

$$D \sim N(0, 2\sigma^2)$$

## Example

- ▶ Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.
- a) Find  $P(156 < X < 164)$ .

## Example

- ▶ Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.

a) Find  $P(156 < X < 164)$ .

$$\begin{aligned}P(156 < X < 164) &= P\left(\frac{156 - 160}{8} < \frac{X - 160}{8} < \frac{164 - 160}{8}\right) \\&= P(-0.5 < Z < 0.5) \\&= 0.3830\end{aligned}$$

## Example

- Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.

- a) Find  $P(156 < X < 164)$ .

$$\begin{aligned}P(156 < X < 164) &= P\left(\frac{156 - 160}{8} < \frac{X - 160}{8} < \frac{164 - 160}{8}\right) \\&= P(-0.5 < Z < 0.5) \\&= 0.3830\end{aligned}$$

- b) Find the probability that the average weight of 16 men chosen at random is between 156 and 164 pounds.

## Example

- Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.

- a) Find  $P(156 < X < 164)$ .

$$\begin{aligned}P(156 < X < 164) &= P\left(\frac{156 - 160}{8} < \frac{X - 160}{8} < \frac{164 - 160}{8}\right) \\&= P(-0.5 < Z < 0.5) \\&= 0.3830\end{aligned}$$

- b) Find the probability that the average weight of 16 men chosen at random is between 156 and 164 pounds.

$$\begin{aligned}\bar{X} &\sim N(160, 64/16 = 4) \\P(156 < \bar{X} < 164) &= P(-2 < Z < 2) \\&\approx 0.95\end{aligned}$$

## Example

- ▶ Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.
- c) Calculate the numbers  $A$  and  $B$  such that  $P(A < X < B) \approx 0.95$ .

## Example

- Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.
- c) Calculate the numbers  $A$  and  $B$  such that  $P(A < X < B) \approx 0.95$ .

$$0.95 \approx P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = P(-2\sigma + \mu < X < 2\sigma + \mu)$$

$$A = -2(8) + 160 = 144, \quad B = 2(8) + 160 = 176$$

## Example

- ▶ Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.
- c) Calculate the numbers  $A$  and  $B$  such that  $P(A < X < B) \approx 0.95$ .

$$0.95 \approx P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = P(-2\sigma + \mu < X < 2\sigma + \mu)$$

$$A = -2(8) + 160 = 144, \quad B = 2(8) + 160 = 176$$

- d) Calculate the numbers  $C$  and  $D$  such that the average of 256 randomly chosen adults is between  $C$  and  $D$  with probability approximately 0.95.



## Example

- ▶ Suppose we know the weight  $X$  of an adult man chosen at random is normally distributed with mean 160 pounds and variance 64 pounds<sup>2</sup>.
- c) Calculate the numbers  $A$  and  $B$  such that  $P(A < X < B) \approx 0.95$ .

$$0.95 \approx P\left(-2 < \frac{X - \mu}{\sigma} < 2\right) = P(-2\sigma + \mu < X < 2\sigma + \mu)$$

$$A = -2(8) + 160 = 144, \quad B = 2(8) + 160 = 176$$

- d) Calculate the numbers  $C$  and  $D$  such that the average of 256 randomly chosen adults is between  $C$  and  $D$  with probability approximately 0.95.

$$\bar{X}_{256} \sim N(160, 64/256 = 1/4)$$

$$C = -2\sigma + \mu = -2(1/2) + 160 = 159$$

$$D = 2\sigma + \mu = 2(1/2) + 160 = 161$$

# Central Limit Theorem

The Central Limit Theorem:

- ▶ Suppose  $X_1, X_2, \dots, X_n$  are *iid* with mean  $\mu$  and variance  $\sigma^2$ .
- ▶ Then, for large  $n$

$$T_n \sim N(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

*no matter the distribution of the individual  $X_i$*

- ▶ Allows approximation of all distributions using the normal distribution if you know the mean and variance.

# Central Limit Theorem

The Central Limit Theorem:

- ▶ Suppose  $X_1, X_2, \dots, X_n$  are *iid* with mean  $\mu$  and variance  $\sigma^2$ .
- ▶ Then, for large  $n$

$$T_n \sim N(n\mu, n\sigma^2) \quad \text{and} \quad \bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

*no matter the distribution of the individual  $X_i$*

- ▶ Allows approximation of all distributions using the normal distribution if you know the mean and variance.

**Note:** if  $X_1, \dots, X_n$  are normally distributed, then this applies for *all*  $n$ , not just large  $n$ .

## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .

## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .
- ▶ For each  $X_i$ ,  $\text{Mean}(X_i) = \theta$  and  $\text{Var}(X_i) = \theta(1 - \theta)$ .

## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .
- ▶ For each  $X_i$ ,  $\text{Mean}(X_i) = \theta$  and  $\text{Var}(X_i) = \theta(1 - \theta)$ .
- ▶ The sum is:  $T_n = X_1 + X_2 + \dots + X_n$ .

## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .
- ▶ For each  $X_i$ ,  $\text{Mean}(X_i) = \theta$  and  $\text{Var}(X_i) = \theta(1 - \theta)$ .
- ▶ The sum is:  $T_n = X_1 + X_2 + \dots + X_n$ .
- ▶ The proportion is:

$$P = \frac{X_1 + \dots + X_n}{n}.$$

## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .
- ▶ For each  $X_i$ ,  $\text{Mean}(X_i) = \theta$  and  $\text{Var}(X_i) = \theta(1 - \theta)$ .
- ▶ The sum is:  $T_n = X_1 + X_2 + \dots + X_n$ .
- ▶ The proportion is:

$$P = \frac{X_1 + \dots + X_n}{n}.$$



## Central Limit Theorem: Example

- ▶ Let  $X_1, X_2, \dots, X_n \stackrel{iid}{\sim} \text{Binomial}(1, \theta)$ .
- ▶ For each  $X_i$ ,  $\text{Mean}(X_i) = \theta$  and  $\text{Var}(X_i) = \theta(1 - \theta)$ .
- ▶ The sum is:  $T_n = X_1 + X_2 + \dots + X_n$ .
- ▶ The proportion is:

$$P = \frac{X_1 + \dots + X_n}{n}.$$

- ▶ For large  $n$ ,

$$T_n \sim N(n\theta, n\theta[1 - \theta])$$

$$P \sim N\left(\theta, \frac{\theta(1 - \theta)}{n}\right)$$

## Central Limit Theorem: Problem

- ▶ Suppose you are rolling a fair die 1000 times. Calculate the numbers  $A$  and  $B$  such that the average of the 1000 rolls is between  $A$  and  $B$  with probability approximately 0.95. You may assume the mean of one roll is 3.5 and the variance is  $35/12$ .

## Central Limit Theorem: Problem

- Suppose you are rolling a fair die 1000 times. Calculate the numbers  $A$  and  $B$  such that the average of the 1000 rolls is between  $A$  and  $B$  with probability approximately 0.95. You may assume the mean of one roll is 3.5 and the variance is  $35/12$ .

$$\text{Mean}(X_i) = 3.5, \quad \text{Var}(X_i) = 35/12$$

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right) = N\left(3.5, \frac{35}{12000}\right) \quad \text{by CLT}$$

$$A = -2\sigma + \mu = -2\sqrt{35/12000} + 3.5 \approx 3.392$$

$$B = 2\sigma + \mu = 2\sqrt{35/12000} + 3.5 \approx 3.608$$

# Statistics

- ▶ We have finished the first half of the course on [probability](#). Now, we move on to [statistics](#).

# Statistics

- ▶ We have finished the first half of the course on **probability**. Now, we move on to **statistics**.
- ▶ **Statistics** is used to make inductive statements about some phenomenon (coin-flipping, dice rolling) **after** observing data.

# Statistics

- ▶ We have finished the first half of the course on **probability**. Now, we move on to **statistics**.
- ▶ **Statistics** is used to make inductive statements about some phenomenon (coin-flipping, dice rolling) **after** observing data.
- ▶ Three main activities of statistics:
  1. Estimating numerical values of a parameter or parameters.
  2. Assessing accuracy of these estimates.
  3. Testing hypotheses about the numerical values of parameters.

# Statistics

- ▶ We have finished the first half of the course on **probability**. Now, we move on to **statistics**.
- ▶ **Statistics** is used to make inductive statements about some phenomenon (coin-flipping, dice rolling) **after** observing data.
- ▶ Three main activities of statistics:
  1. Estimating numerical values of a parameter or parameters.
  2. Assessing accuracy of these estimates.
  3. Testing hypotheses about the numerical values of parameters.
- ▶ Example: Suppose flip a coin 1,000 times and observe 700 heads.
  1. How do I estimate the probability  $\theta$  of achieving a head?
  2. How accurate is my estimate of  $\theta$ ?
  3. Is this a fair coin ( $\theta = 0.5$ )?

## Estimation of a parameter: Binomial parameter $\theta$

- Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?



## Estimation of a parameter: Binomial parameter $\theta$

- ▶ Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?
- ▶ An intuitive estimator for  $\theta$  is  $p = x/n$ , the **observed** proportion of successes.

## Estimation of a parameter: Binomial parameter $\theta$

- ▶ Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?
- ▶ An intuitive estimator for  $\theta$  is  $p = x/n$ , the **observed** proportion of successes.
- ▶ Consider the random variable  $P$ , the proportion of successes **prior** to performing the experiment.

## Estimation of a parameter: Binomial parameter $\theta$

- ▶ Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?
- ▶ An intuitive estimator for  $\theta$  is  $p = x/n$ , the **observed** proportion of successes.
- ▶ Consider the random variable  $P$ , the proportion of successes **prior** to performing the experiment.
  - ▶  $\text{Mean}(P) = \theta$  so  $p$  is “shooting at the right target”.  $p$  is then referred to as an **unbiased** estimate of  $\theta$ .

## Estimation of a parameter: Binomial parameter $\theta$

- ▶ Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?
- ▶ An intuitive estimator for  $\theta$  is  $p = x/n$ , the **observed** proportion of successes.
- ▶ Consider the random variable  $P$ , the proportion of successes **prior** to performing the experiment.
  - ▶  $\text{Mean}(P) = \theta$  so  $p$  is “shooting at the right target”.  $p$  is then referred to as an **unbiased** estimate of  $\theta$ .
- ▶ Difference between estimate and estimator:
  - ▶ **Estimate:** A function of the observed data used to estimate a given parameter. Ex:  $p$ .
  - ▶ **Estimator:** The random variable whose realization is the estimate. Ex:  $P$ .

# Estimation of a parameter: Binomial parameter $\theta$

- ▶ Recall a binomial random variable  $X \sim \text{Bin}(n, \theta)$ . How do we estimate the probability of success  $\theta$ ?
- ▶ An intuitive estimator for  $\theta$  is  $p = x/n$ , the **observed** proportion of successes.
- ▶ Consider the random variable  $P$ , the proportion of successes **prior** to performing the experiment.
  - ▶  $\text{Mean}(P) = \theta$  so  $p$  is “shooting at the right target”.  $p$  is then referred to as an **unbiased** estimate of  $\theta$ .
- ▶ Difference between estimate and estimator:
  - ▶ **Estimate:** A function of the observed data used to estimate a given parameter. Ex:  $p$ .
  - ▶ **Estimator:** The random variable whose realization is the estimate. Ex:  $P$ .
- ▶ To investigate the precision of an estimate, we need to consider the random variable estimator.

## Precision of an estimate: Binomial parameter $\theta$

- Precision of  $p$  as an estimate of  $\theta$  depends on the variance of random variable  $P$ .

## Precision of an estimate: Binomial parameter $\theta$

- ▶ Precision of  $p$  as an estimate of  $\theta$  depends on the variance of random variable  $P$ .
- ▶ By the CLT, two standard deviation rule, and approximations (see pg. 40-41), we get the approximate 95% confidence interval for  $\theta$  as

$$p \pm 2\sqrt{p(1-p)/n}$$

## Precision of an estimate: Binomial parameter $\theta$

- Precision of  $p$  as an estimate of  $\theta$  depends on the variance of random variable  $P$ .
- By the CLT, two standard deviation rule, and approximations (see pg. 40-41), we get the approximate 95% confidence interval for  $\theta$  as

$$p \pm 2\sqrt{p(1-p)/n}$$

- We can further approximate the 95% confidence interval with  $p(1-p) \leq 1/4$  to get

$$p \pm \sqrt{1/n} \tag{66}$$



## Precision of an estimate: Binomial parameter $\theta$

- Precision of  $p$  as an estimate of  $\theta$  depends on the variance of random variable  $P$ .
- By the CLT, two standard deviation rule, and approximations (see pg. 40-41), we get the approximate **95% confidence interval** for  $\theta$  as

$$p \pm 2\sqrt{p(1-p)/n}$$

- We can further approximate the 95% confidence interval with  $p(1-p) \leq 1/4$  to get

$$p \pm \sqrt{1/n} \tag{66}$$

- Correspondingly, the 99% confidence interval is

$$p \pm 2.576\sqrt{p(1-p)/n} \approx p \pm 1.288\sqrt{1/n}$$

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

$$p = x/n = 388/531 = 0.7307$$

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

$$p = x/n = 388/531 = 0.7307$$

Q2: Calculate the 95% confidence interval for  $\theta$  using the approximate 95% interval formula (66):

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

$$p = x/n = 388/531 = 0.7307$$

Q2: Calculate the 95% confidence interval for  $\theta$  using the approximate 95% interval formula (66):

$$p \pm \sqrt{1/n} = 0.7307 \pm \sqrt{1/531} = 0.7307 \pm 0.0434$$

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

$$p = x/n = 388/531 = 0.7307$$

Q2: Calculate the 95% confidence interval for  $\theta$  using the approximate 95% interval formula (66):

$$p \pm \sqrt{1/n} = 0.7307 \pm \sqrt{1/531} = 0.7307 \pm 0.0434$$

Q3: What is the sample size if we want the width of the confidence interval to be 0.02?

## Example

In the 2017-2018 NBA season, LeBron James shot 531 free throws and made 388. We want to estimate the probability  $\theta$  that LeBron James makes a free throw.

Q1: What is the estimate for  $\theta$ ?

$$p = x/n = 388/531 = 0.7307$$

Q2: Calculate the 95% confidence interval for  $\theta$  using the approximate 95% interval formula (66):

$$p \pm \sqrt{1/n} = 0.7307 \pm \sqrt{1/531} = 0.7307 \pm 0.0434$$

Q3: What is the sample size if we want the width of the confidence interval to be 0.02?

$$\text{We want } \sqrt{1/n} = 0.01 = 0.02/2.$$

$$1/n = 0.01^2$$

$$n = 1/0.01^2 = 10000$$