

STAT 111

Recitation 1

Mo Huang

Email: mohuang@wharton.upenn.edu

Office Hours: Wednesdays 3:00 - 4:00 pm, JMHH F96

Slides: github.com/mohuangx/STAT111-Spring2019

September 7, 2018

Logistics

- ▶ Section 207: 1:00 pm - 1:50 pm
- ▶ Section 208: 2:00 pm - 2:50 pm

- ▶ Every Friday, I will:
 1. Collect the homework for that week;
 2. Give you your graded homework; and
 3. Give you the homework for the week after (also posted on Canvas).

- ▶ If you do not collect your homework, it will be available in the STAT 111 box in the Statistics Department (4th Floor, JMHH).

- ▶ If you know in advance that you will not be able to attend a recitation, give your homework to Professor Ewens on Thursdays or put it in my mailbox (at entrance to the Statistics Department)

- ▶ Questions about course materials should be asked during office hours or in an email to Professor Ewens.

Statistics and Probability


- ▶ **Probability theory** is *deductive* (or “top-down logic”). We start with a theory and then consider the implications resulting from that theory.
- ▶ **Statistics** is *inductive* (or “bottom-up logic”). We start with finite observations and then attempt to make objective statements about the world from a sample.
- ▶ Statistical methods use probability theory to draw conclusions from observed data.

Statistics and Probability: Example

- ▶ We currently cure an illness with some medicine (“current medicine”) which cures 80% of patients.
- ▶ A new medicine is proposed (“new medicine”) which cures 1,643 people out of 2,000.

Is the new medicine more effective?

Probability theory




Statistics


Probability: Axioms

- ▶ $0 \leq P(A) \leq 1$, where A is any event.
- ▶ $P(A^C) = 1 - P(A)$ where A^C is the complement of A .
- ▶ $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Intersection $P(A \cap B)$



Union $P(A \cup B)$



Probability: Independence

- ▶ A and B are independent if and only if $P(A \cap B) = P(A)P(B)$
- ▶ Conditional probability: $P(A|B) = \frac{P(A \cap B)}{P(B)}$

Suppose A and B are independent. Then

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

So we have two definitions of independence:

1. $P(A \cap B) = P(A)P(B)$
2. $P(A|B) = P(A)$

Note: This year, conditional probability is not on the syllabus but it's still an interesting and useful concept!

Probability: Questions

Q1: A smoke alarm consists of two parts, A and B . If there is smoke, part A will detect it with probability 0.96 and part B will detect it with probability 0.98. The probability that they both detect it is 0.95. What is the probability that smoke will not be detected?

A1: Let A be the event that part A detects smoke and let B be the event that part B detects smoke. What is the event that smoke is not detected?

$A \cup B = \text{Smoke is detected} \rightarrow [A \cup B]^C = \text{Smoke is not detected}$

$$\begin{aligned}P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\&= 0.96 + 0.98 - 0.95 \\&= 0.99\end{aligned}$$

$$\begin{aligned}P([A \cup B]^C) &= 1 - P(A \cup B) \\&= 0.01\end{aligned}$$

Probability: Questions

Q2: A six-sided die is biased such that an odd number is twice as likely to occur as an even number. That is, the probability of an odd number is $2/9$ and an even number is $1/9$. Let A be the event that an even number occurs. Let B be the event that a number greater than or equal to 4 occurs. Find $P(A)$, $P(B)$, $P(A \cup B)$ and $P(A \cap B)$.

A2:

$$\begin{aligned}P(A) &= P(2, 4 \text{ or } 6) \\&= 1/9 + 1/9 + 1/9 \\&= 3/9\end{aligned}$$

$$\begin{aligned}P(A \cap B) &= P(4 \text{ or } 6) \\&= 1/9 + 1/9 \\&= 2/9\end{aligned}$$

$$\begin{aligned}P(B) &= P(4, 5 \text{ or } 6) \\&= 1/9 + 2/9 + 1/9 \\&= 4/9\end{aligned}$$

$$\begin{aligned}P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\&= 3/9 + 4/9 - 2/9 \\&= 5/9\end{aligned}$$

Probability: Questions

A2: $P(A) = 3/9$, $P(B) = 4/9$, $P(A \cap B) = 2/9$, $P(A \cup B) = 5/9$

Q3: Following Q2, find $P(A|B)$.

A3:

$$\begin{aligned} P(A|B) &= \frac{P(A \cap B)}{P(B)} \\ &= \frac{2/9}{4/9} \\ &= 1/2 \end{aligned}$$

Q4: Are A and B independent?

A4: No, because $P(A|B) \neq P(A)$. (Recall $P(A) = 1/3$).

Probability: Questions

Q5: An unfair coin ($P(H) = 0.4$) was flipped twice. Given that on at least one flip a head occurred, what is the probability that a head occurred on both flips?

A5: We want to find $P(HH|\text{at least one } H)$.

$$\begin{aligned}P(\text{at least one } H) &= P(HH) + P(HT) + P(TH) \\&= (0.4)^2 + (0.4)(0.6) + (0.6)(0.4) \\&= 0.64.\end{aligned}$$

$$\begin{aligned}P(HH|\text{at least one } H) &= \frac{P(HH \cap \text{at least one } H)}{P(\text{at least one } H)} \\&= \frac{P(HH)}{P(\text{at least one } H)} \\&= \frac{(0.4)^2}{0.64} \\&= 0.25\end{aligned}$$