# Heart stroke-volume variability in a murine model for heart failure with reduced ejection fraction

Gemma Fernández-Mendoza\*

Departamento de Física, Escuela Superior de Física y Matemáticas, Instituto Politécnico Nacional, 07738 Ciudad de México, México

Moisés Santillán<sup>†</sup>

Centro de Investigación y de Estudios Avanzados, Unidad Monterrey, 66628 Apodaca NL, México

(Dated: January 9, 2023)

## Abstract

Write an abstract

#### I. INTRODUCTION

#### II. MATERIALS AND METHODS

#### III. RESULTS

### A. Experimental Results

#### B. Variability analysis

Heart rate variability has been measured using a variety of techniques. The majority of them are based on the notion of signal stationarity. However, the heart rate's inherently non-stationary nature—which undergoes continuous physiological change to adapt to outside stimuli—presents a significant challenge that could lead to inaccurate results [1]. Although a number of signal preprocessing methodologies have been suggested to address these problems, nonlinear analysis-based strategies are frequently used and seem to produce reliable result [1–7]. One of them that is used in different scientific domains is the Poincairé plot [6, 8], which has numerous applications in different scientific domains [9]. In a Poincairé plot, all values of a time series are plotted against previous values, leading to an ellipsoidal point cloud. Moreover, this diagram may be analyzed quantitatively by the standard deviations of the point projections along the lines y = x (SD2) and y = -x (SD1). The transverse axis of the ellipse (SD1) is a measure of the short-term changes in the the time series, while the longitudinal axis (SD2) reflects long-term changes. In the particular case of heart-beat duration time-series, SD1 is considered as an indicator of the sympathetic activity, whereas SD2 is considered as an inverse indicator of the sympathetic activity [10].

<sup>\*</sup> gemmag.fernandez@gmail.com

<sup>†</sup> http://moises-santillan.github.io; msantillan@cinvestav.mx




FIG. 1. Schematic representation of the model

#### C. Mathematical Model

### IV. CONCLUDING REMARKS

## ACKNOWLEDGMENTS

We wish to acknowledge the support of the author community in using REVT<sub>E</sub>X, offering suggestions and encouragement, testing new versions, . . . .

- [1] N. MARWAN, M. CARMENROMANO, M. THIEL, and J. KURTHS, Recurrence plots for the analysis of complex systems, Physics Reports 438, 237 (2007).
- [2] N. Marwan, N. Wessel, U. Meyerfeldt, A. Schirdewan, and J. Kurths, Recurrence-plot-based measures of complexity and their application to heart-rate-variability data, Physical Review E 66, 10.1103/physreve.66.026702 (2002).
- [3] A. E. Aubert, B. Seps, and F. Beckers, Heart rate variability in athletes, Sports Medicine 33, 889 (2003).
- [4] A. Giuliani, G. Piccirillo, V. Marigliano, and A. Colosimo, A nonlinear explanation of aginginduced changes in heartbeat dynamics, American Journal of Physiology-Heart and Circulatory Physiology 275, H1455 (1998).

- [5] U. R. Acharya, K. P. Joseph, N. Kannathal, C. M. Lim, and J. S. Suri, Heart rate variability: a review, Medical & Biological Engineering & Computing 44, 1031 (2006).
- [6] C. L. Webber and J. P. Zbilut, Dynamical assessment of physiological systems and states using recurrence plot strategies, Journal of Applied Physiology **76**, 965 (1994).
- [7] T. Henriques, M. Ribeiro, A. Teixeira, L. Castro, L. Antunes, and C. Costa-Santos, Nonlinear methods most applied to heart-rate time series: A review, Entropy 22, 309 (2020).
- [8] R. A. Hoshi, C. M. Pastre, L. C. M. Vanderlei, and M. F. Godoy, Assessment of heart rate complexity recovery from maximal exercise using recurrence quantification analysis, in *Springer Proceedings in Physics* (Springer International Publishing, 2016) pp. 157–168.
- [9] A. Voss, S. Schulz, R. Schroeder, M. Baumert, and P. Caminal, Methods derived from non-linear dynamics for analysing heart rate variability, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences 367, 277 (2008).
- [10] G. Zimatore, M. C. Gallotta, M. Campanella, P. H. Skarzynski, G. Maulucci, C. Serantoni, M. D. Spirito, D. Curzi, L. Guidetti, C. Baldari, and S. Hatzopoulos, Detecting metabolic thresholds from nonlinear analysis of heart rate time series: A review, International Journal of Environmental Research and Public Health 19, 12719 (2022).