

Dietary quality in adolescents as a risk factor for painful temporomandibular disorders and headache in young adult members of the Danish National Birth Cohort



Mojdeh Mansoori¹, Cristina Exposto¹, Bodil Hammer Bech², Luda Diatchenko³, Lene Baad-Hansen¹

1. Department of Dentistry and Oral Health, Section for Orofacial Pain and Jaw Function, Aarhus University, Denmark
2. Department of Public Health - Epidemiology, Aarhus University, Denmark
3. Departments of Anesthesia & Faculty of Dentistry, McGill University, and Genome Quebec Innovation Centre, Montreal, Quebec, Canada

Background & Hypothesis:

- Painful temporomandibular disorder (TMD-P) - The most common non-odontogenic-related chronic orofacial pain
- Most well-established risk factors: Gender, age, psychosocial factors, and pain sensitivity phenotype
- Although there is some evidence that the presence of TMD-P affects dietary choice and nutritional intake, the way that nutritional factors may influence the development of TMD-P has not been explored.
- One proposed reason of this research: Proinflammatory markers increased in chronic pain and obesity situations.
- Identifying risk factors for TMD-P and headache could eventually be a target for either prevention or early intervention.

Study design:



Data analyses:

- Regression modelling adjusted for known confounders
- Analysis is ongoing and results therefore “pending”

Keywords:

- Painful Temporomandibular Disorders, Healthy Eating Index, Headache