

# Complex Exponentials

R. Mok BSc(AdvMath), MTeach, GradCertDS

AI NSW Mathematics Heads of Department Conference, October 2022

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula
- 4 What is an exponential?
- 5 Bibliography

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula
- 4 What is an exponential?
- 5 Bibliography

# Opening Question

Here's a few questions to warm you up for the session to see if you're awake!

- 1 Solve for  $x \in \mathbb{N} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 2 Solve for  $x \in \mathbb{R} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 3 Solve for  $x \in \mathbb{C} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

# Opening Question

Here's a few questions to warm you up for the session to see if you're awake!

- 1 Solve for  $x \in \mathbb{N} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 2 Solve for  $x \in \mathbb{R} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 3 Solve for  $x \in \mathbb{C} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Answers:

- 1 A, B and D
- 2 C
- 3 B and C

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula
- 4 What is an exponential?
- 5 Bibliography

# De Moivre's Theorem

## De Moivre's Theorem (DMT)[1]

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

# De Moivre's Theorem

## De Moivre's Theorem (DMT)[1]

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

## Pythagoras' Theorem

$$c^2 = a^2 + b^2$$



# De Moivre's Theorem

## De Moivre's Theorem (DMT)[1]

For all integers  $n$

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

## Pythagoras' Theorem

A triangle with side lengths  $a, b, c$  is right angled if and only if

$$c^2 = a^2 + b^2$$

# De Moivre's Theorem

## De Moivre's Theorem (DMT)[1]

For all **integers**  $n$

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

This is to ensure the exponential of a complex number yields one value for all arguments of that complex number.

## Pythagoras' Theorem

A triangle with side lengths  $a, b, c$  is right angled if and only if

$$c^2 = a^2 + b^2$$

# Exploring De Moivre's Theorem with DESMOS

<https://www.desmos.com/calculator/igqisujucx>

# Extending De Moivre's Theorem to Rational Exponents

If we want to extend DMT to rational exponents, this happens:

## Example

$$\begin{aligned}1 &= (\cos 0 + i \sin 0) \text{ or } (\cos 2\pi + i \sin 2\pi) \\1^{\frac{1}{2}} &= (\cos 0 + i \sin 0)^{\frac{1}{2}} \text{ or } (\cos 2\pi + i \sin 2\pi)^{\frac{1}{2}} \\&= \cos 0 + i \sin 0 \text{ or } \cos \pi + i \sin \pi \\&= 1 \text{ or } -1\end{aligned}$$

# Extending De Moivre's Theorem to Rational Exponents

If we want to extend DMT to rational exponents, this happens:

## Example

$$\begin{aligned}1 &= (\cos 0 + i \sin 0) \text{ or } (\cos 2\pi + i \sin 2\pi) \\1^{\frac{1}{2}} &= (\cos 0 + i \sin 0)^{\frac{1}{2}} \text{ or } (\cos 2\pi + i \sin 2\pi)^{\frac{1}{2}} \\&= \cos 0 + i \sin 0 \text{ or } \cos \pi + i \sin \pi \\&= 1 \text{ or } -1\end{aligned}$$

Exponentials of complex numbers are **multi-valued** functions<sup>1</sup>.

---

<sup>1</sup>HSC uses the terminology 'one-to-many' relation.

# Roots of Unity

This process is essentially how Roots of Unity are found:

## Example

$$z^n = 1$$

$$= (\cos(2k\pi) + i \sin(2k\pi)) \text{ for } k \in \mathbb{Z}$$

$$z = \cos\left(\frac{2k\pi}{n}\right) + i \sin\left(\frac{2k\pi}{n}\right) \text{ by the extended DMT}$$

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$



# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$

$\sqrt{\cdot} : \mathbb{C} \rightarrow \mathbb{C}$  is single-valued and the input is restricted to the principal argument. However, the exponential is multi-valued.

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$

$\sqrt{\cdot} : \mathbb{C} \rightarrow \mathbb{C}$  is single-valued and the input is restricted to the principal argument. However, the exponential is multi-valued.

What even is an exponential then? More on that later...

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .
- Irrational exponents do not divide the revolutions nicely and hence have infinite values.

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .
- Irrational exponents do not divide the revolutions nicely and hence have infinite values.




Figure:  $1^{\frac{1}{5}}$  vs. irrational exponents such as  $1^{\pi}$ ,  $(-1)^{\sqrt{2}}$ , etc

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.
- Extending DMT to irrational exponents yields infinite values.



# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.
- Extending DMT to irrational exponents yields infinite values.
- What's next? Complex exponents.

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula**
- 4 What is an exponential?
- 5 Bibliography

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula [1] which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula [1] which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

This opens up new avenues of discussion that were previously not possible in the old syllabus...

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula [1] which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

This opens up new avenues of discussion that were previously not possible in the old syllabus...

... such as:

## De Moivre's Theorem Restated

$$(e^{ix})^n = e^{inx}$$

with the same discussions about  $n$  as in the previous section.

# Complex Logarithms

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

# Complex Logarithms

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

Does such a function  $\log : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  that is the inverse of complex exponentiation exist? Yes it does - proof beyond the time we have now.

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

Does such a function  $\log : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  that is the inverse of complex exponentiation exist? Yes it does - proof beyond the time we have now.

## Complex Logarithm [2, p. 23]

If  $z$  is a complex number  $\neq 0$ , then there exist complex numbers  $\omega$  such that  $e^{\omega} = z$  where  $\omega$  is in the form  $\log_e |z| + i \operatorname{Arg} z + 2n\pi i$  where  $n \in \mathbb{Z}$ .



# Complex Exponential Equations

## Example

Solve for  $z \in \mathbb{C} : e^z = -1$ .

# Complex Exponential Equations

## Example

Solve for  $z \in \mathbb{C} : e^z = -1$ .

$$\begin{aligned} z &= \log(-1) \\ &= \log_e |-1| + i \arg(-1) \\ &= 0 + i \arg(-1) \\ &= i(2k + 1)\pi \text{ for } k \in \mathbb{Z} \end{aligned}$$

# Complex Exponential Equations

## Example

Solve for  $z \in \mathbb{C} : e^z = -1$ .

$$\begin{aligned} z &= \log(-1) \\ &= \log_e |-1| + i \arg(-1) \\ &= 0 + i \arg(-1) \\ &= i(2k + 1)\pi \text{ for } k \in \mathbb{Z} \end{aligned}$$

If we test any of these solutions such as  $z_1 = i3\pi$ :

$$e^{z_1} = e^{i3\pi} = \cos(3\pi) + i \sin(3\pi) = -1$$

Hence,  $i3\pi$  is indeed a solution to  $e^z = -1$ .

# Complex Trigonometric Equations

It can be shown using Euler's Formula that:

## Trigonometric Functions

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}$$

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i}$$

# Complex Trigonometric Equations

## Example

Solve for  $z \in \mathbb{C} : \sin z = 2$

# Complex Trigonometric Equations

## Example

Solve for  $z \in \mathbb{C}$  :  $\sin z = 2$

$$\frac{e^{iz} + e^{-iz}}{2} = 2$$

$$e^{iz} + e^{-iz} = 4$$

$$e^{2iz} - 4e^{iz} + 1 = 0$$

$$e^{iz} = \frac{4 \pm \sqrt{4^2 - 4(1)(1)}}{2} = 2 \pm \sqrt{3}$$

$$iz = \log_e(2 \pm \sqrt{3}) + i2k\pi \text{ for } k \in \mathbb{Z}$$

$$z = \frac{1}{i} \log_e(2 \pm \sqrt{3}) + 2k\pi$$

## Example

Evaluate  $i^i$ .

## Example

Evaluate  $i^i$ .

$$\begin{aligned} i^i &= e^{\log i^i} \\ &= e^{i \log i} \\ &= e^{i(\log_e |i| + i \arg i)} \\ &= e^{i^2 \arg i} \\ &= e^{-1(\frac{\pi}{2} + 2k\pi)} \text{ for } k \in \mathbb{Z} \\ &= \dots, 111.3177784899\dots, 0.2078795764\dots, 0.0003882032039\dots, \dots \end{aligned}$$



# More examples

## Example

- Evaluate  $1^i$ .
- Evaluate  $(-1)^{\sqrt{2}}$ .

## Example

- Evaluate  $1^i$ .
- Evaluate  $(-1)^{\sqrt{2}}$ .

Answers:

$$1^i = e^{-2k\pi} \text{ for } k \in \mathbb{Z}$$

$$(-1)^{\sqrt{2}} = e^{\sqrt{2}i\pi(2k+1)} \text{ for } k \in \mathbb{Z}$$

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula
- 4 What is an exponential?
- 5 Bibliography

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}?$

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}?$

- Stage 5: Extend the concept of exponentials continuously with index laws that fixes this problem.

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}$ ?

- Stage 5: Extend the concept of exponentials continuously with index laws that fixes this problem.
- Stage 6: The number  $e$  is introduced in the topic of calculus. Jacob Bernoulli derived this value in 1683 when considering the compound interest problem of compounding a principal amount of \$1 at an ever increasing compounding period over a year. He arrived at the discovery:

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

# Exponential Function

e

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

if and only if

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}$$

The proof uses concepts in Binomial Theorem and Analysis [3, pp. 64-65]. It can also be shown (using similar methods as in the proof) that:

Exponential Function [3, p. 174]

$$\exp(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!} = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n$$



## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive:

<https://www.desmos.com/calculator/deicpgcinp>

Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{n})^n$  for  $n = 1, 2, 3$

## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive:

<https://www.desmos.com/calculator/deicpgcinp>

Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{10})^n$  for  $n = 1, 2, \dots, 10$

## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive:

<https://www.desmos.com/calculator/deicpgcinp>

Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{1000})^n$  for  $n = 1, 2, \dots, 1000$

# What is an exponential?

What does  $a^b$  really mean?

# What is an exponential?

## What does $a^b$ really mean?

$a^b$  is equivalent to  $\exp(b \log a)$  [4, p. 46].

Ahlfors [4] adopts the convention that if  $a$  is restricted to positive numbers,  $\log a$  shall be real (unless the contrary is stated) and  $a^b$  has a single value. Otherwise,  $\log a$  is the complex logarithm and  $a^b$  has in general infinitely many values.

# What is an exponential?

## What does $a^b$ really mean?

$a^b$  is equivalent to  $\exp(b \log a)$  [4, p. 46].

Ahlfors [4] adopts the convention that if  $a$  is restricted to positive numbers,  $\log a$  shall be real (unless the contrary is stated) and  $a^b$  has a single value. Otherwise,  $\log a$  is the complex logarithm and  $a^b$  has in general infinitely many values.

We already need, and use, this definition in deriving Calculus results such as:

$$\begin{aligned}y &= a^x \\&= e^{\log_e(a^x)} \\&= e^{x \log_e(a)} \\ \frac{dy}{dx} &= \log_e(a) e^{x \log_e(a)} = \log_e(a) a^x\end{aligned}$$

# Table of Contents

- 1 Opening Question
- 2 Revisiting De Moivre's Theorem
- 3 Euler's Formula
- 4 What is an exponential?
- 5 Bibliography**

# Bibliography

- [1] NSW Education Standards Authority, *Mathematics extension 2 stage 6 syllabus*, 2017.
- [2] T. M. Apostol, *Mathematical Analysis*, 2nd ed. Addison-Wesley, 1974, p. 23, ISBN: 0-201-00288-4.
- [3] W. Rudin, *Principles of Mathematical Analysis*, 3rd ed. McGraw-Hill, Inc., 1976, pp. 64–65, 178, ISBN: 0-07-054235-X.
- [4] L. V. Ahlfors, *Complex Analysis*, 3rd ed. McGraw-Hill, Inc., 1979, p. 46, ISBN: 0-07-000657-1.

Source files and PDF for this presentation can be found on:

[https://github.com/moksifu/complex\\_exponentials](https://github.com/moksifu/complex_exponentials) under the MIT Licence.