

# Complex Exponentials

A can of worms we shall open!

R. Mok BSc(AdvMath), MTeach, GradCertDS

AISSNSW Mathematics Heads of Department Conference, August 2021

# Table of Contents

# Table of Contents

# Opening Question

Here's a few questions to warm you up for the session to see if you're awake!

- 1 Solve for  $x \in \mathbb{N} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 2 Solve for  $x \in \mathbb{R} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 3 Solve for  $x \in \mathbb{C} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

# Opening Question

Here's a few questions to warm you up for the session to see if you're awake!

- 1 Solve for  $x \in \mathbb{N} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 2 Solve for  $x \in \mathbb{R} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$
- 3 Solve for  $x \in \mathbb{C} : x^2 = 4$   
A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Answers:

- 1 A, B and D
- 2 C
- 3 B and C

# Table of Contents

# De Moivre's Theorem

## De Moivre's Theorem (DMT) (syllabus)

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

# De Moivre's Theorem

## De Moivre's Theorem (DMT) (syllabus)

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

## Pythagoras' Theorem

$$c^2 = a^2 + b^2$$



# De Moivre's Theorem

## De Moivre's Theorem (DMT) (syllabus)

For all integers  $n$

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

## Pythagoras' Theorem

A triangle with side lengths  $a, b, c$  is right angled if and only if

$$c^2 = a^2 + b^2$$

DESMOS Interactive: <https://www.desmos.com/calculator/szywgvtxm8>

# De Moivre's Theorem

## De Moivre's Theorem (DMT) (syllabus)

For all integers  $n$

$$[r(\cos \theta + i \sin \theta)]^n = r^n(\cos(n\theta) + i \sin(n\theta))$$

This is to ensure the exponential of a complex number yields one value for all arguments of that complex number.

## Pythagoras' Theorem

A triangle with side lengths  $a, b, c$  is right angled if and only if

$$c^2 = a^2 + b^2$$

DESMOS Interactive: <https://www.desmos.com/calculator/szywgvtxm8>

# Extending De Moivre's Theorem to Rational Exponents

If we want to extend DMT to rational exponents, this happens:

## Example

$$\begin{aligned}1 &= (\cos 0 + i \sin 0) \text{ or } (\cos 2\pi + i \sin 2\pi) \\1^{\frac{1}{2}} &= (\cos 0 + i \sin 0)^{\frac{1}{2}} \text{ or } (\cos 2\pi + i \sin 2\pi)^{\frac{1}{2}} \\&= \cos 0 + i \sin 0 \text{ or } \cos \pi + i \sin \pi \\&= 1 \text{ or } -1\end{aligned}$$

# Extending De Moivre's Theorem to Rational Exponents

If we want to extend DMT to rational exponents, this happens:

## Example

$$\begin{aligned}1 &= (\cos 0 + i \sin 0) \text{ or } (\cos 2\pi + i \sin 2\pi) \\1^{\frac{1}{2}} &= (\cos 0 + i \sin 0)^{\frac{1}{2}} \text{ or } (\cos 2\pi + i \sin 2\pi)^{\frac{1}{2}} \\&= \cos 0 + i \sin 0 \text{ or } \cos \pi + i \sin \pi \\&= 1 \text{ or } -1\end{aligned}$$

Exponentials of complex numbers are **multi-valued** functions<sup>1</sup>.

---

<sup>1</sup>HSC uses the terminology 'one-to-many' relation.

This process is essentially how Roots of Unity are found:

## Example

$$z^n = 1$$

$$= (\cos(2k\pi) + i \sin(2k\pi)) \text{ for } k \in \mathbb{Z}$$

$$z = \cos\left(\frac{2k\pi}{n}\right) + i \sin\left(\frac{2k\pi}{n}\right) \text{ by the extended DMT}$$

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

- A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$

# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$     B)  $x = 4^{\frac{1}{2}}$     C)  $x = \pm 2$     D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$

$\sqrt{\cdot} : \mathbb{C} \rightarrow \mathbb{C}$  is single-valued and is restricted to the principal argument.  
However, the exponential is multi-valued.



# Back to the opening question

## Opening Question

Solve for  $x \in \mathbb{C} : x^2 = 4$

A)  $x = 2$    B)  $x = 4^{\frac{1}{2}}$    C)  $x = \pm 2$    D)  $x = \sqrt{4}$

Note:  $\sqrt{4} = 2$ ,  $4^{\frac{1}{2}} = \pm 2$

$\sqrt{\cdot} : \mathbb{C} \rightarrow \mathbb{C}$  is single-valued and is restricted to the principal argument.  
However, the exponential is multi-valued.

What even is an exponential then? More on that later...

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .
- Irrational exponents do not divide the revolutions nicely and hence have infinite values.

# Complex Exponentials with Irrationals

- Rational exponents  $m/n$  divide the revolutions of arguments nicely such that their results repeat themselves with periodicity (at most)  $n$ .
- Irrational exponents do not divide the revolutions nicely and hence have infinite values.




Figure:  $1^{\frac{1}{5}}$  vs. irrational exponents such as  $1^{\pi}$ ,  $(-1)^{\sqrt{2}}$ , etc

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.
- Extending DMT to irrational exponents yields infinite values.

# Quick Recap

- De Moivre's Theorem (DMT) for integer exponents yields single values.
- Extending DMT to rational exponents  $m/n$  yields (at most)  $n$  values.
- Extending DMT to irrational exponents yields infinite values.
- What's next? Complex exponents.



# Table of Contents

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula (**syllabus**) which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula (**syllabus**) which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

This opens up new avenues of discussion that were previously not possible in the old syllabus...

# Euler's Formula

HSC Dotpoint MEX-N1.3 introduces Euler's Formula (**syllabus**) which was not in the old syllabus:

## Euler's Formula

$$e^{ix} = \cos x + i \sin x \text{ for real } x$$

This opens up new avenues of discussion that were previously not possible in the old syllabus...

... such as:

## De Moivre's Theorem Restated

$$(e^{ix})^n = e^{inx}$$

with the same discussions about  $n$  as in the previous section.

# Complex Logarithms

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

Does such a function  $\log : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  that is the inverse of complex exponentiation exist? Yes it does - proof beyond the time we have now.

Euler's Formula opens up discussion about:

$$z = |z|e^{i \arg z}$$
$$\log z = \log(|z|e^{i \arg z})$$

Does such a function  $\log : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$  that is the inverse of complex exponentiation exist? Yes it does - proof beyond the time we have now.

(Principal) Complex Logarithm (**daners**)

$$\text{Log} z = \log_e |z| + i \text{Arg} z$$

## Example

Solve for  $z \in \mathbb{C} : e^z = 1$ .



# Complex Exponential Equations

## Example

Solve for  $z \in \mathbb{C} : e^z = 1$ .

$$\begin{aligned} z &= \log(1) \\ &= \log_e |1| + i \arg(1) \\ &= 0 + i \arg(1) \\ &= i2k\pi \text{ for } k \in \mathbb{Z} \end{aligned}$$

# Complex Exponential Equations

## Example

Solve for  $z \in \mathbb{C} : e^z = 1$ .

$$\begin{aligned} z &= \log(1) \\ &= \log_e |1| + i \arg(1) \\ &= 0 + i \arg(1) \\ &= i2k\pi \text{ for } k \in \mathbb{Z} \end{aligned}$$

If we test any of these solutions such as  $z_1 = i2\pi$ :

$$e^{z_1} = e^{i2\pi} = 1$$

Hence,  $i2\pi$  is indeed a solution to  $e^z = 1$ .

# Complex Trigonometric Equations

It can be shown using Euler's Formula that:

## Trigonometric Functions

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}$$

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i}$$

# Complex Trigonometric Equations

## Example

Solve for  $z \in \mathbb{C} : \sin z = 2$

# Complex Trigonometric Equations

## Example

Solve for  $z \in \mathbb{C}$  :  $\sin z = 2$

$$\frac{e^{iz} + e^{-iz}}{2} = 2$$

$$e^{iz} + e^{-iz} = 4$$

$$e^{2iz} - 4e^{iz} + 1 = 0$$

$$e^{iz} = \frac{4 \pm \sqrt{4^2 - 4(1)(1)}}{2} = 2 \pm \sqrt{3}$$

$$iz = \log_e(2 \pm \sqrt{3}) + i2k\pi \text{ for } k \in \mathbb{Z}$$

$$z = \frac{1}{i} \log_e(2 \pm \sqrt{3}) + 2k\pi$$

## Example

Evaluate  $i^i$ .

## Example

Evaluate  $i^i$ .

$$\begin{aligned} i^i &= e^{\log i^i} \\ &= e^{i \log i} \\ &= e^{i(\log_e |i| + i \arg i)} \\ &= e^{i^2 \arg i} \\ &= e^{-1(\frac{\pi}{2} + 2k\pi)} \text{ for } k \in \mathbb{Z} \\ &= \dots, 111.3177784899\dots, 0.2078795764\dots, 0.0003882032039\dots, \dots \end{aligned}$$

# More examples

## Example

- Evaluate  $1^i$ .
- Evaluate  $(-1)^{\sqrt{2}}$ .



## Example

- Evaluate  $1^i$ .
- Evaluate  $(-1)^{\sqrt{2}}$ .

Answers:

$$1^i = e^{-2k\pi} \text{ for } k \in \mathbb{Z}$$

$$(-1)^{\sqrt{2}} = e^{\sqrt{2}i\pi(2k+1)} \text{ for } k \in \mathbb{Z}$$

# Table of Contents

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}?$

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}?$

- Stage 5: Extend the concept of exponentials with index laws that fixes this problem.

# A progressive relearning at each stage

- Stage 4: A power is how many times that base multiplies itself.

$$2^3 = 2 \times 2 \times 2 = 8$$

Problem: How can  $2^{\frac{-1}{2}}$  mean  $\underbrace{2 \times \dots \times 2}_{\frac{-1}{2} \text{ times}}$ ?

- Stage 5: Extend the concept of exponentials with index laws that fixes this problem.
- Stage 6: The number  $e$  is introduced in the topic of calculus. Jacob Bernoulli derived this value in 1683 when considering the compound interest problem of compounding a principal amount of \$1 at an ever increasing compounding period over a year. He arrived at the discovery:

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

# Exponential Function

e

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

if and only if

$$e = \sum_{k=0}^{\infty} \frac{1}{k!}$$

The proof uses concepts in Binomial Theorem and Analysis (**rudin**). It can also be shown (using similar methods as in the proof) that:

Exponential Function (**rudin**)

$$\exp(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!} = \lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n$$

## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive: <https://www.desmos.com/calculator/p4rzq24eck>  
Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{3})^n$  for  $n = 1, 2, 3$



## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive: <https://www.desmos.com/calculator/p4rzq24eck>  
Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{n})^n$  for  $n = 1, 2, \dots, 10$

## Plots of increasing $n$ for $(1 + \frac{x}{n})^n$

DESMOS Interactive: <https://www.desmos.com/calculator/p4rzq24eck>

Plots of  $(1 + \frac{x}{n})^n$  for  $x = 1$  or  $x = i\pi$  and differing values of  $n$  are shown below:




Figure:  $(1 + \frac{x}{1000})^n$  for  $n = 1, 2, \dots, 1000$

# Table of Contents

# Bibliography