

# Adapting the MRIQC structural workflow for estimation of FLAIR scan quality



We have developed a modified version of the MRIQC workflow for FLAIR scans and validated it across multiple sites and head-motion degraded scans

Molly Ireland<sup>1</sup>, Heath Pardoe<sup>1</sup>, Perchyonok Yuliya<sup>2</sup>, Ricky Lu<sup>2</sup>, Greg Fitt<sup>2</sup>, David N Vaughan<sup>1,2</sup>, Chris Tailby<sup>1,3</sup>, David F Abbott<sup>1</sup> and Graeme D Jackson<sup>1,2</sup> for the Australian Epilepsy Project Investigators

1. Florey Institute of Neuroscience and Mental Health, Melbourne, Australia  
2. Department of Neurology, Austin Health, Melbourne, Australia  
3. Department of Neuropsychology, Austin Health, Melbourne, Australia

**Background**

MRIQC is a widely used tool to generate **automated assessments of MRI scan quality**; however its structural workflow does not include FLAIR scans

FLAIR-specific challenges:


- Low signal-to-noise (SNR) due to CSF suppression
- White-matter (WM) hyperintensities

This work **adapted the MRIQC tool** for assessment of 3D isotropic **FLAIR** scans


**Methods**

Iteratively modified the MRIQC workflow to process FLAIR scans from the Australia Epilepsy Project (AEP) and OpenNeuro ds004332 dataset.

| Dataset  | N   | Age            | Multi-site         |
|----------|-----|----------------|--------------------|
| AEP      | 272 | 37y (18-67y)   | Yes (235/25/7/3/2) |
| ds004332 | 29  | 23.5y (19-36y) | No                 |




**Figure 1:** Adapted MRIQC structural workflow for FLAIR scans. Modifications to the workflow are embedded within preexisting processing modules. Changes: ▲ = increase, ▼ = decrease, + = addition.



**Results**

Three changes significantly improved overall preprocessing for FLAIR:

1. Spatial transforms using FLAIR templates in MNI space
2. Registration with ANTs B-spline symmetric normalization (SyN) algorithm showed significant improvements for motion-affected FLAIR scans relative to default SyN
3. Tissue segmentation with a nonparametric likelihood model



**Figure 2:** Comparisons of overall scan quality ratings against selected IQMs (snr\_total, cnr and fber) for T1w and FLAIR scans.

- Many more IQMs correlated with visual ratings for FLAIR (N = 44) compared to T1 (N = 18)
- FLAIR and T1w IQMs were consistent with each other. Same direction for 13/15 IQMs with statistically significant associations for both modalities
- Mean adjusted R<sup>2</sup> higher for FLAIR (0.2) than T1w (0.04)

To evaluate our FLAIR pipeline, the relationship between expert neuroradiological ratings and output image quality metrics (IQMs) was assessed with linear regression.

Finally, we compared these relationships with the same analyses applied to output from T1w scans from the unmodified MRIQC pipeline.

**Conclusions**

- Modified MRIQC workflow to estimate image quality metrics for 3D isotropic FLAIR
- Validated relationship between FLAIR IQMs from multisite, motion-affected and motion-free scans against neuroradiologist visual image quality ratings
- Predominantly developed with standardised Australian Epilepsy Project data meaning more multisite data required to validate performance