

Paper Title*

1st Given Name Surname
 dept. name of organization (of Aff.)
 name of organization (of Aff.)
 City, Country
 email address

2nd Given Name Surname
 dept. name of organization (of Aff.)
 name of organization (of Aff.)
 City, Country
 email address

Abstract—
Index Terms—

So each IaaS provider try to solve following optimizaiton problem:

I. INTRODUCTION

II. SYSTEM MODEL

$$\varphi_s(R_s) = \log\left(\sum_{i=1}^{N_I} r_{i,s}\right) - \sum_{i=1}^{N_I} p_i r_{i,s} F_s \quad (1)$$

$$t_s = \frac{1}{\sum_{i=1}^{N_I} r_{i,s}} \sum_{i=1}^{N_I} r_{s,i} (t_{i,s}^{sensors} + t_{i,s} + t_{i,s}^{actuators}) \quad (2)$$

$$t_{i,s} = \frac{1}{u_{i,s} C_i - r_{i,s} F_s} \quad (3)$$

$$\max_{\Lambda_s} \varphi_s(R_s) \quad (4a)$$

subject to:

$$R_s^{min} \leq \sum_{k=1}^{N_I} r_{k,s} \quad (4b)$$

$$0 \leq r_{j,s}, \forall j \in \{1, \dots, N_I\} \quad (4c)$$

$$r_{j,s} R_s \leq 0.9 u_{j,s} C_j, \forall j \in \{1, \dots, N_I\} \quad (4d)$$

$$t_{j,s} + \frac{1}{u_{j,s} C_j - r_{j,s} F_s} \leq 0 \quad (4e)$$

$$\varphi_i(p_i, U_i, \Lambda_S^i) = \sum_{s=1}^{N_S} p_i \lambda_{i,s}^i R_s - P^{electrical}(P_i(\sum_{s=1}^{N_S} (u_{i,s}))) \quad (5)$$

$$P_i(u_i) = P_i^{idle} + (P_i^{max} - P_i^{idle}) u_i \quad (6)$$

$$\max_{p_i, U_i, \Lambda_S^i} \varphi_i(p_i, U_i, \Lambda_S^i) \quad (7a)$$

subject to:

$$0 \leq u_s^i, \forall s \in \{1, \dots, N_S\} \quad (7b)$$

$$\sum_{s=1}^{N_S} u_s^i \leq 1 \quad (7c)$$

$$P_{idle}^i + (P_{max}^i - P_{idle}^i) \sum_{s=1}^{N_S} u_s^i \leq \bar{P}^i \quad (7d)$$

$$\Lambda_s^i \in SOL(F_s), \forall s \quad (7e)$$

$$\pi(x^i, x^{-i}) - \pi(y^i, x^{-i}) = \varphi_i(x^i, x^{-i}) - \varphi_i(y^i, x^{-i}) \quad (8)$$

It's easy to show that any global minimum of function π is a nash equilibrium of corresponding game.

show or not??? For IaaS providers exact potential function can be writen as:

$$\pi(x^i, x^{-i}) = \sum_{i=1}^{N_I} \varphi_i(x^i, x^{-i}) \quad (9)$$

III. PROBLEM FORMULATION

$$\max_{p^i, U^i, \Lambda_S^i, \sigma_S^i, \gamma_S^i, \nu_S^i, \eta_S^i} \varphi(p_i, U^i, \Lambda_S^i) \quad (10a)$$

subject to:

$$\sum_{s=1}^{N_S} u_s^i \leq 1 \quad (10b)$$

$$0 \leq u_s^i, \forall s \in \{1, \dots, N_S\} \quad (10c)$$

$$P_{idle}^i + (P_{max}^i - P_{idle}^i) \sum_{s=1}^{N_S} u_s^i \leq \bar{P}^i \quad (10d)$$

$$0 \leq \lambda_{s,i}^i \quad (10e)$$

$$\lambda_s^{min} - \sum_{j=1}^{N_I} \lambda_{j,s}^i \leq 0, \forall s \in \{1, \dots, N_S\} \quad (10f)$$

$$\lambda_{j,s}^i R_s \leq 0.9 \mu_{j,s} C_j, \forall s \& j \quad (10g)$$

$$\sum_{j=1}^{N_I} \lambda_{j,s}^i (t_{j,s}^{sensors} + \frac{1}{\mu_{j,s}^i - \lambda_{j,s}^i} + t_{j,s}^{actuators} - t_s^{max}) \leq 0, \forall s \quad (10h)$$

$$\sigma_{j,s}^i \lambda_{j,s}^i = 0, \forall j \& s \quad (10i)$$

$$\gamma_s^i (\lambda_s^{min} - \sum_{j=1}^{N_I} \lambda_{j,s}^i) = 0, \forall s \quad (10j)$$

$$\nu_{j,s}^i (\lambda_{j,s}^i R_s - 0.9 \mu_{j,s} C_j) = 0 \forall j \& s \quad (10k)$$

$$\eta_s^i \sum_{j=1}^{N_I} \lambda_{j,s}^i (t_{j,s}^{sensors} + \frac{1}{\mu_{j,s} C_j - \lambda_{j,s}^i R_s} + t_{j,s}^{actuators} - t_s^{max}) = 0, \forall s \quad (10l)$$

$$\frac{1}{\sum_{k=1}^{N_I} \lambda_{k,s}^i} - p_j R_s + \sigma_{j,s}^i + \gamma_s^i - R_s \nu_{j,s}^i - \eta_s^i (t_{j,s}^{sensors} + t_{j,s}^{actuators} - t_s^{max} + \frac{\mu_{j,s} C_j}{\mu_{j,s} C_j - \lambda_{j,s}^i R_s}) = 0, \forall s \& j \quad (10m)$$

$$0 \leq \sigma_{j,s}^i, \sigma_{j,s}^i, \nu_{j,s}^i \& \sigma_{j,s}^i, \forall s \& j \quad (10n)$$

Here $\lambda_{s,i}$ is $[\lambda_1^i, \dots, \lambda_{N_S}^i]$ and λ_s^i is conjecture of λ_s by PaaS provider i , u^i is $[u_{i,1}, \dots, u_{i,N_S}]$ and $u_i = \sum_{i=1}^{N_S} u_{i,s}$. A potential function can be defined for this game.

Function π is potential function of this game:

$$\pi(x^i, y^i, x^{-i}, y^{-i}) = \sum_{i=1}^{N_I} P(u_i) - p_i u_i \quad (11)$$

here x^i and y^i are a tuple of (p^i, u^i) and $(\lambda_{s,i}, \nu_S^i, \gamma_S^i)$

REFERENCES