

Formation of strongly shifted EIT resonances using “forbidden” transitions of Cesium

R. Momier^{1,2,*}, A. Sargsyan², A. Tonoyan², D. Sarkisyan² and C. Leroy¹

¹Laboratoire Interdisciplinaire Carnot de Bourgogne, UMR 6303 CNRS - Université de Bourgogne, 21000 Dijon, France

²Institute for Physical Research, NAS of Armenia, Ashtarak-2, 0203 Armenia

*rodolphe.momier@u-bourgogne.fr / momier.rodolphe@gmail.com

Abstract

Atomic transitions satisfying $\Delta F = \pm 2$ of alkali atoms have zero probability in zero magnetic field (they are so-called “forbidden” transitions) but experience a large probability increase in an external magnetic field. These transitions are called magnetically induced (MI) transitions. In this paper, we use for the first time the σ^+ MI $F_g = 3 \rightarrow F_e = 5$ transitions of Cesium as probe radiation to form EIT resonances in strong magnetic fields (1 - 3 kG) while the coupling radiation frequency is resonant with $F_g = 4 \rightarrow F_e = 5$ σ^+ transitions. The experiment is performed using a nanometric-thin cell filled with Cs vapor and a strong permanent magnet. Due to the large frequency shift slope of the MI transitions (4 MHz/G), it is possible to form contrasted and strongly frequency-shifted EIT resonances. Preliminary calculations performed considering Doppler-broadened three level systems in a nanocell are in reasonable agreement with the experimental measurements.

Theory

We perform the calculations using a simple density matrix model for a three-level Λ -system:

$$\begin{aligned}\dot{\rho}_{32} &= i\Omega_c(\rho_{22} - \rho_{33}) + i\Omega_p\rho_{21} - (i\Delta_c + \gamma_{31}/2)\rho_{32} \\ \dot{\rho}_{31} &= i\Omega_c(\rho_{11} - \rho_{33}) + i\Omega_p\rho_{21} - (i\Delta_p + \gamma_{32}/2)\rho_{31} \\ \dot{\rho}_{21} &= i\Omega_c^*\rho_{31} - i\Omega_p^*\rho_{32} + (i\Delta_R + \Gamma_{21})\rho_{21}\end{aligned}$$

where Δ_c (resp. Δ_p) is the detuning of the coupling (resp. probe) laser to the upper state.




Figure 1: Three level system used in the calculations. The total decay rate $\Gamma_{33} = (\gamma_{31} + \gamma_{32})/2$. The dephasing rate of coherence between the ground state is $\Gamma_{21} = (2\pi t)^{-1}$ where t is the time of flight of the atoms through the cell (at the most probable thermal velocity).

The absorption profile is given by [2,3]:

$$\begin{aligned}\langle A \rangle &= \frac{-4\pi\omega_p N t_2^2 t_1}{cu\sqrt{u}} \frac{E_p}{|F|^2} \int_0^{+\infty} v_z M(v_z) dv_z \int_0^{L/v} dt \\ &\times \text{Im} \left\{ \mu_{31} \left[\rho_{31}^+(t, \Delta_p^+, E_{p0}(v_z t)) (1 - r_1 e^{2ik_p v_z t}) \right. \right. \\ &\left. \left. + \rho_{31}^-(t, \Delta_p^-, E_{p0}(L - v_z t)) (1 - r_1 e^{2ik_p (L - v_z t)}) \right] \right\}.\end{aligned}$$

Magnetically induced transitions





Figure 3: Magnetic field dependence of the D_2 line Zeeman transitions of Cs. a) $F_g = 3 \rightarrow F_e = 5$ σ^+ transitions. b) $F_g = 4 \rightarrow F_e = 5$ σ^+ transitions. c) Transition $[4, -1] \rightarrow [5, -2]$ (σ). This transition forms a Λ -system with the transition $[3, -3] \rightarrow [5, -2]$ (as shown in the inset). Its probability tends to 0 as the magnetic field increases, therefore forming EIT resonances at high magnetic fields requires both probe and coupling to be σ^+ -polarized.



$|F, m_F\rangle$ is just a notation for visualization, as the states would be better described in the $|J, m_J, I, m_I\rangle$ basis.

Figure 4: Scheme of the Cs D_2 line σ^+ transitions between $F_g = 3, 4$ and $F_e = 5$. The probe frequency is scanned across transitions 1 to 7, and the coupling frequencies are resonant with the $F_g = 4 \rightarrow F_e = 5$ transitions, forming seven Λ -systems. Only the states involved in the process are shown.

Experimental results




Figure 5: Probe transmission spectra of the Cs nanocell ($L = \lambda = 852$ nm). Lines 1 to 3 show four EIT resonances, labelled EIT 4, EIT 5 and EIT 6. The external longitudinal magnetic field is $B = 1770$ G on the left, and $B = 2880$ G on the right. The left part of the inset is a zoom on EIT 6, fitted with a Gaussian profile (FWHM 35 MHz) and compared with theory. The intensity of the coupling radiation was 18 mW/cm². Red: coupling on, black: coupling off. Small VSOP peaks are visible on each atomic transitions formed by the probe radiation. Their typical linewidth is 40-50 MHz. Zero frequency corresponds to the transition frequency of Cs D_2 line.

Transitions 1 to 7 (Fig. 4) can still be easily recorded up to $B = 8$ kG. In that case, their frequency shift reaches around 34 GHz from the Cs D_2 line transition frequency, allowing to use these EIT resonances as frequency reference [3].

Experimental setup




Figure 2: Scheme of the experimental setup. The orange channel allows to form EIT resonances in the absorption spectrum of Cs using a nanometric-thin cell, and the green channel is used to measure the saturated absorption (SA) of a usual long cell for reference.

References

- [1] Shore, B. W. (1990) ‘The Theory of Coherent Atomic Excitation’, Wiley
- [2] Pashayan-Leroy, Y. et al. (2007) ‘Electromagnetically induced transparency: the thickness of the vapor column is of the order of a light wavelength’, *J. Opt. Soc. Am. B*, **24**(8), pp. 1829-1838.
- [3] Nikogosyan, G.V. et al. (2004) ‘Absorption of resonance radiation and fluorescence of a layer of an atomic gas with thickness of the order of a wavelength’, *J. Opt. Technol.* **71**(9), pp. 602-607
- [4] Sargsyan, A. et al. (2023) ‘Formation of strongly shifted EIT resonances using “forbidden” transitions of Cesium’, *J. Quant. Spectrosc. Radiat. Transf.*, **303**, p. 108582.

Conclusion

We use for the first time forbidden transitions of Cs to form EIT resonances. This is made possible by using a nanometric-thin cell. Due to the behavior of the Zeeman transitions, these EIT resonances are strongly shifted with respect to the Cs D_2 line transition frequency and are good candidates to serve as frequency reference. Narrower EIT resonances could be obtained by using coherently coupled beams derived from a single laser.

Acknowledgements

This research was sponsored in part by the NATO Science for Peace and Security Programme under grant G5794 and by the Science Committee of RA in the frame of the project n°1-6/23-I/IPR. R.M. acknowledges support from the ERASMUS+ agency through the Staff Mobility for Traineeship programme.

To learn more

