


First Question: Visualization, Chi-Square and t-test

Arnau Abella

03/03/2020


- a. Analyze the distribution of “X100m” according to the type of competition by using boxplot. Write your conclusion.

```
decathlon <- read.csv( "/home/arnau/MIRI/SMDE/hw1/decathlon.csv", header = TRUE, sep = ",")
dim(decathlon)
head(decathlon)
summary(decathlon)
boxplot(X100m~Competition, ylab="x100m", xlab="competition", data=decathlon)
```



```
decastar<-which(decathlon$Competition=="Decastar")
olympic <-which(decathlon$Competition=="OlympicG")
plot(density(decathlon$X100m[decastar]),main="Density curves of X100m for Decastar and OlympicG", xlim=
lines(density(decathlon$X100m[olympic]),col=2)
```

Density curves of X100m for Decastar and OlympicG



The discret random variable X_{100m} follows a normal distribution by the shape of its density function.

Both distributions are very similar but the medians do not coincide.

Notice that we have less samples from Decastar (13) than from OlympicG (28) so we need to make some probabilistic analysis before refusing that the medians are equals.

From the statistics, it is easy to see that, in average, the runners from OlympicG are one second faster than the ones from Decastar.

- b. Create a new categorical variable with two categories from the variable “X100m” by using 11 seconds as the cut-off point. Make a cross table from the new categorical variable and the “Competition”. Are these two variables independent? Write your conclusion by checking marginal probabilities and test the independency of two variables by using Chi-Square test.

```
decathlon$X100m_11s<- cut(decathlon$X100m, c(11,0 , 12.0))
levels(decathlon$X100m_11s)<-c("< 11s","> 11s")
tab<-table(decathlon$X100m_11s, decathlon$Competition)
tab
```

```
##
##      Decastar  OlympicG
## < 11s         2         19
## > 11s        11         9
```

```
prop.table(tab)
```

```
##
##      Decastar  OlympicG
## < 11s 0.04878049 0.46341463
## > 11s 0.26829268 0.21951220
```

```
chisq.test(tab) # Independent and normally
```


```
##
## Pearson's Chi-squared test with Yates' continuity correction
##
## data:  tab
## X-squared = 7.7962, df = 1, p-value = 0.005236
```

As the p-value 0.0052 from χ^2 is smaller than the .05 significance level, we can reject the null hypothesis that running 100 meters in less/more than 11 seconds is independent of the kind of competition.

- c. Visualize the distribution of quantitative variables by using proper graph. Which of these variables follows a Normal distribution?


All quantitative variables follow a Normal distribution except for **Rank** and **Points**. Here are some sample plots:

```
plot(density(decathlon$X100m), main="Probability density function: X100m")
```



```
plot(density(decathlon$X400m), main="Probability density function: X400m")
```


Probability density function: X400m



N = 41 Bandwidth = 0.4378

```
plot(density(decathlon$Javeline), main="Probability density function: Javeline")
```

Probability density function: Javeline




N = 41 Bandwidth = 1.796

To check that Rank and Position do not follow a Normal distribution we used qqnorm and Shapiro-Wilk normality test.

```
plot(density(decathlon$Rank), main="Probability density function: X100m")
```


Probability density function: X100m



N = 41 Bandwidth = 3.391

```
qqnorm(decathlon$Rank); qqline(decathlon$Rank, col = 2)
```

Normal Q-Q Plot



```
shapiro.test(decathlon$Rank)
```


```
##  
##  Shapiro-Wilk normality test  
##  
## data:  decathlon$Rank  
## W = 0.94188, p-value = 0.03649
```

- d. Generate three Normally distributed random variables of length 50. Two of them should have the same mean, different standard deviations while the third one has a different mean but the same standard deviation with the first distribution. Use t test to compare mean differences between three variables.

```
s1<-rnorm(50, mean=25,sd=4)
s2<-rnorm(50, mean=25,sd=8)
s3<-rnorm(50, mean=35,sd=4)
```

```
plot(density(s1),xlim=c(0,50),ylim=c(0,0.13), main="Three different Normal distributed random variables",
lines(density(s2),col=2)
lines(density(s3),col=3)
```

Three different Normal distributed random variables



Let's compare the mean difference using t-test:

```
t.test(s1, s2, var.equal=TRUE)
```

```
##
## Two Sample t-test
##
## data: s1 and s2
## t = 1.1862, df = 98, p-value = 0.2384
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
## -1.070083 4.250051
## sample estimates:
## mean of x mean of y
## 25.62440 24.03441
```

We can't refuse that the mean of **s1** and **s2** are different because the p-value 0.56 is greater than α . However, we can refuse this for **s1** and **s3** as expected because the p-value is way slower than α .


```
t.test(s1, s3, var.equal=TRUE)
```

```
##  
## Two Sample t-test  
##  
## data: s1 and s3  
## t = -11.264, df = 98, p-value < 2.2e-16  
## alternative hypothesis: true difference in means is not equal to 0  
## 95 percent confidence interval:  
## -10.83724 -7.59059  
## sample estimates:  
## mean of x mean of y  
## 25.62440 34.83832
```

Additionally, we can test the variance of the distributions. The distributions **s1** and **s2** have different variance as expected.

```
var.test(s1, s2, var.equal=TRUE)
```

```
##  
## F test to compare two variances  
##  
## data: s1 and s2  
## F = 0.28903, num df = 49, denom df = 49, p-value = 2.73e-05  
## alternative hypothesis: true ratio of variances is not equal to 1  
## 95 percent confidence interval:  
## 0.1640200 0.5093327  
## sample estimates:  
## ratio of variances  
## 0.2890342
```

- e. Test if there is a difference between two type of competitions according to the variables “X100m” and “X400m” by using t-test.

After performing the Student’s t-test on the *X100m*, we **can** refuse the null hypothesis H_0 that there is no significant difference in the means because the p-value 0.00407 is smaller than α . So we can state with 95% of confidence that there is a statistical difference in the times for the 100 meters race dependening on the competition type.

```
t.test(X100m ~ Competition, data = decathlon)

##
##  Welch Two Sample t-test
##
## data:  X100m by Competition
## t = 3.2037, df = 22.168, p-value = 0.00407
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  0.09164794 0.42769272
## sample estimates:
## mean in group Decastar mean in group OlympicG
##           11.17538           10.91571
```

After performing the Student’s t-test on the *X400m*, we **can’t** refuse the null hypothesis H_0 that there is no significant difference in the means because the p-value 0.9543 is greater than α . So there is no significative evidence that the mean time on the 400 meters race is different depending on the competition type.

```
t.test(X400m ~ Competition, data = decathlon)

##
##  Welch Two Sample t-test
##
## data:  X400m by Competition
## t = 0.05771, df = 32.106, p-value = 0.9543
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  -0.6858299 0.7258299
## sample estimates:
## mean in group Decastar mean in group OlympicG
##           49.63           49.61
```