Neurodevelopmental Disorder With Hypotonia, Seizures, and Absent Language (NDHSAL): A Comprehensive Disease Profile

Disease: Neurodevelopmental Disorder With Hypotonia, Seizures, And Absent Language (NDHSAL) **MONDO:** MONDO:0014995 · **OMIM:** #617268 · **Causal gene: HECW2** (HGNC:29853; NCBI Gene 57520; OMIM gene 617245)

Summary

Neurodevelopmental Disorder With Hypotonia, Seizures, and Absent Language (NDHSAL) is an ultra-rare Mendelian neurodevelopmental disorder caused by variants in **HECW2 — HECT, C2 and WW domain-containing E3 ubiquitin protein ligase 2** (alias **NEDL2**). HECW2 encodes a HECT-type E3 ubiquitin ligase that ubiquitinates and **stabilizes the transcription factor p73 (TP73)**, a crucial mediator of neurogenesis and brain development. The disorder was first defined by Berko et al. in 2017, who identified de novo deleterious missense variants in HECW2 in probands with intellectual disability, developmental delay, and hypotonia (PMID: 27389779).

Clinically, NDHSAL presents with a **completely penetrant core triad** — neonatal/congenital **hypotonia** (with or without spasticity), **global developmental delay/intellectual disability**, and **developmental language disorder/absent language** (each 100% in the largest cohort) — accompanied by frequent **seizures (~61%)**, behavioural problems (88.9%), vision problems (83.9%), motor/coordination problems (75%), and gastrointestinal issues (70%) (PMID: 34321324). A subset develop severe **developmental and epileptic encephalopathy (DEE)**, including West syndrome/infantile spasms and burst-suppression patterns, and rare cases show **cardiac long-QT syndrome** (PMID: 37280227).

Two genetic mechanisms operate. The **predominant form is autosomal dominant**, driven by **de novo missense variants clustered in or near the C-terminal HECT catalytic domain (88.2%)** acting through a **dominant-negative** mechanism (PMID: 34047014). A **rarer autosomal recessive** form arises from **biallelic loss-of-function/hypomorphic variants** in consanguineous families (PMID: 35487419; PMID: 35753050; PMID: 40812465). HECW2 is one of the most constraint-intolerant genes in the genome (pLI ≈ 1, missense Z = 4.54). There is **no disease-modifying or gene-targeted therapy**; management is entirely **symptomatic** (antiseizure drugs, ACTH/ketogenic diet, perampanel for refractory DEE, rehabilitation, and cardiac surveillance). **No validated animal model and no biochemical biomarker** currently exist.

> **Evidence-base note:** This report synthesizes primary human clinical literature (case reports and one ~35-patient multicenter cohort) plus authoritative molecular/population databases (UniProt, NCBI Gene, Ensembl, gnomAD, ClinVar, HomoloGene) queried live during the investigation. NDHSAL is ultra-rare; most statements derive from aggregated small-cohort/case data, not large registries. Where a mechanistic step is inferred rather than experimentally demonstrated in the disease context, this is stated explicitly.

Section 1 — Disease Information

NDHSAL is a Mendelian neurodevelopmental disorder characterized by the symptom triad embedded in its name. The information base is largely **individual-patient derived** (case reports and one multicenter cohort of ~35 individuals) aggregated into disease-level resources (OMIM, MONDO).

Key identifiers:

Resource	Identifier
MONDO	MONDO:0014995
OMIM (phenotype)	#617268
OMIM (gene)	617245 (HECW2)
HGNC	29853
NCBI Gene	57520
Ensembl	ENSG00000138411
UniProt	Q9P2P5
Cytoband	2q32.3

Synonyms: NDHSAL; HECW2-related neurodevelopmental disorder; HECW2-related disorder; neurodevelopmental disorder with hypotonia, seizures, and absent language. Gene aliases: **NEDL2**, NDHSAL.

ICD-10/ICD-11, MeSH and Orphanet do not carry a dedicated, disease-specific code for this ultra-rare entity; it is captured under broad neurodevelopmental/ID categories.

Section 2 — Etiology

Primary cause — genetic. NDHSAL is monogenic, caused by pathogenic variants in **HECW2**. There is **no known environmental, infectious, toxic, or lifestyle cause**, and no gene–environment interaction has been described. No environmental or genetic **protective factors** are documented; given the de novo dominant mechanism, classical protective-allele epidemiology is not applicable.

Genetic risk / causal variants. - **Autosomal dominant (most common):** heterozygous **de novo missense** variants, predominantly in or near the C-terminal HECT domain; dominant-negative effect (PMID: 34047014). - **Autosomal recessive (rare):** biallelic loss-of-function/ hypomorphic variants (nonsense, splice, frameshift) in consanguineous families (PMID: 35487419; PMID: 35753050; PMID: 40812465).

Risk-factor context. The only established risk factor for the recessive form is **consanguinity**. For the dominant form, **de novo mutation** (sporadic) is the rule, with advanced parental age a general — but unproven-specific — background risk for de novo events. No modifier genes have been identified.

Section 3 — Phenotypes

Phenotype frequencies are drawn principally from the **35-case cohort of Acharya et al. 2022** (PMID: 34321324) and the founding series (PMID: 27389779).

Phenotype	Type	Frequency	Suggested HPO term
Hypotonia (± spasticity)	Clinical sign	100%	HP:0001252 (± HP:0001257)
Developmental delay / intellectual disability	Clinical/ behavioural	100%	HP:0001263 / HP:0001249
Developmental language disorder / absent language	Behavioural/ clinical	100%	HP:0000750 / HP:0001344
Behavioural problems	Behavioural	88.9%	HP:0000708
Vision problems (incl. cortical visual impairment)	Clinical sign	83.9%	HP:0100704 / HP:0000505
Motor coordination/movement problems	Clinical sign	75%	HP:0001310 / HP:0002066
Gastrointestinal issues (incl. GERD)	Clinical sign	70%	HP:0011024 / HP:0002020
Seizures	Clinical sign	61.3%	HP:0001250
Strabismus	Clinical sign	Common	HP:0000486
Nystagmus	Clinical sign	Common	HP:0000639
Dysmorphic facial features	Physical	Common	HP:0001999
Microcephaly (severe cases)	Physical	Subset	HP:0000252
Cerebral/cerebellar atrophy (severe cases)	Imaging	Subset	HP:0002059 / HP:0001272
Long QT syndrome (rare)	Lab/ECG	Rare	HP:0001657

Berko et al. summarize the associated features: *"Other common features include seizures, strabismus, nystagmus, cortical visual impairment and dysmorphic facial features"* (PMID: 27389779). Acharya et al. quantify penetrance: *"Clinical characterisation suggests complete penetrance for hypotonia with or without spasticity (100%), developmental delay/intellectual disability (100%) and developmental language disorder (100%). Other common features are*

behavioural problems (88.9%), vision problems (83.9%), motor coordination/movement (75%) and gastrointestinal issues (70%). Seizures were present in 61.3% of individuals" (PMID: 34321324).

Characteristics. Onset is **congenital/neonatal** (hypotonia) to **infancy** (seizures, developmental delay). Severity is **variable but generally moderate-to-severe**; the disease is **chronic and lifelong** rather than classically neurodegenerative, although DEE presentations can show early cerebral/cerebellar atrophy and drug-resistant epilepsy. **Quality-of-life impact** is substantial: absent/severely limited language, frequent non-ambulatory status, and intellectual disability imply lifelong dependence for daily functioning; no formal EQ-5D/SF-36 data exist for this rare disorder.

Section 4 — Genetic / Molecular Information

Causal gene: HECW2 (HGNC:29853; NCBI Gene 57520; Ensembl ENSG00000138411; cytoband 2q32.3; UniProt Q9P2P5, 1572 aa).

Variant spectrum. In the autosomal dominant form **all reported variants are missense**, and **88.2%** of likely-pathogenic/pathogenic variants lie in or near the C-terminal HECT domain (PMID: 34321324). Four **recurrent variants** anchor the hotspot:

Variant	Note
p.(Arg1191Gln)	~22.9% of cases
p.(Arg1330Trp)	~20% of cases
p.(Asn1199Lys)	Recurrent
p.(Phe1327Ser)	Recurrent

Recessive variants include homozygous nonsense **p.Arg246*** (PMID: 35487419), a homozygous **splice-donor** variant eliminating exon 22 (p.Leu1256_Trp1306del; ~60% reduced protein) (PMID: 35753050), and a homozygous **frameshift p.Y1201Lfs*7** predicted to undergo nonsense-mediated decay (PMID: 40812465).

Variant classification (ClinVar, live query of HECW2[gene], 732 records): 4 Pathogenic, 11 Likely pathogenic, 7 Conflicting, 237 Uncertain significance, 122 Likely benign, 2 Benign. The vast majority are single-nucleotide variants; a handful are small deletions/duplications/CNVs.

Allele frequency / constraint. Pathogenic alleles are **essentially absent from gnomAD** (private de novo or rare biallelic events). HECW2 is among the most **constraint-intolerant** genes: gnomAD **pLI = 0.99999**, **LOEUF = 0.43**, **missense Z = 4.54**, **LoF Z = 7.32**, observed/expected missense = 0.76.

Functional consequences. The dominant missense variants act by a **dominant-negative** mechanism on the HECT catalytic machinery; recessive variants act by **loss of function** (NMD or reduced protein). No modifier genes, no disease-specific epigenetic signature, and no recurrent chromosomal abnormality are established (HECW2 is not detectable by karyotype and rarely implicated by CMA).

Section 5 — Environmental Information


Not applicable. NDHSAL is a monogenic disorder. No environmental toxins, radiation, pollution, occupational exposures, lifestyle factors, or infectious agents are implicated in its causation or triggering. The only non-genetic contextual factor is **consanguinity**, which raises the probability of the recessive form.

Section 6 — Mechanism / Pathophysiology

Ordered causal chain

1. A **HECW2 variant** (de novo heterozygous missense in the HECT domain, or biallelic LoF) **alters the E3 ubiquitin ligase activity** of HECW2/NEDL2.
2. *Dominant branch:* the HECT-domain missense product **results in** a **dominant-negative** enzyme that poisons normal ligase function (inferred from exclusively missense, HECT-clustered variants with no LoF in the AD form; PMID: 34047014).
3. *Recessive branch:* biallelic LoF variants **lead to absent/reduced HECW2 protein** (NMD or ~60% reduced expression; PMID: 40812465; PMID: 35753050).
4. Altered HECW2 activity **leads to dysregulated ubiquitination and reduced stabilization of p73 (TP73)** (inferred from HECW2's established role as a p73-stabilizing ligase; PMID: 27389779; PMID: 35487419).

5. Reduced p73 stability/transcriptional activity **results in impaired p73-dependent programs of neuronal proliferation, apoptosis, and neuronal differentiation** essential for brain development (PMID: 35487419).
6. Disrupted neurodevelopment **leads to abnormal cortical/cerebellar development and neuronal network dysfunction** (manifest in severe cases as cerebral/cerebellar atrophy and epileptogenic networks).
7. These developmental and network abnormalities **result in** the **clinical phenotype**: hypotonia, developmental delay/ID, absent language, seizures/DEE, visual and GI dysfunction; and, via HECW2's broad substrate roles, occasional **cardiac long-QT** (PMID: 37280227).



Supporting molecular detail

- **Molecular pathways / protein function:** HECW2/NEDL2 is a **NEDD4-family HECT E3 ubiquitin ligase**. UniProt Q9P2P5 (1572 aa) has an N-terminal **C2 domain** (aa 167–301; membrane/Ca^{2+} targeting), two **WW domains** (aa 807–840 and 985–1018; PPxY substrate binding), and a C-terminal **HECT catalytic domain** (aa 1237–1572) with the catalytic cysteine (~C1540) forming the ubiquitin thioester intermediate. A defined **TP73-interaction region** spans aa 737–1068.

- Berko et al.: *"HECW2 is an ubiquitin ligase that stabilises p73, a crucial mediator of neurodevelopment and neurogenesis. This study implicates pathogenic genetic variants in HECW2 as potential causes of neurodevelopmental disorders in humans"* (PMID: 27389779).
- A subsequent report: *"HECW2 encodes an E3 ubiquitin-protein ligase that stabilizes and enhances transcriptional activity of p73, a key factor regulating proliferation, apoptosis, and neuronal differentiation, which are together essential for proper brain development"* (PMID: 35487419).
- **Cellular processes:** Beyond p73, HECW2 regulates the **mitotic metaphase/anaphase transition**, interacting with FZR1/Cdh1 and being degraded by APC/C-Cdh1 at mitotic exit — consistent with a role in **cell-cycle control** relevant to neurogenesis.
- **Subcellular localization (GO cellular component):** cytoplasm (GO:0005737) and cytoskeleton/**mitotic spindle** (GO:0072686).
- **Suggested GO biological processes:** protein polyubiquitination (GO:0000209), protein stabilization (GO:0050821), nervous system development (GO:0007399), neuron differentiation (GO:0030182), regulation of mitotic cell cycle (GO:0007346).
- **Suggested CL cell types:** neuron (CL:0000540), glutamatergic neuron (CL:0000679), cerebellar Purkinje cell (CL:0000121), neural progenitor cell (CL:0011020).

No metabolomic, proteomic, lipidomic, single-cell, spatial, or CRISPR-screen data specific to NDHSAL are available; disease-variant characterization to date is in vitro/in silico.

Section 7 — Anatomical Structures Affected

- **Primary organ / body system:** central nervous system (**brain**, UBERON:0000955; nervous system, UBERON:0001016). Involved regions include cerebral cortex (UBERON:0000956) and cerebellum (UBERON:0002037) — both showing atrophy in severe cases.
- **Secondary involvement:** eye/visual pathway (cortical visual impairment, strabismus, nystagmus; UBERON:0000970), gastrointestinal tract (GERD; UBERON:0001555), and **heart** (UBERON:0000948) in rare long-QT cases.
- **Tissue/cell level:** nervous tissue; affected cell populations include **neurons** (CL:0000540) and **neural progenitors** (CL:0011020); HECW2 is expressed predominantly in adult **brain, lung, and heart**.
- **Subcellular level (GO CC):** cytoplasm (GO:0005737), mitotic spindle (GO:0072686), nucleus-associated transcriptional regulation via p73.

- **Lateralization:** CNS involvement is **bilateral/symmetric**.

Section 8 — Temporal Development

- **Onset:** congenital/neonatal **hypotonia**; infantile-onset **seizures** and **developmental delay**. Pattern is typically **chronic/insidious** developmental, with acute seizure presentations in DEE.
- **Progression:** Overall **static-to-slowly-evolving** encephalopathy — **not classically neurodegenerative**. However, severe DEE presentations evolve rapidly in infancy (e.g., burst-suppression → myoclonic → spasms; PMID: 40879666) and may show early cerebral/cerebellar atrophy (PMID: 40812465).
- **Course / duration: lifelong, chronic** disability.
- **Critical periods:** infancy/early childhood is the key window for seizure control and developmental intervention (physiotherapy, speech therapy).
- **Remission:** no spontaneous remission of the neurodevelopmental disorder; seizures may be partially controlled but are frequently **drug-resistant**.

Section 9 — Inheritance and Population

- **Epidemiology:** ultra-rare; **no formal prevalence or incidence** estimates exist. The literature comprises case reports plus one multicenter cohort (~35 individuals; PMID: 34321324), so **fewer than ~100 patients** are reported worldwide. No Orphanet prevalence class is established.
- **Inheritance:** predominantly **autosomal dominant** from **de novo heterozygous missense** variants (18/22 confirmed de novo in Acharya 2022). Rare **autosomal recessive** forms arise from biallelic LoF/hypomorphic variants in consanguineous families (PMID: 35487419; PMID: 35753050; PMID: 40812465).
- **Penetrance: complete (100%)** for hypotonia, DD/ID, and language disorder.
- **Expressivity: variable**, including marked intrafamilial variability for a transmitted splice variant (PMID: 40879666).
- **Anticipation:** none (no repeat-expansion mechanism).

- **Mosaicism:** germline mosaicism is plausible (as for other de novo disorders) but not specifically documented.
- **Founder effects / consanguinity:** no founder allele; **consanguinity** underlies recessive cases.
- **Carrier frequency:** not established; pathogenic alleles are essentially absent from population databases.
- **Sex ratio:** no clear sex predilection. **Geographic distribution:** worldwide, no endemic region.

Section 10 — Diagnostics

Molecular diagnosis is definitive. Because variants are private de novo missense (or rare biallelic), **trio whole-exome sequencing (WES)** is the highest-yield test (18/22 de novo in Acharya 2022; trio-WES diagnoses also in PMID: 40469502). **Whole-genome sequencing (WGS)** and **multigene epilepsy/ID/DEE NGS panels** are alternatives; **single-gene testing** supports familial/cascade confirmation. HECW2 is **not detectable by karyotype** and is **rarely implicated by chromosomal microarray** (occasional 2q32.3 CNVs in ClinVar).

Variant interpretation challenge: ClinVar shows only **4 Pathogenic + 11 Likely pathogenic** against **237 VUS** and 122 Likely benign (732 records), underscoring the burden of uncertain variants and the value of de novo status and HECT-domain localization in ACMG/AMP classification.

Supportive diagnostics: - **EEG** to characterize seizures (burst-suppression, hypsarrhythmia/spasms; PMID: 40879666, PMID: 40705299). - **Brain MRI** (cerebral/cerebellar atrophy, nonspecific anomalies; PMID: 40812465, PMID: 35753050). - **Ophthalmologic assessment** (cortical visual impairment, strabismus, nystagmus). - **ECG** for QT monitoring (PMID: 37280227).

No specific biochemical/metabolite biomarker exists; there is no newborn-screening test. **Differential diagnosis** includes other genetic hypotonia–DEE–ID syndromes; distinction is made by gene identification.

Section 11 — Outcome / Prognosis

- **Survival/mortality:** no published survival statistics. The disorder is **not intrinsically lethal**, but severe DEE and **rare cardiac long-QT** (PMID: 37280227) can be **life-threatening**.
- **Morbidity/disability:** high and lifelong — **absent/severely limited language**, intellectual disability, and frequently **non-ambulatory** status, implying dependence for activities of daily living.
- **Course:** chronic, largely static-to-slowly-evolving; **drug-resistant epilepsy** worsens prognosis in a subset.
- **Prognostic factors:** HECT-domain variants correlate with cortical visual impairment and GI issues; **seizures were observed only with variants in or near the HECT domain**. Severe/DEE presentations with early atrophy portend worse developmental outcomes.
- **No validated prognostic biomarker** exists.

Section 12 — Treatment

There is no disease-modifying or gene-targeted therapy. Management is entirely symptomatic and supportive.

Domain	Intervention	Evidence	NCIT suggestion
Epilepsy (general)	Antiseizure medications; often drug-resistant	PMID: 35753050	NCIT:C264 (Anticonvulsant Agent)
West syndrome / spasms	ACTH + modified Atkins (ketogenic-type) diet	PMID: 40705299	NCIT:C2366 (ACTH); NCIT:C92948 (Ketogenic Diet)
Refractory DEE	Perampanel (AMPA-receptor antagonist), personalized	PMID: 40879666	NCIT:C73043 (Perampanel)
Hypotonia/ motor	Physiotherapy	—	NCIT:C15415 (Physical Therapy)
Language	Speech therapy	—	NCIT:C15401 (Speech Therapy)
Vision	Ophthalmologic management	—	—
GI (GERD)	Reflux management	PMID: 35753050	—
Cardiac	QT surveillance / LQTS management	PMID: 37280227	—

ACTH plus ketogenic-type diet was reported with benefit in pharmacoresistant HECW2-related West syndrome: *"We describe an infant with a pathogenic HECW2 gene variation who presented with excessive floppiness, global delay, tonic eye deviations, and epileptic spasms treated with a combination of adrenocorticotropic hormone and modified Atkin's diet"* (PMID: 40705299). Perampanel exemplifies personalized strategy: *"It also underscores the importance of personalized therapeutic strategies, as demonstrated by the use of perampanel in this patient"* (PMID: 40879666).

No pharmacogenomic, gene, cell, RNA, targeted, or immunotherapy is established, and **no registered clinical trials** specific to NDHSAL exist.

Section 13 — Prevention

Because NDHSAL is monogenic with no environmental contribution, prevention is **genetic/reproductive**, not public-health. - **Primary prevention:** none for the dominant de novo form (sporadic, unpredictable). - **Genetic counseling:** central — recurrence risk is low for de novo AD cases (but germline mosaicism cannot be fully excluded), and **25%** for biallelic recessive families. - **Prenatal / preimplantation genetic testing** is possible once a familial variant is known (especially in consanguineous/recessive families). - **Cascade testing** of relatives for known familial variants. - **Tertiary prevention:** minimizing complications via seizure control, cardiac QT surveillance, nutritional/GI management, and developmental therapies. - **No immunization, population screening, or behavioural/environmental intervention** is applicable.

Section 14 — Other Species / Natural Disease

HECW2 is **evolutionarily conserved** with one-to-one orthologs across vertebrates (HomoloGene group 66192):

Species	Gene	NCBI Gene ID
Mouse (*Mus musculus*)	Hecw2	329152
Rat (*Rattus norvegicus*)	Hecw2	316395
Zebrafish (*Danio rerio*)	hecw2	100331355
Chicken (*Gallus gallus*)	HECW2	424050

Additional orthologs exist in dog, cow, macaque, and chimpanzee. HECW2/NEDL2 belongs to the **NEDD4/HECT E3 ligase family**. **No naturally occurring NDHSAL-equivalent disease** is documented in companion animals or wildlife (no OMIA entry equivalent to the human disorder), and there is **no zoonotic or transmission dimension** (it is a genetic disorder).


Section 15 — Model Organisms

No published knockout/knock-in mouse, rat, or zebrafish model specifically recapitulates the human NDHSAL phenotype as of this review. The founding functional studies defined NEDL2 as a **p73-stabilizing ubiquitin ligase** in cell systems and a regulator of the **APC/C-Cdh1 mitotic transition**, i.e., **in vitro/cell-based** characterization. Disease-variant work to date is in vitro/in silico (e.g., ~60% reduced protein for the recessive splice variant, PMID: 35753050; NMD prediction for the frameshift, PMID: 40812465).

Model resources for future work: MGI/IMPC/KOMP (mouse Hecw2, Gene 329152), RGD (rat), ZFIN (zebrafish hecw2). Conserved orthology makes mouse and zebrafish strong candidate systems. **Limitation:** absence of any validated model means genotype–phenotype mechanisms (dominant-negative vs. LoF) and candidate therapeutics cannot yet be tested in vivo.

Mechanistic Model / Interpretation

The unifying model is **p73 dysregulation downstream of a defective HECT ubiquitin ligase**. HECW2 normally ubiquitinates and **stabilizes p73**, amplifying a transcriptional program (proliferation, apoptosis, neuronal differentiation) that brain development depends upon. Two mutational routes converge on reduced effective p73 activity:



The **genotype–phenotype correlation** is informative: variants in/near the HECT domain are associated with cortical visual impairment and GI issues, and **seizures occur only with HECT-region variants**, implying that the catalytic core is the most phenotypically consequential region (PMID: 34321324). The extreme population constraint (pLI ≈ 1, missense Z = 4.54) independently corroborates that both dosage reduction and missense perturbation of HECW2 are strongly deleterious — fully consistent with the observed AD (dominant-negative) and AR (LoF) mechanisms. Yanagishita et al. articulate the dominant branch directly: *"All HECW2*

variants, including those that were previously reported, are missense, and no loss-of-function variants have been identified. Most of the identified variants are located around the HECT domain. These findings suggest that the dominant negative effects of missense variants around the HECT domain may be the mechanism underlying HECW2-related disorder" (PMID: 34047014). The recessive branch is explicitly distinguished: *"This frameshift variant likely results in nonsense-mediated decay (NMD) of the mRNA, suggesting a loss-of-function mechanism distinct from the mechanism implicated in missense variants that causing the disease with autosomal dominant inheritance"* (PMID: 40812465).

Evidence Base

PMID	Title (abbrev.)	Contribution
27389779	De novo missense variants in HECW2 (Berko 2017)	Disease-defining paper; names HECW2 as causal; p73-stabilizing ligase; lists core/associated phenotypes
34321324	Delineating genotype/ phenotype spectrum (Acharya 2022)	35-case cohort; penetrance 100% for triad; HECT hotspot 88.2%; seizures 61.3%; genotype–phenotype correlation
34047014	HECW2-related disorder in 4 Japanese patients	Dominant-negative, HECT-clustered missense; no LoF in AD form
35487419	Homozygous nonsense HECW2 variant	Recessive form; p.Arg246*; p73/brain-development mechanism
35753050	First splicing variant, AR inheritance	AR splice variant; ~60% reduced protein (partial LoF); drug-resistant epilepsy, GERD, MRI anomalies
40812465	AR frameshift broadens spectrum	p.Y1201Lfs*7; NMD/LoF mechanism distinct from AD missense; microcephaly, atrophy
37280227	Novel variant, congenital long QT	Cardiac LQTS comorbidity (p.Leu1448Ser)
40705299	HECW2 as a cause of West syndrome	ACTH + modified Atkins diet management
40879666	Rapid electroclinical evolution in DEE	Perampanel; familial transmission; intrafamilial variability
40469502	Novel variant, Chinese girl	Trio-WES diagnosis; DLD, hypotonia

Context papers on HECW2 substrate biology in non-neural settings (e.g., NSCLC via SCNN1A/ AKT-mTOR, PMID: 42537707; ATR degradation after lamin misexpression, PMID: 22382637; infantile hemangioma glycolysis, PMID: 40520873) confirm HECW2 is a broadly acting K48-linked E3 ligase but are **not disease-specific** to NDHSAL.

Limitations and Knowledge Gaps

1. **Small evidence base:** fewer than ~100 reported patients; no prevalence/incidence, survival, or validated QoL data.
2. **No in vivo model:** the dominant-negative vs. LoF mechanisms are inferred from variant distribution and in vitro data, not demonstrated in an animal model.
3. **Direct p73-pathway link unproven in patient neurons:** the p73 mechanism is extrapolated from cell-line biology; patient-derived neuronal/organoid confirmation is lacking.
4. **Variant interpretation burden:** 237 VUS in ClinVar; functional assays for ligase activity are not routine.
5. **No biomarker and no disease-modifying therapy;** management evidence is anecdotal (single cases).
6. **Recessive form rare and under-characterized;** genotype–phenotype rules for biallelic variants are preliminary.

Proposed Follow-up Experiments / Actions

1. **Generate in vivo models:** create *Hecw2* knock-in mice/zebrafish carrying recurrent HECT-domain missense alleles (e.g., p.Arg1191Gln, p.Arg1330Trp) and biallelic LoF alleles to test dominant-negative vs. LoF mechanisms and recapitulate hypotonia/seizures.
2. **Patient-derived iPSC neurons/organoids:** quantify p73 stability and downstream transcriptional targets in mutant vs. control neurons; test whether p73 activity can be pharmacologically restored.
3. **Functional ligase assays:** establish a standardized in vitro ubiquitination/auto-ubiquitination assay to reclassify the 237 VUS and confirm dominant-negative behavior of HECT-domain missenses.
4. **Natural history registry:** build an international registry to estimate prevalence, define developmental trajectories, seizure outcomes, and cardiac risk, and to standardize QoL assessment.
5. **Cardiac phenotyping:** systematic ECG screening across the cohort to determine the true frequency of QT prolongation and inform surveillance guidelines.

6. **Therapeutic screening:** once a model exists, screen antiseizure agents (including perampanel) and p73-pathway modulators for genotype-specific efficacy.

Report compiled from 5 investigation iterations, 11 confirmed findings, and 18 reviewed papers. Evidence source types span human clinical case reports/cohorts, in vitro/cell-based functional studies, and computational/population-genetics analyses (gnomAD, ClinVar, UniProt, HomoloGene).
