

Joint Laxity, Short Stature, and Myopia (GZF1-Related Phenotype): Comprehensive Disease Report

OMIM #617662 · Autosomal recessive · Mendelian connective-tissue disorder

Summary

Joint Laxity, Short Stature, and Myopia (OMIM #617662), also referred to as the **GZF1-related phenotype (GZF1RP)** or **autosomal-recessive Larsen-like syndrome**, is an ultra-rare Mendelian connective-tissue disorder caused by **biallelic loss-of-function variants in the GZF1 gene** (GDNF-inducible zinc finger protein 1; chromosome 20p11.21). It was first defined molecularly in 2017 in two multiplex consanguineous Saudi families ascertained through combined autozygome-plus-exome analysis, and the clinical and genetic spectrum was expanded in 2026. As of 2026, only ~13 patients from ~6 families have been reported worldwide, all carrying biallelic *GZF1* variants — establishing autosomal recessive inheritance with apparently high penetrance in biallelic carriers.

The disorder is characterized by a recognizable triad of **severe/high myopia** (with retinal detachment and congenital glaucoma), **short stature**, and **joint hypermobility with recurrent large-joint dislocation**, accompanied by facial dysmorphism, scoliosis, thoracic deformity, progressive sensorineural/mixed hearing loss, umbilical hernia, and hypodontia. The severe ocular involvement distinguishes GZF1RP from classic *FLNB*-related Larsen syndrome. GZF1 is a BTB/POZ-domain transcriptional repressor bearing 10 tandem zinc-finger motifs; it is induced downstream of GDNF/RET signaling and is expressed in the developing eye and limb — the very tissues affected in patients.

The proposed pathomechanism runs from **GZF1 loss of function → dysregulation of extracellular-matrix/collagen genes → most notably *P3H2* (LEPREL1)-mediated collagen under-3-hydroxylation → structurally defective scleral and ocular collagen → axial elongation and high myopia**, with parallel connective-tissue laxity producing joint and skeletal manifestations. Because *P3H2/LEPREL1* biallelic loss is an independently established

cause of non-syndromic high myopia, it provides a strong mechanistic anchor connecting GZF1 dysregulation to the ocular phenotype. There is no disease-modifying therapy; management is entirely supportive and multidisciplinary (ophthalmology, orthopedics, audiology, genetic counseling).

1. Disease Information

Overview. Joint Laxity, Short Stature, and Myopia is a Mendelian, autosomal-recessive connective-tissue disorder combining ocular and skeletal manifestations. The disease was delineated as a distinct entity when biallelic *GZF1* variants were identified as its cause, situating it within — yet distinct from — the Larsen syndrome spectrum.

Key identifiers.

Resource	Identifier
OMIM (phenotype)	#617662 (Joint laxity, short stature, and myopia)
Gene	<i>GZF1</i> (GDNF-inducible zinc finger protein 1)
Cytoband	20p11.21
MONDO	Corresponds to the GZF1-related phenotype (MONDO ID not asserted in outline; map to OMIM:617662)
Category	Mendelian, autosomal recessive

Synonyms / alternative names. GZF1-related phenotype (GZF1RP); GZF1-related ocular and skeletal disorder; autosomal-recessive Larsen-like syndrome; "a specific ocular and skeletal disorder distinguishable from Larsen syndrome" [PMID: 42170786].

Source of information. Information is derived from **individual patient reports and small case series** (aggregated at the disease level through OMIM), not from large EHR or population registries — reflecting the ultra-rare status of the condition.

2. Etiology

Primary cause — genetic. The disorder is caused by **biallelic (homozygous or compound heterozygous) loss-of-function variants in *GZF1***. The first families were consanguineous, and the causal homozygous truncating variant was identified via combined autozygome and exome analysis: *"In a multiplex consanguineous Saudi family affected by severe and recurrent large joint dislocation and severe myopia, we identified a homozygous truncating variant in *GZF1* through a combined autozygome and exome approach"* [PMID: 28475863].

Genetic risk factors. - **Causal variants:** biallelic truncating/frameshift *GZF1* variants (see Section 4). - **Consanguinity:** a major risk/ascertainment factor, given the recessive inheritance and multiplex consanguineous pedigrees. - **Recurrent alleles:** the c.1440del (p.His481IlefsTer26) variant was seen in homozygosity in two sisters and in compound heterozygosity in a third patient, suggesting possible recurrent/founder alleles [PMID: 42170786].

Environmental risk factors. None identified. This is a monogenic disorder without established environmental contributors.

Protective factors. None described. As a fully penetrant recessive disorder in biallelic carriers, no protective genetic or environmental modifiers have been reported.

Gene–environment interactions. None reported; the phenotype appears determined by genotype.

3. Phenotypes

Across all reported patients, the recurrent **core triad** is short stature, large-joint dislocation/joint hypermobility, and severe/high myopia [PMID: 28475863;

P 33009817

P 42170786

The 2026 series defined a recognizable *GZF1*RP: *"The comparison allows us to define a recognizable *GZF1*RP that includes severe ocular defects, short stature, facial dysmorphism, joint hypermobility/dislocations, scoliosis, thoracic deformity, progressive hearing loss, umbilical hernia, and hypodontia"* [PMID: 42170786].

Phenotype	Type	Onset	Progression	Suggested HPO term
High/severe myopia	Physical/ ophthalmologic	Congenital/ early childhood	Progressive	HP:0011003 (High myopia)
Retinal detachment	Clinical sign	Childhood	Episodic/ progressive	HP:0000541
Congenital glaucoma	Clinical sign	Congenital	Progressive	HP:0001087
Abnormal iris morphology	Physical	Congenital	Stable	HP:0000525
Short stature	Physical	Childhood	Stable/ progressive	HP:0004322
Joint hypermobility	Physical	Congenital	Stable	HP:0001382
Large-joint dislocation (recurrent)	Clinical sign	Congenital/ infancy	Recurrent/ episodic	HP:0002828 (Joint dislocation)
Scoliosis	Physical	Childhood	Progressive	HP:0002650
Thoracic deformity	Physical	Childhood	Progressive	HP:0000765
Sensorineural/mixed hearing loss	Sensory	Childhood	Progressive	HP:0000407 / HP:0000405
Facial dysmorphism	Physical	Congenital	Stable	HP:0001999
Umbilical hernia	Physical	Congenital	Stable	HP:0001537
Hypodontia	Physical	Childhood	Stable	HP:0000668
Congenital heart disease (rarer)	Clinical sign	Congenital	Variable	HP:0001627
Cervical segmentation defects, carpal shortening, lumbar sacralization	Radiological	Congenital	Stable	HP:0000925 / HP:0001180

Severity and frequency. Ocular features are severe and distinguishing (high myopia, retinal detachment, congenital glaucoma) [PMID: 28475863;

].

P 42170786

Hearing loss and scoliosis are described as **progressive** [PMID: 42170786]. Given only ~13 reported patients, precise per-phenotype frequencies cannot be established, but short stature, joint dislocation/hypermobility, and myopia appear in essentially all patients (qualitatively "very frequent").

Quality-of-life impact. Not formally measured with standardized instruments (EQ-5D/SF-36). Inferentially, severe visual impairment (myopia, retinal detachment, glaucoma) and recurrent large-joint dislocations with scoliosis substantially affect mobility, vision, and daily functioning; progressive hearing loss adds communication burden.

4. Genetic/Molecular Information

Causal gene. *GZF1* (GDNF-inducible zinc finger protein 1), 20p11.21. *GZF1* encodes "a novel GDNF-inducible gene (named *GZF1*) with a BTB/POZ (broad complex, tramtrack, and bric-a-brac)/ (poxvirus and zinc finger) domain and 10 tandemly repeated zinc finger motifs" [PMID: 14522971] — i.e., a sequence-specific transcriptional repressor.

Pathogenic variants (all loss-of-function).

Variant (cDNA)	Protein	Type	Zygosity / cohort	PMID
Homozygous truncating variant	Truncation	Nonsense/ frameshift	Homozygous, Saudi families	28475863
c.397_400del	p.Leu133fs	Frameshift	Compound het (Chinese)	33009817
c.1474del	p.Met492fs	Frameshift	Compound het (Chinese)	33009817
c.1440del	p.His481IlefsTer26	Frameshift	Homozygous (2 sisters) / compound het	42170786
c.1451_1452del	p.Cys484fs	Frameshift	Compound het (3rd patient)	42170786

Variant classification. All reported variants are **truncating/frameshift** and classified as pathogenic under ACMG/AMP criteria (predicted loss of function + functional evidence + segregation).

Functional consequences — loss of function. Functional assays showed mutant GZF1 protein is undetectable or cytoplasmically mislocalized with reduced mRNA/protein: *"no HA-conjugated mutant protein was detected by western blotting, which was also confirmed by immunofluorescence staining"* [PMID: 33009817], and the authors concluded *"these results suggested that the two variants could lead to loss of function of GZF1"* [PMID: 33009817]. The mechanism is therefore **loss of function**, not gain of function or dominant negative.

Downstream transcriptional dysregulation. *"Global transcriptional profiling of cells from affected individuals revealed a shared pattern of gene dysregulation and significant enrichment of genes encoding matrix proteins, including P3H2, which hints at a potential disease mechanism"* [PMID: 28475863].

Modifier genes / epigenetics / chromosomal abnormalities. No modifier genes, epigenetic marks, or large-scale chromosomal abnormalities have been reported for this disorder. Diagnosis is at the single-nucleotide/indel level.

Allele frequency. Reported variants are private/ultra-rare and not established in gnomAD at appreciable frequency; the recurrence of c.1440del across families suggests a possible recurrent/founder allele [PMID: 42170786].

5. Environmental Information

- **Environmental factors:** None identified. Monogenic Mendelian disorder.
- **Lifestyle factors:** None implicated in causation. (General myopia-management behavioral advice is not disease-specific.)
- **Infectious agents:** Not applicable.

6. Mechanism / Pathophysiology

Causal chain (initiating lesion → clinical manifestation)

1. **Biallelic loss-of-function variant in *GZF1*** (truncating/frameshift) **results in** absent or mislocalized GZF1 protein [PMID: 33009817].
2. Loss of the GZF1 BTB/POZ + 10-zinc-finger transcriptional repressor **leads to** dysregulated transcription of its target genes in developing eye and limb tissues (where GZF1 is normally expressed) [PMID: 28475863];

P 14522971
3. This dysregulation **results in** significant enrichment of aberrantly expressed **extracellular-matrix/collagen genes, most notably *P3H2* (*LEPREL1*)** [PMID: 28475863] (*demonstrated as transcriptomic enrichment; the precise direction of *P3H2* change and its sufficiency are inferred*).


P 21885030
4. Altered P3H2 (prolyl 3-hydroxylase 2) activity **leads to** collagen under-3-hydroxylation. In the P3h2-null model, "*almost every known site of prolyl 3-hydroxylation in types I and IV collagen from P3h2(n/n) mouse eye tissues was significantly under-hydroxylated*" [PMID: 25645914] (*inferred as the operative lesion in *GZF1RP* by analogy*).
5. Structurally defective scleral (type I) and lens-capsule (type IV) collagen **results in** biomechanically weak, elongating sclera → **axial high myopia**, with associated vitreoretinal degeneration, retinal detachment, cataract and glaucoma [PMID: 25645914];

P 24172257

6. **In parallel (branch):** generalized connective-tissue/ECM collagen dysfunction **leads to** ligamentous/capsular laxity → **joint hypermobility, recurrent large-joint dislocation, scoliosis, thoracic deformity, umbilical hernia** [PMID: 28475863;

P 42170786

7. **In parallel (branch):** GZF1's role in cell proliferation and development (GZF1 supports proliferation; *"knockdown of GZF1 and nucleolin expression markedly impaired cell proliferation"* [PMID: 17674968]) **contributes to short stature (inferred).**



Detail by category

- **Molecular pathways:** GDNF/RET signaling (GZF1 is GDNF-inducible and required downstream of GDNF/RET for renal branching morphogenesis) [PMID: 14522971]; transcriptional repression; collagen prolyl-3-hydroxylation (2-oxoglutarate-dependent dioxygenase pathway) [PMID: 21885030].
- **Cellular processes:** transcriptional regulation, cell proliferation [PMID: 17674968], extracellular-matrix assembly/collagen post-translational modification.
- **Protein dysfunction:** GZF1 — loss of function via truncation/mislocalization [PMID: 33009817]. P3H2 — reduced enzymatic 3-hydroxylation of collagen prolines [PMID: 25645914].

- **Biochemical abnormality:** deficient collagen prolyl 3-hydroxylation (a 2-oxoglutarate-dependent dioxygenase reaction) [PMID: 21885030;

P 25645914

- **Tissue-damage mechanism:** biomechanical failure of collagen-rich sclera and joint capsules/ligaments → axial elongation and laxity.
- **Immune/metabolic involvement:** none reported.

Suggested GO terms: GO:0006355 (regulation of transcription, DNA-templated); GO:0019511 (peptidyl-proline hydroxylation); GO:0030199 (collagen fibril organization); GO:0008283 (cell population proliferation); GO:0030198 (extracellular matrix organization). **Suggested CL terms:** CL:0000057 (fibroblast); scleral/ocular fibroblast (approximate); CL:0000062 (osteoblast); CL:0000138 (chondrocyte).

7. Anatomical Structures Affected

Organ/system level. - **Primary:** Eye (sclera, retina, lens, iris, anterior-chamber angle) — UBERON:0000970; skeletal/musculoskeletal system (large joints, spine, thorax) — UBERON:0002204. - **Secondary:** Ear/auditory system (progressive hearing loss) — UBERON:0001690; teeth (hypodontia); abdominal wall (umbilical hernia); heart (rarer congenital heart disease) — UBERON:0000948. - **Body systems:** ocular/visual, musculoskeletal/connective tissue, auditory, occasionally cardiovascular.

Tissue/cell level. Connective tissue (collagen-rich ECM); scleral/ocular fibroblasts; joint-capsule and ligamentous fibroblasts; growth-plate chondrocytes/osteoblasts (short stature). GZF1 is "*expressed in the eyes and limbs of developing mice*" [PMID: 28475863].

Subcellular level. Nucleus (GZF1 transcriptional repressor; note pathogenic cytoplasmic mislocalization — GO:0005634); nucleolus (nucleolin interaction, GO:0005730) [PMID: 17674968]; endoplasmic reticulum/extracellular region (collagen synthesis and modification, GO:0005783 / GO:0005576).

Localization / laterality. Bilateral ocular and skeletal involvement; joint dislocations affect large joints (hips, knees) and can be recurrent.

Suggested UBERON terms: UBERON:0001801 (sclera), UBERON:0000966 (retina), UBERON:0000965 (lens), UBERON:0001769 (iris), UBERON:0002481 (bone tissue), UBERON:0002217 (joint), UBERON:0001130 (vertebral column).

8. Temporal Development

- **Onset:** Congenital/early. Recurrent large-joint dislocations, congenital glaucoma, and high myopia present from infancy/early childhood [PMID: 42170786].
- **Onset pattern:** Chronic/congenital (structural developmental disorder).
- **Progression:** Myopia is progressive; **hearing loss and scoliosis are explicitly described as progressive** [PMID: 42170786]; retinal detachment can present episodically.
- **Course:** Chronic, lifelong. No remission.
- **Critical periods:** Early childhood is the key window for ophthalmologic surveillance (glaucoma, retinal detachment) and orthopedic management of dislocations and scoliosis.

9. Inheritance and Population

Epidemiology. Ultra-rare: *"Only ten patients from five families have been reported, all of whom carry biallelic variants of GZF1"* [PMID: 42170786], plus three new patients in 2026 (~13 patients from ~6 families total). No prevalence or incidence figures are established. No sex-ratio bias reported (sample too small).

Inheritance. Autosomal recessive, with biallelic *GZF1* variants in all cases [PMID: 42170786]. Original families were multiplex and consanguineous (Saudi), ascertained by autozygome/homozygosity mapping [PMID: 28475863].

Penetrance / expressivity. Apparently high penetrance in biallelic carriers; expressivity is broadly consistent for the core triad with variable additional features (e.g., congenital heart disease is rarer).

Zygosity states observed: homozygous (c.1440del in two sisters) and compound heterozygous (c.1440del + c.1451_1452del; and c.397_400del + c.1474del in Chinese patients) [PMID: 42170786];

].

P 33009817

Founder/recurrent effects. Recurrence of c.1440del across families suggests a possible recurrent/founder allele [PMID: 42170786].

Consanguinity. A significant contributing factor in reported pedigrees [PMID: 28475863].

Population demographics. Reported in Saudi (consanguineous) and Chinese patients. No formal geographic prevalence data. Carrier frequency not established.

10. Diagnostics

Molecular diagnosis is definitive. The gene was identified and diagnoses confirmed by **whole-exome sequencing / autozygosity (homozygosity) mapping** [PMID: 28475863] and by WES in subsequent patients [PMID: 33009817].

Recommended genetic testing approach: - WES/WGS with autozygosity mapping (high yield in consanguineous families). - **Targeted *GZF1* sequencing / connective-tissue or Larsen-syndrome gene panels** (should include *GZF1*, *FLNB*, *CHST3*, *B4GALT7*). - **Functional confirmation** of novel variants: mRNA/protein expression and immunofluorescence in HEK293T cells demonstrated absent mutant protein [PMID: 33009817].

Supportive clinical workup (documented in patients): - **Ophthalmologic examination:** high myopia, retinal detachment, congenital glaucoma, abnormal iris [PMID: 28475863; 42170786].

P 42170786

- **Skeletal radiography:** joint dislocations, scoliosis, thoracic deformity, and newly reported "*cervical segmentation defects, carpal shortening, and lower lumbar sacralization*" [PMID: 42170786]. - **Audiology:** progressive hearing loss. - **Cardiac evaluation:** congenital heart disease reported [PMID: 42170786].

Differential diagnosis. Larsen syndrome and Larsen-like disorders. *GZF1*RP is distinguished by **severe ocular involvement** (high myopia, retinal detachment, congenital glaucoma) not typical of classic *FLNB*-Larsen syndrome: "*GZF1 mutations cause a phenotype of severe myopia and significant articular involvement not previously described in Larsen syndrome*" [PMID: 28475863]. Other differentials: recessive Larsen-like *CHST3/B4GALT7* disorders; other connective-tissue/high-myopia syndromes (e.g., Stickler, Marfan, Ehlers-Danlos, and non-syndromic *LEPREL1* high myopia).

Screening. In known families, **cascade genetic/carrier testing** and prenatal/preimplantation genetic testing are appropriate. No population newborn-screening program exists (ultra-rare).

11. Outcome / Prognosis

Survival/mortality. No excess mortality data reported; the disorder is not described as inherently life-limiting, though rarer congenital heart disease could affect individual prognosis. No survival statistics exist given tiny cohorts.

Morbidity/function. Significant morbidity from **visual impairment** (high myopia, retinal detachment, congenital glaucoma — risk of blindness if untreated), **musculoskeletal disability** (recurrent large-joint dislocations, scoliosis, thoracic deformity), and **progressive hearing loss**. Short stature and facial dysmorphism add to the phenotype.

Disease course. Chronic, lifelong, with progressive ocular, spinal, and auditory features. Recovery potential is limited to what surgical/supportive interventions provide (e.g., retinal-detachment repair, glaucoma control, orthopedic stabilization).

Prognostic factors. Not formally studied; severity of ocular complications and spinal deformity likely drive functional prognosis. No molecular prognostic biomarkers established.

12. Treatment

No disease-modifying or pharmacological therapy exists, and no clinical trials are registered for this ultra-rare disorder. Management is **supportive and multidisciplinary**:

Domain	Intervention	Suggested NCIT concept
Ophthalmologic	Corrective lenses for high myopia; monitoring/surgery for retinal detachment; congenital glaucoma management (medical/surgical IOP control)	NCIT:C15277 (Ophthalmologic procedure)
Orthopedic/surgical	Reduction/stabilization of large-joint dislocations; scoliosis management (bracing/surgery); thoracic deformity care	NCIT:C15329 (Orthopedic surgery)
Audiologic	Hearing aids / audiologic rehabilitation for progressive hearing loss	NCIT:C15194 (Hearing aid)
Dental	Management of hypodontia	—
Surgical	Umbilical hernia repair; cardiac evaluation/repair if congenital heart disease present	NCIT:C51826 (Herniorrhaphy)
Rehabilitative	Physical/occupational therapy for joint stability and mobility	NCIT:C15516 (Physical therapy)
Genetic	Genetic counseling for families	—

Pharmacogenomics, gene/cell/RNA therapy, targeted/immunotherapy: none applicable or available at present.

13. Prevention

- **Primary prevention:** Not possible for a monogenic recessive disorder beyond reproductive counseling. **Genetic counseling** for consanguineous families and known carriers is central.
- **Secondary prevention:** Early ophthalmologic surveillance (glaucoma, retinal detachment) and orthopedic monitoring to prevent complications; audiologic monitoring.
- **Tertiary prevention:** Management of established complications (vision preservation, joint stabilization, scoliosis control, hearing support).
- **Genetic screening: Cascade/carrier testing** in affected families; prenatal diagnosis and preimplantation genetic testing available where the familial variant is known.

- **Counseling:** 25% recurrence risk per pregnancy for carrier couples (autosomal recessive); recurrence and recurrent/founder-allele considerations in specific populations.

14. Other Species / Natural Disease

- **Taxonomy:** *Homo sapiens* (NCBI:txid9606). A relevant mouse model exists for the downstream mechanism (*Mus musculus*, NCBI:txid10090).
- **Orthologous genes:** *Gzfl* (mouse ortholog); *P3h2/Leprel1* (mouse ortholog used in the P3h2-null model).
- **Natural disease in other species:** No naturally occurring GZF1RP counterpart reported in companion animals or wildlife (OMIA entry not established in outline).
- **Comparative biology:** GZF1 developmental function is conserved (expressed in developing mouse eye and limb) [PMID: 28475863]; the collagen prolyl-3-hydroxylation defect is conserved and modeled in mouse [PMID: 25645914].

15. Model Organisms

- **Mouse — P3h2-null (downstream mechanism model):** *"Post-translationally abnormal collagens of prolyl 3-hydroxylase-2 null mice offer a pathobiological mechanism for the high myopia linked to human LEPREL1 mutations"* — the null mice show under-3-hydroxylated types I and IV collagen in eye tissue, recapitulating the ocular collagen defect [PMID: 25645914]. This models the myopia arm of the pathway (not GZF1 itself).
- **Developmental expression model:** GZF1 shown expressed in eyes and limbs of developing mice [PMID: 28475863].
- **Renal organ-culture model:** antisense knockdown of GZF1 impaired ureteric-bud branching morphogenesis, establishing its developmental role downstream of GDNF/RET [PMID: 14522971].
- **In vitro cellular models:** patient-derived cells for transcriptional profiling [PMID: 28475863]; HEK293T over-expression assays for variant functional characterization [PMID: 33009817]; nucleolin-interaction/proliferation assays [PMID: 17674968].
- **Model limitations:** No published *Gzfl*-knockout mouse recapitulating the full human ocular-skeletal syndrome is described; the P3h2-null mouse captures only the myopia/

collagen arm. A dedicated *Gzfl* animal model recapitulating joint laxity and short stature is a key gap.

Key Findings (Expanded)

Finding 1 — Disease identity: GZF1-related phenotype (OMIM #617662), autosomal recessive

Two independent multiplex consanguineous Saudi families with severe recurrent large-joint dislocation and severe myopia each carried homozygous truncating *GZF1* variants, identified via combined autozygome + exome analysis [PMID: 28475863]. As of 2026, ~10 patients from 5 families (plus 3 new patients) all carry biallelic *GZF1* variants, confirming recessive inheritance [PMID: 42170786]. This firmly establishes *GZF1* as the single causal gene and the disorder as autosomal recessive.

Finding 2 — Causal mechanism: biallelic GZF1 LoF dysregulates ECM genes (P3H2)

GZF1 is a BTB/POZ-domain transcriptional repressor with 10 tandem zinc fingers [PMID: 14522971]. Disease variants are truncating/frameshift and functionally loss-of-function, with mutant protein undetectable or mislocalized [PMID: 33009817]. Patient-cell transcriptomics showed enrichment of dysregulated matrix-protein genes including *P3H2* [PMID: 28475863], defining the mechanistic hypothesis.

Finding 3 — Clinical phenotype spectrum distinct from FLNB-Larsen

Recurrent core features (short stature, large-joint dislocation/hypermobility, severe myopia) plus severe ocular defects, facial dysmorphism, scoliosis, thoracic deformity, progressive hearing loss, umbilical hernia, hypodontia, and newly reported skeletal radiological findings define a **recognizable GZF1RP** [PMID: 42170786], distinguished from classic Larsen by severe ocular involvement [PMID: 28475863].

Finding 4 — Mechanistic anchor: P3H2/LEPREL1 collagen under-hydroxylation causes high myopia

Independent recessive kindreds with biallelic *LEPREL1* loss-of-function cause non-syndromic high axial myopia with early cataract, vitreoretinal degeneration, and retinal detachment [PMID: 21885030;

].

P 24172257

P3h2-null mice show under-3-hydroxylated eye collagen, providing a scleral-collagen pathomechanism [PMID: 25645914]. This links GZF1-driven P3H2 dysregulation to the ocular phenotype.

Finding 5 — GZF1 normal biology and model systems

GZF1 is a GDNF/RET-inducible transcriptional repressor required for renal branching morphogenesis [PMID: 14522971], modulated by nucleolin and supporting cell proliferation [PMID: 17674968], and expressed in developing eye and limb — the affected tissues [PMID: 28475863].

Finding 6 — Differential diagnosis and genetic heterogeneity of the Larsen spectrum

Larsen syndrome is genetically heterogeneous: dominant *FLNB* accounts for the majority; recessive forms arise from biallelic *CHST3* and *B4GALT7*; and biallelic *GZF1* defines a further recessive form [PMID: 28475863]. GZF1RP is a distinct, recognizable entity distinguishable from Larsen [PMID: 42170786].

Finding 7 — Epidemiology, inheritance mechanics, natural history

Ultra-rare (~13 patients / ~6 families), fully recessive with biallelic variants, consanguinity-associated, with recurrent alleles (c.1440del) and both homozygous and compound-heterozygous states; congenital/early onset with progressive hearing loss and scoliosis [PMID: 42170786;

;

P 28475863

].

P 33009817

Finding 8 — Diagnostics and management framework

Diagnosis is molecular (WES/autozygosity mapping) with supportive ophthalmologic, skeletal-radiographic, audiologic, and cardiac workup; variant pathogenicity confirmed functionally in HEK293T cells [PMID: 42170786;

P 33009817

P 28475863

Management is supportive; no disease-modifying therapy or registered trials exist.

Mechanistic Model / Interpretation

The coherent narrative is a **transcription-factor → extracellular-matrix → connective-tissue-biomechanics** cascade. Loss of the GZF1 repressor derails the collagen/ECM transcriptional program in developing eye and limb. The best-supported downstream node is **P3H2 (LEPREL1)**: because independent human and mouse evidence shows that P3H2 loss under-hydroxylates ocular type I/IV collagen and causes high axial myopia, the GZF1→P3H2 link plausibly explains the severe ocular phenotype. The same generalized ECM/collagen defect explains the joint laxity, dislocations, scoliosis, thoracic deformity, and hernia arm, while GZF1's proliferative/developmental role plausibly underlies short stature. The strongest evidentiary chain is the ocular arm (transcriptomic enrichment of P3H2 in patients + established P3H2-myopia biology). The skeletal and growth arms are more inferential and warrant a dedicated *Gzf1* animal model.

Mechanistic arm	Strength of evidence	Basis
Ocular (GZF1→P3H2→collagen→myopia)	Strong (partly inferred)	Patient transcriptomics [28475863] + independent P3H2/LEPREL1 human & mouse data [21885030; 24172257; 25645914]
Joint laxity / skeletal	Moderate	ECM dysregulation [28475863]; clinical spectrum [42170786]
Short stature	Weak/ inferred	GZF1 proliferation role [17674968]; developmental expression [28475863]

Evidence Base

PMID	Title (abbrev.)	Role
28475863	<i>GZF1 Mutations Expand the Genetic Heterogeneity of Larsen Syndrome</i>	Landmark: gene discovery, recessive inheritance, P3H2 mechanism, developmental expression, differential Dx
42170786	<i>Expanding the Genetic and Clinical Spectrum of GZF1-Related Phenotype</i>	Defines recognizable GZF1RP, epidemiology, recurrent alleles, new radiological findings
33009817	<i>Novel GZF1 pathogenic variants in two Chinese patients with Larsen syndrome</i>	Functional LoF evidence; compound-het variants; broadens population
14522971	<i>GDNF-inducible gene required for renal branching morphogenesis</i>	GZF1 normal biology: structure, GDNF/RET induction, developmental role
17674968	<i>Nucleolin modulates GZF1 localization/transcription/proliferation</i>	GZF1 in cell proliferation; subcellular regulation
21885030	<i>High myopia caused by a mutation in LEPREL1 (P3H2)</i>	Establishes P3H2/LEPREL1 LoF → high myopia (mechanistic anchor)
24172257	<i>Homozygous LoF LEPREL1 causes high myopia with early cataract</i>	Confirms collagen-modification disruption → high myopia + cataract
25645914	<i>Abnormal collagens of P3h2-null mice / high myopia mechanism</i>	Mouse model: eye-tissue collagen under-3-hydroxylation

No papers in the evidence set challenge the core conclusions; the P3H2 papers strengthen but do not by themselves prove the GZF1→P3H2 causal link in patients (that link rests on transcriptomic enrichment plus mechanistic analogy).

Limitations and Knowledge Gaps

1. **Very small evidence base** (~13 patients / ~6 families) — precludes reliable prevalence, penetrance, sex-ratio, and per-phenotype frequency estimates.
 2. **GZF1→P3H2 causal link is partly inferred** — supported by transcriptomic enrichment in patient cells [PMID: 28475863] plus independent P3H2 biology, but not yet proven by direct rescue/functional epistasis in a GZF1-deficient system.
 3. **No published *Gzf1* animal model** recapitulating the full ocular-skeletal syndrome; the P3h2-null mouse models only the myopia arm.
 4. **Mechanism of short stature is unestablished** (inferred from GZF1's proliferative role).
 5. **No natural-history or QoL instrument data**, no prognostic biomarkers, no therapeutic trials.
 6. **Full target-gene set of GZF1** beyond P3H2 not characterized; other dysregulated ECM genes may contribute.
-

Proposed Follow-up Experiments / Actions

1. **Generate a *Gzf1* loss-of-function mouse** (constitutive and conditional in eye/limb/growth plate) and phenotype for myopia, joint laxity, and skeletal growth to test the full causal chain.
2. **Directly test the GZF1→P3H2 axis:** ChIP/CUT&RUN for GZF1 occupancy at the *P3H2* locus; quantify P3H2 mRNA/protein and collagen 3-hydroxylation in patient-derived fibroblasts/iPSC-derived scleral cells; attempt P3H2 rescue.
3. **Expand the patient registry** via GeneMatcher/consortia to refine phenotype frequencies, penetrance, genotype–phenotype correlations, and possible founder alleles (e.g., c.1440del).
4. **Deep-phenotype the ECM transcriptome/proteome** of patient cells to enumerate additional dysregulated collagen/ECM genes contributing to the skeletal and growth phenotypes.
5. **Establish natural-history and QoL data** (visual, orthopedic, audiology outcomes) to inform surveillance guidelines and care standards.
6. **Assess carrier/allele frequencies** in relevant populations (e.g., Saudi, Chinese) to guide targeted carrier screening.

Report compiled from 8 confirmed findings and 8 primary papers over a 5-iteration investigation. Evidence types: human clinical (case series/genetics), model organism (mouse P3h2-null; mouse expression; renal organ culture), and in vitro (HEK293T functional assays, patient-cell transcriptomics).

Generated by OpenScientist — Scientific Hypothesis Agent for Novel Discovery