

Familial Defective Apolipoprotein B-100 (FDB): A Comprehensive Disease Profile

Disease: Familial Defective Apolipoprotein B-100 (FDB) **MONDO ID:** MONDO:0007751 | **OMIM:** #144010 | **Category:** Mendelian (autosomal dominant / codominant)

Summary

Familial Defective Apolipoprotein B-100 (FDB) is a monogenic autosomal dominant (codominant) form of primary hypercholesterolemia caused by missense variants in the *APOB* gene that impair the ability of apolipoprotein B-100 (apoB-100) to bind the LDL receptor (LDLR). The archetypal and most common cause is the **R3500Q** variant (c.10580G>A; rs5742904; legacy p.Arg3500Gln, mature-protein numbering p.Arg3527Gln), which disrupts a critical intramolecular interaction between **Arg3500** and **Trp4369** that is required for apoB-100 to adopt the conformation necessary for normal LDLR engagement [PMID: 11115503](#). Because defective LDL particles are cleared slowly from the plasma, LDL cholesterol (LDL-C) accumulates, leading to arterial LDL retention, macrophage foam-cell formation, tendon xanthomas, and premature atherosclerotic cardiovascular disease.

The R3500Q variant is a **single European founder mutation** that arose in an ancestor roughly 6,000–7,000 years ago ($\approx 6,750$ years; ~ 270 generations), and virtually all carriers worldwide are identical-by-descent [PMID: 9339363](#). Clinically, FDB is a **milder, frequently underdiagnosed** cousin of LDLR-mediated familial hypercholesterolemia (FH): LDL-C elevation is on average ~ 1.2 mmol/L lower than in LDLR mutation carriers, coronary heart disease occurs less often and later, and standard clinical FH scoring criteria frequently fail to flag FDB carriers [PMID: 15528480](#); [PMID: 25126774](#); [PMID: 18279815](#).

Because the LDL receptor is structurally intact in FDB, the disorder responds well to LDLR-upregulating pharmacotherapy — statins, ezetimibe, and PCSK9 inhibitors — and definitive diagnosis is molecular (single-gene codon-3500 assay or an NGS FH gene panel), enabling reverse-cascade screening of relatives [PMID: 30910740](#); [PMID: 29284604](#). Allelic heterogeneity exists at and near codon 3500 (R3500W, R3531C, R3480W, H3543Y), with **R3500W being an important contributor to FH among East Asians** [PMID: 27919345](#); [PMID: 15135245](#).

1. Disease Information

Overview. FDB is an inherited disorder of LDL catabolism in which a mutant apoB-100 — the sole apolipoprotein of LDL and the ligand recognized by the hepatic LDL receptor — binds the receptor with reduced affinity. The consequence is delayed clearance of LDL, elevated plasma LDL-C and apoB, and an increased lifetime risk of premature coronary artery disease. Clinically it is essentially indistinguishable from heterozygous LDLR-mediated FH without molecular testing; both present with elevated LDL-C, normal triglycerides and HDL-C, tendon xanthomas, and premature atherosclerosis [PMID: 18325181](#).

Key identifiers.

Resource	Identifier
MONDO	MONDO:0007751
OMIM	#144010 (Hypercholesterolemia, due to ligand-defective apoB, or FDB)
Gene	<i>APOB</i> (HGNC:603), chromosome 2p24–23
Classic variant	rs5742904 (R3500Q; c.10580G>A)
Category	Mendelian, autosomal dominant/codominant

Synonyms / alternative names. Familial ligand-defective apolipoprotein B-100; familial defective apoB-100; FDB; ligand-defective apoB; hypercholesterolemia due to ligand-defective apoB. FDB is one molecular subtype within the broader clinical umbrella of *autosomal dominant hypercholesterolemia* (ADH) / familial hypercholesterolemia (FH), alongside *LDLR* and *PCSK9* causes [PMID: 31883481](#).

Data source type. The knowledge in this report derives from **aggregated disease-level resources** — molecular genetics studies, FH registries, population cohorts (e.g., Copenhagen General Population Study), and functional/biochemical studies — rather than individual EHR-derived patient records.

2. Etiology

Primary cause — genetic. FDB is a **monogenic disorder** caused by heterozygous (or, rarely, homozygous/compound heterozygous) missense variants in *APOB* that localize to the LDLR-binding region of apoB-100 near codon 3500. The disease is not infectious or primarily environmental; environmental/lifestyle factors modify severity but do not cause the disorder.

Genetic risk factors. - **Causal variant:** *APOB* R3500Q (c.10580G>A; rs5742904) is the classic and most frequent cause. Additional ligand-defective variants clustering in the same functional region include **R3500W**, **R3531C**, **R3480W**, and **H3543Y** [PMID: 15135245](#); [PMID: 11115503](#). - **Zygosity:** heterozygotes are affected (codominant); homozygous *APOB* cases exist but are rare (e.g., 4 homozygous *APOB* patients among 49 molecularly defined homozygous ADH cases in the Netherlands) [PMID: 24585268](#).

Environmental / modifying risk factors. Background diet (saturated fat, cholesterol intake), obesity, smoking, and coexisting metabolic syndrome raise absolute LDL-C and cardiovascular risk on top of the genetic defect. Phenotypic expression depends jointly on the mutation and the genetic/dietary background of the population studied; ascertainment in lipid clinics overestimates the true population effect size [PMID: 15528480](#).

Protective factors. Because the underlying lesion is impaired receptor-mediated LDL clearance, any factor that upregulates LDLR expression or lowers LDL production is protective — this is the pharmacological basis of statins, ezetimibe, and PCSK9 inhibition. Rare *APOB* missense variants can also lower LDL (e.g., an apoB S2429T variant proposed to have an LDL-lowering effect within an FH family), illustrating that *APOB* variation can push LDL in either direction [PMID: 27578127](#); [PMID: 7585299](#).

Gene–environment interactions. The measured LDL-C increment attributable to a given *APOB* or *LDLR* variant differs by background population and by ascertainment context — general-population carriers and clinically ascertained FH carriers differed by ~1.6 mmol/L (P=0.03) — confirming that the phenotype is a product of genotype × background/lifestyle rather than genotype alone [PMID: 15528480](#).

3. Phenotypes

FDB phenotypes are those of heterozygous FH but generally milder.

Phenotype	Type	HPO suggestion	Onset / severity / frequency
Hypercholesterolemia (elevated LDL-C, elevated apoB)	Laboratory abnormality	HP:0003124 (Hypercholesterolemia); HP:0003563 (Hyperbetalipoproteinemia)	Congenital/lifelong biochemical trait; milder than LDLR-FH; near-universal but LDL-C may fall within normal range in some carriers PMID: 18279815
Tendon xanthomas	Clinical sign / physical manifestation	HP:0001229 / HP:0010874 (Xanthomatosis)	Adult-onset; less frequent than in LDLR-FH; present in a minority PMID: 10559517
Premature coronary artery disease	Clinical sign / complication	HP:0001677 (Coronary artery atherosclerosis); HP:0005110	Adult-onset, later than in LDLR-FH; CHD in ~5.6% of FDB (mean age 52) vs 41.8% of FH (mean age 41) PMID: 25126774
Normal triglycerides and HDL-C	Laboratory (distinguishing feature)	—	Characteristic normal-TG/normal-HDL profile helps distinguish from combined hyperlipidemias PMID: 18325181

Severity and progression. LDL-C elevation is stable/lifelong; atherosclerosis is slowly progressive. Notably, FH/high-LDL-C (including *APOB* R3500Q) increases myocardial-infarction risk but **not** ischemic stroke risk in the general population [PMID: 29593013](#).

Quality of life. No FDB-specific EQ-5D/SF-36 data were identified. QoL impact derives chiefly from cardiovascular events and lifelong pharmacotherapy; because FDB is generally milder and later-onset than LDLR-FH, the average burden is correspondingly lower.

4. Genetic / Molecular Information

Causal gene. *APOB* (apolipoprotein B; HGNC:603; OMIM *107730), chromosome 2p24–23. ApoB-100 (4,536 residues in the mature protein) is the structural scaffold of VLDL/IDL/LDL and the LDLR ligand.

Pathogenic variants.

Variant (legacy)	cDNA / rsID	Type	Notes
R3500Q	c.10580G>A; rs5742904	Missense	Classic FDB variant; most common; European founder PMID: 9339363
R3500W	—	Missense	Significant FH contributor in East Asians; more prevalent than R3500Q in Chinese hyperlipidemics (2.4% vs 0.3%) PMID: 27919345 ; PMID: 10998466
R3531C	—	Missense	Ligand-defective, LDLR-binding region PMID: 15135245
R3480W	—	Missense	Ligand-defective, LDLR-binding region PMID: 15135245
H3543Y	—	Missense	~4× more frequent than R3500Q in a German cardiology cohort (0.47% vs 0.12%) PMID: 15135245

Variant classification (ACMG/AMP). R3500Q is well established as **pathogenic** (functional evidence of defective LDLR binding, cosegregation, founder haplotype). Several additional exon 26/29 variants (e.g., S3476L, S3488G, Y3533C, T3540M, I4350T, G4368D) have been reported as FDB-associated in Dutch cohorts and warrant case-by-case interpretation [PMID: 18325181](#).

Allele frequency. R3500Q carrier frequency is on the order of ~0.1% (≈ 1 in 500–1,000) in Northern/Central European populations, lower in Southern Europe (e.g., rare in general Spain but locally enriched in Galicia due to Celtic ancestry) [PMID: 12208478](#).

Somatic vs germline. Germline; there is no somatic component.

Functional consequence. A **conformational loss-of-ligand-function**: the variant does not abolish the protein but changes the local conformation around the receptor-binding site, reducing LDLR affinity to ~32% of normal [PMID: 10657378](#). This behaves as a dominant/codominant trait because each affected allele produces defective LDL particles.

Modifier genes. Polygenic LDL-raising SNP burden and coexisting variants in other lipid genes (*LDLR*, *PCSK9*, *APOE*, *ANGPTL3*, *APOC3*) modulate expressivity; rare *APOB* variants (e.g., S2429T) may lower LDL and partially offset severity [PMID: 27578127](#); [PMID: 31883481](#).

Epigenetic / chromosomal. No recurrent epigenetic mechanism or chromosomal abnormality is implicated in FDB; it is a point-mutation disorder.

5. Environmental Information

- **Environmental factors:** No specific toxin, radiation, or occupational exposure causes FDB.
- **Lifestyle factors:** Dietary saturated fat/cholesterol, obesity, physical inactivity, and smoking increase absolute LDL-C and atherosclerotic risk on top of the genetic defect and are the primary modifiable levers.
- **Infectious agents:** None. FDB is a purely genetic, non-communicable disorder.


6. Mechanism / Pathophysiology

Ordered causal chain

1. A germline missense variant in *APOB* (most commonly **R3500Q**) is inherited → **leads to** an amino-acid substitution in apoB-100 at/near the LDLR-binding region.
2. The substitution disrupts the intramolecular **Arg3500–Trp4369** interaction (and requires the apoB-100 C-terminus) → **results in** an altered local conformation of apoB-100 around

the receptor-binding site (demonstrated by higher affinity for the conformation-sensitive MB47 antibody) [PMID: 11115503](#).

3. The altered conformation → **reduces** binding of LDL particles to the hepatic LDL receptor (to ~32% of normal affinity) [PMID: 10657378](#).
4. Reduced receptor binding → **decreases** receptor-mediated hepatic uptake/catabolism of LDL → **prolongs** LDL residence time in plasma.
5. Delayed clearance → **raises** plasma LDL-C and apoB concentrations (the biochemical phenotype).
6. Elevated (and modified/oxidized) LDL → **accumulates and is retained** in the arterial intima.
7. Retained LDL → **is internalized by macrophages** → **drives** foam-cell formation and an immune-inflammatory response [PMID: 33959626](#).
8. Chronic foam-cell/inflammatory burden → **produces** atherosclerotic plaques → **manifests** clinically as tendon xanthomas and premature coronary artery disease [PMID: 18325181](#).



Upstream vs downstream. The conformational defect and reduced receptor binding are the **upstream initiating lesions**; hypercholesterolemia is the intermediate biochemical readout; foam-cell formation and atherosclerosis are **downstream** consequences shared with all LDL-driven hypercholesterolemias.

Molecular pathways / cellular processes. Receptor-mediated endocytosis of LDL (KEGG hsa04979, "Cholesterol metabolism"); reverse cholesterol transport (ABCA1-mediated efflux); macrophage lipid handling and NF-κB-driven vascular inflammation are engaged downstream [PMID: 33568202](#); [PMID: 36838686](#).

Protein dysfunction. Conformational loss-of-ligand-function of apoB-100 (not misfolding/aggregation, not loss of expression). UniProt P04114 (APOB_HUMAN).

Metabolic changes. Isolated elevation of LDL-C and apoB with characteristically normal triglycerides and HDL-C — distinguishing FDB from combined hyperlipidemias.

Suggested ontology terms. GO:0006629 (lipid metabolic process); GO:0034383 (low-density lipoprotein particle clearance); GO:0006898 (receptor-mediated endocytosis); GO:0034364 (low-density lipoprotein particle); CHEBI:39025 (low-density lipoprotein). Cell types: CL:0000182 (hepatocyte), CL:0000235 (macrophage/foam cell), CL:0000891 (aortic/vascular smooth muscle context).

7. Anatomical Structures Affected

- **Primary organ:** Liver (hepatocytes) — the principal site of LDL-receptor-mediated clearance whose function is bypassed by the defective ligand. UBERON:0002107 (liver); CL:0000182 (hepatocyte).
- **Secondary/target organs:** Arterial wall, especially coronary arteries (and femoral/peripheral arteries). UBERON:0001981 (blood vessel); UBERON:0001621 (coronary artery). Femoral intima-media thickness is increased in FH patients including *APOB* defects, in relation to mutational class [PMID: 18096825](#).
- **Body systems:** Cardiovascular (atherosclerosis, CAD) and integumentary/musculoskeletal (tendon xanthomas). Blood/plasma lipoprotein compartment is the biochemical stage.
- **Tissue/cell level:** Vascular intima (connective tissue), arterial macrophages/foam cells (CL:0000235), and hepatocytes.
- **Subcellular:** Plasma-membrane LDL receptor and clathrin-coated pits (GO:0005905); endosomes/lysosomes for LDL degradation (GO:0005764). The extracellular lipoprotein particle (GO:0034364) is the affected entity.
- **Lateralization:** Systemic/bilateral (xanthomas often symmetric); vascular disease is diffuse rather than lateralized.

8. Temporal Development

- **Onset:** The biochemical trait (elevated LDL-C/apoB) is **congenital and lifelong**. Clinical complications are **adult-onset**, typically later than in LDLR-FH.
 - **Onset pattern:** Chronic/insidious.
 - **Progression:** Slowly progressive atherosclerosis. FDB CHD occurs at a mean age of ~52 years versus ~41 years in LDLR-FH, and far less frequently (5.6% vs 41.8%) [PMID: 25126774](#).
 - **Course:** Stable biochemical elevation with progressive vascular risk; chronic lifelong disease.
 - **Critical window:** Cumulative LDL exposure drives risk, so the critical period for intervention is **early and sustained LDL lowering** from young adulthood (or childhood in higher-risk cases), before substantial plaque burden accrues.
-

9. Inheritance and Population

Inheritance. Autosomal **dominant / codominant**; heterozygotes are affected. Homozygous/compound-heterozygous *APOB* cases are rare and more severe [PMID: 24585268](#).

Penetrance / expressivity. Biochemical penetrance is high but **expressivity is variable** — some carriers have LDL-C within the normal range, and clinical FH scores (Dutch Lipid Clinic Network, Simon Broome) have low sensitivity for *APOB* variants (DLCN sensitivity 13.8%; SBDC 22.5%) [PMID: 30270060](#); [PMID: 18279815](#).

Founder effect. R3500Q is a single European founder mutation ~6,000–7,000 years old (~270 generations); all probands are identical-by-descent, carried on the rare *APOB* haplotype 194 (XbaI–/MspI+/EcoRI–) [PMID: 9339363](#); [PMID: 12208478](#).

Epidemiology. Estimated carrier frequency ~0.1% (~1 in 500–1,000) in Northern/Central Europe; combined LDLR+*APOB* FH prevalence ~1 in 125–135 in Switzerland, with FDB a minority subtype [PMID: 30270060](#). FDB is a minor cause of clinically diagnosed FH in Southern Europe (e.g., 1.4% of Spanish clinical-FH probands; population prevalence ~2.8×10⁻⁵), but locally enriched in Galicia [PMID: 12208478](#); [PMID: 16596945](#).

Population distribution of variants. R3500Q predominates in European-descent populations; **R3500W is the major ligand-defective *APOB* variant in East Asians** (identified in 8 Taiwanese FH probands; heterozygote prevalence 2.4% vs 0.3% for R3500Q in Chinese hyperlipidemics) [PMID: 20538126](#); [PMID: 10998466](#).

Sex ratio / consanguinity. No strong sex bias (autosomal dominant); consanguinity is not required for the dominant heterozygous phenotype but underlies rare homozygous cases.

10. Diagnostics

Laboratory. Elevated LDL-C and apoB with normal triglycerides and HDL-C (LOINC: 13457-7 LDL-C; 1884-6 apoB) [PMID: 18325181](#). A functional flow-cytometric LDL-ligand assay can detect heterozygous FDB by demonstrating reduced LDL binding [PMID: 10657378](#).

Genetic testing (definitive). FDB **cannot be distinguished clinically from LDLR-FH** and requires molecular confirmation [PMID: 18325181](#). Approaches: - Targeted single-variant/codon-3500 assays (PCR-RFLP, high-resolution melting) for R3500Q/H3543Y [PMID: 18325181](#). - NGS FH gene panels covering *LDLR*, *APOB*, *PCSK9* ($\pm$ *LDLRAP1*, *ABCG5/8*) [PMID: 41412630](#); [PMID: 41760042](#). - WES/WGS are useful when panels are negative.

Clinical criteria & differential diagnosis. DLCN and Simon Broome criteria are standard for FH but **under-detect FDB** [PMID: 30270060](#). Differential diagnosis includes LDLR-mediated FH, *PCSK9* gain-of-function FH, polygenic hypercholesterolemia, familial combined hyperlipidemia [PMID: 34906840](#), and rare recessive/other catabolic defects (a clinical HoFH phenotype has been reported with intact apoB/LDLR, implying additional catabolic genes) [PMID: 9626156](#).

Screening. Cascade (reverse-cascade) screening of first-degree relatives after an index molecular diagnosis is the key strategy; universal newborn/pediatric screening for FH is supported by healthcare professionals but not yet routine [PMID: 41251905](#); [PMID: 41608359](#).

11. Outcome / Prognosis

- **Survival/mortality:** No FDB-specific survival tables identified. Prognosis is driven by cardiovascular event risk, which is **lower and later** than in LDLR-FH.

- **Morbidity:** Premature CAD is the dominant morbidity; CHD prevalence ~5.6% in FDB vs 41.8% in LDLR-FH [PMID: 25126774](#). FH/high LDL-C raises myocardial infarction risk but not ischemic stroke risk [PMID: 29593013](#).
- **Prognostic factors:** Mutational class (null LDLR alleles > receptor-defective > *APOB* defects for atherosclerosis severity and LDL-C), absolute/cumulative LDL-C, age, sex, smoking, and blood pressure [PMID: 18096825](#); [PMID: 28475941](#).
- **Recovery potential:** With early sustained LDL lowering, cardiovascular risk is substantially reducible; the intact LDL receptor predicts good pharmacologic response.

12. Treatment

FDB is managed as **heterozygous FH**, and — critically — because the LDL receptor itself is intact, LDLR-upregulating therapies are highly effective.

Therapy (NCIT suggestion)	Class / mechanism	Evidence in FDB context
Statins (e.g., rosuvastatin, atorvastatin) — NCIT:C1518	HMG-CoA reductase inhibitor → ↑ hepatic LDLR expression	Mevastatin increases LDLR expression in vitro; statins are first-line for FH including <i>APOB</i> PMID: 29284604 ; PMID: 30910740
Ezetimibe — NCIT:C61708	NPC1L1 inhibitor → ↓ intestinal cholesterol absorption	Standard add-on in heterozygous FH PMID: 30910740
PCSK9 inhibitors (evolocumab, alirocumab) — NCIT:C121787	Monoclonal antibody → prevents LDLR degradation → ↑ cell-surface LDLR	PCSK9-neutralizing antibody restores LDLR expression; effective because receptor is intact PMID: 29284604

Rationale. "There is today a large availability of drugs (i.e., statins, ezetimibe and PCSK9 inhibitors) allowing theoretically the normalization of plasma LDL cholesterol levels in this population" [PMID: 30910740](#). Because FDB LDL clearance depends on functional LDLR, upregulating the receptor compensates for the ligand defect — mechanistically supported by the observation that "Mevastatin increased, whereas rPCSK9 reduced LDLR expression. The PCSK9-neutralizing antibody restored LDLR expression" [PMID: 29284604](#).

Experimental / advanced. Gene- and cell-based therapies (LDLR gene delivery, CRISPR/Cas9 editing) are in development primarily for homozygous FH; RNA therapies and MTP/ANGPTL3-directed agents extend options for severe/refractory disease [PMID: 36572377](#); [PMID: 30910740](#).

Personalized medicine. Molecular subtyping (APOB vs LDLR-null vs LDLR-defective) informs prognosis and intensity of therapy; genotype-guided cascade screening extends benefit to relatives [PMID: 41412630](#).

13. Prevention

- **Primary prevention:** Lifelong healthy diet, avoidance of smoking, weight and blood-pressure control to minimize cumulative atherosclerotic risk.
- **Secondary prevention (early detection):** Reverse-cascade genetic and lipid screening of relatives after an index diagnosis; consideration of pediatric/newborn FH screening [PMID: 41251905](#); [PMID: 41608359](#).
- **Tertiary prevention:** Aggressive LDL lowering (statin ± ezetimibe ± PCSK9 inhibitor) to prevent CAD progression and events.
- **Genetic counseling:** 50% transmission risk per offspring (autosomal dominant); counseling supports family planning and cascade testing [PMID: 41412630](#).

14. Other Species / Natural Disease

- **Taxonomy/orthologs:** *APOB* is conserved across mammals (human *APOB*, NCBI Gene 338; mouse *Apob*, NCBI Gene 238). No specific naturally occurring animal disease attributable to an APOB R3500Q-equivalent variant was identified in this investigation.
- **Comparative biology:** The apoB-100/LDLR clearance axis is evolutionarily conserved, and the mechanism has been functionally validated in transgenic mice expressing human R3500Q and W4369Y apoB, which reproduce the defective LDLR binding [PMID: 11115503](#).
- **Transmission/zoonosis:** Not applicable — non-communicable genetic disorder.

15. Model Organisms

- **Transgenic mice:** Mice expressing human apoB-100 carrying **R3500Q** (and the complementary **W4369Y**) were pivotal in defining the mechanism — LDL from both showed identically defective LDLR binding and increased MB47 antibody affinity, demonstrating the Arg3500–Trp4369 conformational interaction [PMID: 11115503](#).
 - **In vitro / cellular models:** Cultured cells (e.g., fibroblasts, HepG2-type systems) are used to quantify LDLR expression and LDL binding under statin/PCSK9 modulation, and to test LDLR-upregulating therapeutics [PMID: 29284604](#). Macrophage foam-cell models (e.g., RAW264.7, ApoE-deficient mouse aortas) recapitulate the downstream atherosclerotic steps [PMID: 33568202](#); [PMID: 36838686](#).
 - **Phenotype recapitulation:** Transgenic mice faithfully reproduce the **ligand-binding defect** (the core lesion). **Limitation:** murine lipoprotein metabolism differs from human, so absolute LDL-C phenotype and atherosclerosis burden require sensitized backgrounds (e.g., ApoE-deficient) and do not fully mirror human clinical disease.
 - **Resources:** MGI (mouse *ApoB*), Cellosaurus/ATCC (cell lines).
-

Mechanistic Model / Interpretation

FDB is best understood as a **ligand-side mirror image of LDLR-mediated FH**: in classic FH the receptor is broken, whereas in FDB the ligand (apoB-100) is broken. Both converge on the same downstream pathology — impaired LDL clearance → hypercholesterolemia → arterial LDL retention → foam cells → atherosclerosis — but they differ quantitatively. The single conformational contact point (Arg3500–Trp4369) is the linchpin: substituting Arg3500 (a specific arginine, not merely any positive charge) collapses the receptor-competent conformation, cutting binding to ~32% of normal. Because only the ligand on the mutant allele is defective and the receptor pool is fully functional, (a) heterozygotes retain substantial clearance capacity (hence a milder phenotype and frequent underdiagnosis), and (b) upregulating the intact receptor pharmacologically (statins, PCSK9 inhibitors) efficiently rescues clearance. This mechanistic logic explains, in one coherent chain, the biochemistry, the milder-than-LDLR-FH clinical course, and the excellent drug response.

Evidence Base

PMID	Contribution	Role
11115503	R3500Q changes apoB-100 conformation near the receptor site; Arg3500–Trp4369 interaction defined via transgenic mice	Core mechanism (F001)
9339363	R3500Q single European founder ~6,000–7,000 yrs ago; identical-by-descent	Founder genetics (F002)
15528480	LDLR carriers ~1.2 mmol/L higher LDL-C than APOB R3500Q; ascertainment effects	Milder phenotype (F003)
25126774	CHD 5.6% (age 52) in FDB vs 41.8% (age 41) in FH	Lower/late CVD burden (F003)
18279815	No FDB subject met confirmed FH diagnosis under Med-Ped criteria	Underdiagnosis (F003)
30910740	Statins/ezetimibe/PCSK9i can normalize LDL-C in heterozygous FH	Treatment rationale (F004)
29284604	Statin ↑ and PCSK9-antibody restores LDLR expression	Drug mechanism (F004)
27919345	R3500W significant FH contributor in East Asians	Allelic heterogeneity (F005)
15135245	R3500W/R3531C/R3480W ligand-defective; H3543Y frequent	Variant spectrum (F005)
10998466	R3500W (2.4%) > R3500Q (0.3%) in Chinese hyperlipidemics	Variant spectrum (F005)
10657378	FDB LDL binds LDLR at 32% of normal affinity	Quantifies initiating step (F006)
33959626	Cholesterol-rich LDL internalized by macrophages → foam cells + inflammation	Downstream chain (F006)
18325181	FDB clinically indistinguishable from FH; HRM detection; cascade screening	Diagnosis/phenotype
30270060	DLCN/SBDC scores under-detect APOB FH	Diagnostic limitation
12208478		

PMID	Contribution	Role
	R3500Q rare in Spain but enriched in Galicia; single haplotype	Epidemiology/ founder
18096825	Femoral atherosclerosis by mutational class (APOB milder than null LDLR)	Prognosis
29593013	FH/high LDL-C raises MI but not ischemic stroke risk	Outcome specificity

Limitations and Knowledge Gaps

1. **No primary omics data were analyzed** in this investigation; conclusions rest on literature synthesis of genetics, biochemistry, and clinical registries.
2. **FDB-specific epidemiology is imprecise** — carrier-frequency estimates vary widely by population and ascertainment; robust general-population prevalence figures for R3500Q specifically remain limited.
3. **Nomenclature ambiguity** — legacy (R3500Q) vs mature-protein (R3527Q) vs full-length numbering can cause confusion; reports should standardize to HGVS.
4. **Variant classification** for non-R3500Q *APOB* variants (e.g., the Dutch exon-26/29 variants) is incomplete; several remain of uncertain functional impact.
5. **No dedicated QoL, survival, or mortality datasets** specific to FDB were located.
6. **Model-organism fidelity** — transgenic mice capture the ligand defect but not the full human atherosclerotic phenotype without sensitizing backgrounds.
7. **Non-European populations** are under-studied; R3500W in East Asians is documented but broad allelic surveys are sparse.

Proposed Follow-up Experiments / Actions

1. **Query gnomAD directly** for population-stratified allele frequencies of rs5742904 (R3500Q) and R3500W to quantify carrier frequencies with confidence intervals.
2. **Pull ClinVar/ClinGen classifications** for all *APOB* codon-3480 to 3543 variants to build a definitive pathogenicity table with ACMG criteria.

3. **Systematic outcomes meta-analysis** comparing cardiovascular event rates and treated LDL-C in molecularly confirmed FDB vs LDLR-FH cohorts (e.g., SAFEHEART, DIAMOND-FH data).
 4. **Structural analysis** (AlphaFold/PDB) of the apoB-100 receptor-binding region to visualize the Arg3500–Trp4369 contact and model each ligand-defective variant's disruption.
 5. **Real-world PCSK9-inhibitor response study** in genotyped FDB patients to confirm the predicted superior response given intact LDLR.
 6. **Cascade-screening yield analysis** to quantify how many relatives per index FDB case are newly diagnosed and treated, informing cost-effectiveness of molecular-first screening.
-

Report compiled from 40 reviewed papers and 6 confirmed findings across 5 investigation iterations. Evidence types span human clinical/registry studies, transgenic mouse and in-vitro functional work, and population genetics. All mechanistic and clinical claims are cited to primary literature (PMIDs above).
