Hypotonia, Ataxia, and Delayed Development Syndrome (HADDS): A Comprehensive Disease Characterization

Disease: Hypotonia, Ataxia, and Delayed Development Syndrome (HADDS) **Causal gene:** *EBF3* (Early B-cell Factor 3; HGNC:19087; gene MIM 607407) **Identifiers:** MONDO:0015021 · OMIM:617330 · Orphanet:658843 **Category:** Mendelian, autosomal dominant neurodevelopmental disorder

Summary

Hypotonia, Ataxia, and Delayed Development Syndrome (HADDS) is a rare, autosomal-dominant neurodevelopmental disorder caused by **heterozygous loss-of-function of *EBF3***, a highly conserved Collier/Olf/EBF (COE) family transcription factor located at chromosome 10q26.3. The disorder was defined simultaneously by three independent whole-exome sequencing studies in 2017, which identified *de novo* missense, nonsense, frameshift, splice-site, and in-frame duplication variants, as well as whole-gene deletions, in individuals with a recognizable syndrome of congenital hypotonia, ataxia, global developmental delay, expressive speech impairment, facial dysmorphism, and genitourinary dysfunction. The unifying molecular mechanism is **haploinsufficiency**: most pathogenic variants reduce the dosage of functional EBF3 protein, and the fact that complete EBF3 deletion (via a 10q26.3 microdeletion) reproduces the phenotype confirms that one functional copy of the gene is insufficient for normal neurodevelopment.

Mechanistically, EBF3 is a dosage-sensitive transcription factor (gnomAD pLI ≈ 1.00; LOEUF upper bound 0.346) that binds DNA through a zinc-knuckle/zinc-finger (ZNF) motif and activates transcription of target genes. Pathogenic missense variants cluster in the DNA-binding domain (recurrent residues Arg163 and Arg209), where they impair DNA binding, cause cytoplasmic mislocalization, and abolish transcriptional activation of reporter genes such as *CDKN1A*. During brain development, EBF3 acts downstream of the homeobox selector gene *ARX* and within a *Prdm8–EBF3–NeuroD1* axis to control neuronal migration and the

multipolar-to-bipolar transition of migrating neurons; disruption of this program is the inferred cause of the cerebellar malformation (vermis hypoplasia with a characteristic radial "dandelion-sign" foliation) and the cortical/white-matter abnormalities seen on imaging.

Clinically, HADDS is lifelong and largely static, with **highly variable expressivity** — developmental delay is the only near-universal feature, while ataxia, abnormal muscle tone, and MRI abnormalities each occur in roughly a third to two-thirds of patients, and rare individuals have normal IQ. Severity correlates with the location of missense variants: those perturbing the zinc-finger domain define a high-risk subtype. There is **no disease-specific or disease-modifying therapy**; management is entirely supportive and multidisciplinary (physical, occupational, and speech therapy; aggressive surveillance and treatment of neurogenic bladder to protect renal function; ophthalmologic management of strabismus; developmental and educational support). The disorder is ultra-rare, known only from small case series and one observational registry.

Key Findings

Finding 1 — HADDS is caused by heterozygous loss-of-function of *EBF3* via haploinsufficiency

Three independent 2017 exome studies established that *de novo* variants in *EBF3* — spanning missense, nonsense, frameshift, splice-site, and a 9-bp duplication — cause a syndromic neurodevelopmental disorder. The variants are predicted to trigger nonsense-mediated mRNA decay (NMD) and reduce functional protein dosage. Critically, complete deletion of *EBF3* within a 10q26.3 microdeletion reproduces the HADDS phenotype, which is the strongest line of evidence that the mechanism is **haploinsufficiency** rather than a dominant-negative or gain-of-function effect.

As stated in the clinical-spectrum review, *"EBF3 pathogenic variants have been predicted to result in nonsense-mediated mRNA decay and haploinsufficiency"* and *"total deletion of EBF3 associated with a 10q26.3 microdeletion also causes HADDS symptoms, supporting the concept that HADDS results from haploinsufficiency of EBF3"* (PMID: 34050706). The founding report described *"de novo variants in EBF3"* as the cause (PMID: 28017372).

Evidence type: human clinical (exome sequencing, cohort studies).

Finding 2 — EBF3 missense variants in the DNA-binding zinc-knuckle impair transcriptional activation

In vitro functional assays in HEK293T cells demonstrated that EBF3 mutant proteins have significantly reduced or absent ability to activate transcription of the reporter gene *CDKN1A*. Most mutants mislocalized to the cytoplasm and were less tightly associated with chromatin, and genome-wide RNA-seq and ChIP-seq showed reduced DNA binding and gene-regulatory activity. The recurrent *de novo* variants affect a single conserved residue in a zinc finger motif that is crucial for DNA binding, and these variants are deleterious in a *Drosophila* model.

Direct quotes: *"By transactivation assays, all EBF3 mutants showed significantly reduced or no ability to activate transcription of the reporter gene CDKN1A"* and *"mutant EBF3 had reduced genome-wide DNA binding and gene-regulatory activity"* (PMID: 28017373). The founding study noted that *"the de novo variants affect a single conserved residue in a zinc finger motif crucial for DNA binding and are deleterious in a fly model"* (PMID: 28017372).

Evidence type: in vitro (cell-based transactivation, RNA-seq, ChIP-seq) and in vivo model organism (*Drosophila*).

Finding 3 — Neurogenic bladder / urologic dysfunction is a frequent, early feature

Genitourinary involvement is a distinctive and clinically important feature of HADDS. In a 52-patient aggregate series (33 female : 19 male), 67% had a urologic diagnosis, with a high prevalence of urinary tract infection, vesicoureteral reflux (VUR), and neurogenic bladder, and females disproportionately affected. In a separate 8-patient series, neurogenic bladder was diagnosed in infancy (median 6.5 months) and required cystostomy in all but one case. The proposed mechanism is impaired relaxation of the urinary sphincter producing detrusor-sphincter dyssynergia.

As stated: *"We hypothesize a high rate of bladder dysfunction secondary to the EBF3 variant's impact on relaxation of the urinary sphincter leading to detrusor sphincter dyssynergia (DSD)"* (PMID: 37718233), and *"neurogenic bladder was diagnosed in infancy (the median 6.5 months), was more frequent than previously reported, and required cystostomy in all but one case"* (PMID: 34050706).

Suggested HPO terms: HP:0000011 (Neurogenic bladder), HP:0000076 (Vesicoureteral reflux), HP:0000010 (Recurrent urinary tract infections).

Evidence type: human clinical (retrospective cohort and case series).

Finding 4 — Cerebellar vermis hypoplasia with a radial "dandelion-sign" foliation is a characteristic neuroimaging feature

In a 6-patient neuroimaging series (2 female : 4 male, ages 2–12), developmental milestones were invariably delayed, all patients showed ataxic signs confirmed by SARA (Scale for the Assessment and Rating of Ataxia) scores — often with hypotonia — and brain MRI in all children revealed a cerebellar malformation with vermis hypoplasia and a peculiar foliation anomaly characterized by radial disposition of the cerebellar folia (the **"dandelion sign"**). Neurophysiologic examinations (EEG, evoked potentials) were unremarkable.

Direct quotes: *"Brain MRI revealed in all children a cerebellar malformation with vermis hypoplasia and a peculiar foliation anomaly characterized by a radial disposition of cerebellar folia (dandelion sign)"* and *"All patients showed ataxic signs, as confirmed by SARA scores, often associated with hypotonia"* (PMID: 37090941).

Suggested HPO terms: HP:0001320 (Cerebellar vermis hypoplasia), HP:0001251 (Ataxia), HP:0001252 (Hypotonia), HP:0002194 (Delayed gross motor development).

Evidence type: human clinical (neuroimaging case series).

Finding 5 — Autosomal dominant, usually *de novo*, with documented parental mosaicism affecting recurrence risk

Across the 2017 cohorts, *EBF3* variants arose *de novo* in the majority of cases (e.g., 8 of 10 individuals in the Harms et al. series). Importantly, the missense variant c.625C>T (p.Arg209Trp) was inherited by two affected siblings from their clinically healthy, mosaic mother — direct evidence of germline/somatic mosaicism with a nonzero recurrence risk for apparently unaffected parents. Recurrent substitutions cluster at DNA-binding residues (e.g., Arg163, Arg209); an independent case confirmed a *de novo* c.625G>A (p.Arg209Trp) variant impairing DNA binding, localization, and transactivation.

Direct quotes: *"the missense variant c.625C>T (p.Arg209Trp) was inherited by two affected siblings from their healthy mother, who is mosaic"* (PMID: 28017373); *"a de novo heterozygous c.625G>A (p.Arg209Trp) variant of the EBF3 gene"* (PMID: 36317217).

Evidence type: human clinical (segregation and mosaicism analysis).

Finding 6 — Missense variants disrupting the zinc-finger (ZNF) domain define a high-severity genotype–phenotype subtype

An integrated phenotypic/genotypic study of 41 individuals, combined with a literature meta-analysis (total n = 83 with EBF3-related neurodevelopmental disorder) using quantitative severity scales, found that patient symptom severity correlates with missense variants perturbing the zinc-finger (ZNF) domain, which stabilizes the EBF3–DNA interaction. In vivo *Drosophila* UAS-GAL4 rescue experiments and in vitro luciferase assays showed that ZNF-associated variants failed to restore viability and impaired transcriptional activation, whereas the recurrent DNA-binding-domain variant p.Arg209Trp partially rescued fly viability and preserved some function — providing a mechanistic basis for the severity gradient.

Direct quotes: *"patient symptom severity correlates with EBF3 missense variants perturbing the ZNF, which is a key protein domain required for stabilizing the interaction between EBF3 and the target DNA sequence"* and *"ZNF-associated variants failed to restore viability in the fruit fly and impaired transcriptional activation"* (PMID: 35340043).

Evidence type: human clinical (genotype–phenotype correlation) plus in vivo/in vitro functional validation.

Finding 7 — *De novo* noncoding regulatory (enhancer hs737) variants also implicate *EBF3* in autism

Whole-genome sequencing of 2,671 autism families (discovery cohort 516, replication 2,155) identified an excess of *de novo* variants in the brain-active enhancer **hs737** (discovery p = 0.0172; significant replication), which targets *EBF3* — a gene that is genome-wide significant in neurodevelopmental disorders. This demonstrates that *EBF3* dysregulation can arise through both coding and noncoding *de novo* variation, broadening the mutational spectrum relevant to the EBF3 phenotypic continuum.

Direct quotes: *"we identify DNVs in the hs737 enhancer in individuals with autism"* and *"we find hs737 targets the gene EBF3 that is genome-wide significant in NDDs"* (PMID: 34256850).

Evidence type: human clinical (whole-genome sequencing, enhancer burden analysis).

Finding 8 — *EBF3* is a highly constrained, dosage-sensitive gene (gnomAD), corroborating haploinsufficiency

gnomAD v2.1.1 constraint metrics for *EBF3* (ENSG00000108001; chr10:129,835,233–129,973,053, minus strand) strongly support intolerance to loss of function:

Metric	Value	Interpretation
pLI	1.00 (0.99999)	Extreme intolerance to LoF
Observed/expected LoF	0.233 (18 obs vs 77.1 exp)	~77% fewer LoF variants than expected
LOEUF (upper bound)	0.346	Below the 0.35 haploinsufficiency threshold
LoF Z	5.71	Strong constraint
Missense Z	4.39 (oe_mis = 0.65)	Missense constraint
Synonymous Z	−0.21	Neutral (as expected)

Identifiers: HGNC:19087; gene MIM 607407; aliases COE3, O/E-2, HADDS. The genic constraint provides an orthogonal, population-genetics line of evidence that a single functional copy of *EBF3* is insufficient.

Evidence type: computational / population genetics (gnomAD).

Finding 9 — ClinVar catalogs hundreds of *EBF3* variants, including ~230 pathogenic/likely pathogenic

A ClinVar query (2026-11) returned 528 *EBF3* variant records total, of which 162 are classified Pathogenic and 68 Likely pathogenic (~230 P/LP); 414 records are single-nucleotide variants. Pathogenic variant classes span missense (clustering in the DNA-binding/zinc-finger domain), nonsense, frameshift, and splice-site changes, plus whole-gene deletions (10q26.3). Recurrent residues include Arg163 and Arg209 in the DNA-binding domain.

Variant class	Representative examples	Functional consequence
Missense (DNA-binding/ZNF)	p.Arg163Gly, p.Arg163Leu, p.Arg209Trp, p.Arg152Cys, p.Phe211Leu	Loss of DNA binding, mislocalization, loss of transactivation
Nonsense	p.Arg470 *(C-terminal), p.Tyr321*	Truncation / NMD → haploinsufficiency
Frameshift	Various	NMD → haploinsufficiency
Splice-site	Various	Aberrant splicing → LoF
Whole-gene deletion	10q26.3 microdeletion	Null allele (phenocopy)

Evidence type: curated variant database (ClinVar).

Finding 10 — Highly variable expressivity, including normal-IQ cases; developmental delay is the only near-universal feature

In a 12-patient Chinese cohort (5 missense + 7 CNVs), developmental delay/intellectual disability was present in 100%, but ataxia in only 33% (6/9), abnormal muscle tone in 55% (6/11), and aberrant brain MRI in 64% (7/11). Novel missense variants localized to the DNA-binding domain (R152C and F211L altered subcellular localization; R163G shifted distribution to a granular pattern). Separately, two unrelated patients with *de novo* variants (p.Arg152Leu; p.Tyr321*) had IQ in the normal range, demonstrating that EBF3-related neurodevelopmental disorder can occur without** intellectual disability; diffusion tensor imaging showed asymmetric longitudinal fasciculus (white-matter long-tract) abnormalities.

Direct quotes: *"All of these patients exhibited developmental delay/intellectual disability. Ataxia was observed in 33% (6/9) of the patients, and abnormal muscle tone was observed in 55% (6/11) of the patients. Aberrant MRI reports were noted in 64% (7/11) of the patients"* (PMID: 36937983); *"Our results demonstrate that EBF3 mutations can underlie neurodevelopmental disorders without intellectual disability"* (PMID: 34177436).

Phenotype	Frequency (12-pt cohort)	Suggested HPO term
Developmental delay / intellectual disability	100%	HP:0001263 / HP:0001249
Abnormal muscle tone (hypotonia)	55% (6/11)	HP:0001252
Aberrant brain MRI	64% (7/11)	HP:0002060
Ataxia	33% (6/9)	HP:0001251

Evidence type: human clinical (cohort with quantitative frequencies).

Finding 11 — *EBF3* is a dosage-sensitive regulator of neuronal migration acting downstream of *ARX* and within a Prdm8–EBF3–NeuroD1 axis

Developmental neurobiology studies in mouse and in vitro place EBF3 in a defined gene-regulatory hierarchy controlling neuronal migration:

- *Ebf2* and *Ebf3*, singly or together, control the migration of Cajal–Retzius cells arising from the cortical hem (PMID: 22421355): *"both Ebf2 and Ebf3, singly or together, control the migration of CR cells arising in the cortical hem."*
- The homeobox selector gene *Arx* directly represses *Ebf3*; ectopic *Ebf3* expression in the ventral telencephalon prevents neuronal tangential migration, and silencing *Ebf3* in *Arx*-mutant tissue partially rescues tangential cell movement (PMID: 19627984: *"its ectopic expression in ventral telencephalon was found to prevent neuronal tangential migration"*; *"Arx is sufficient to repress Ebf3 endogenous expression"*; see also PMID: 18799476).
- *Ebf3* is a *Prdm8* target that drives the multipolar-to-bipolar transition of migrating neocortical neurons via positive feedback on *NeuroD1* (PMID: 29113800: *"Ebf3 is a key regulator of neuronal migration via multipolar-to-bipolar transition"*).

This hierarchy explains the phenotypic overlap between EBF3 and ARX disorders and links reduced EBF3 dosage to defective cortical and cerebellar neuronal migration — the inferred substrate for the ataxia, hypotonia, developmental delay, and the dandelion-sign cerebellar malformation.

Evidence type: model organism (mouse) and in vitro developmental biology.

Finding 12 — No disease-specific therapy exists; management is supportive; the disorder is ultra-rare

A ClinicalTrials.gov query (2026-11) returned no interventional/therapeutic trial specific to *EBF3* or HADDS; the only EBF3-tagged study is the observational Simons Searchlight registry (NCT01238250), which lists *EBF3* among >150 neurodevelopmental genes. HADDS is ultra-rare — the reported literature comprises small case series (first described in 2017; the largest aggregate review assembled 52 patients, PMID: 37718233: *"We identified 52 patients (33F:19M) through literature (28F:18M) and retrospective review (5F:1M)"*) — and no formal prevalence/incidence estimate is published (Orphanet:658843). Reported management across series is multidisciplinary supportive care: physical/occupational/speech therapy; treatment of neurogenic bladder (clean intermittent catheterization/cystostomy, surveillance for VUR/UTI and renal protection); ophthalmologic management of strabismus; and developmental/educational support.

Evidence type: clinical trial registry, case-series synthesis.


Mechanistic Model / Interpretation

Ordered causal chain (initiating lesion → clinical manifestation)

1. A **heterozygous loss-of-function lesion in *EBF3*** arises — most often *de novo* (missense in the DNA-binding/ZNF domain, nonsense, frameshift, splice-site), or as a whole-gene 10q26.3 deletion, or (rarely) a noncoding enhancer hs737 variant. *This leads to →*
2. **Reduced dosage of functional EBF3 protein** (haploinsufficiency). Truncating alleles trigger nonsense-mediated decay; missense alleles in the zinc-knuckle impair folding/DNA contact. *This results in →*
3. **Impaired EBF3 DNA binding and transcriptional activation.** Mutant EBF3 shows reduced/absent transactivation of targets (e.g., *CDKN1A*), cytoplasmic mislocalization, weaker chromatin association, and genome-wide loss of binding. *This leads to →*
4. **Dysregulation of EBF3-dependent neurodevelopmental gene programs.** EBF3 normally acts downstream of *ARX* and within a *Prdm8 → EBF3 → NeuroD1* feedback loop. Loss of EBF3 output disrupts the multipolar-to-bipolar transition and tangential/radial migration of cortical and Cajal–Retzius neurons. *This results in →*

5. **Abnormal neuronal migration and cerebellar/cortical patterning** — inferred to produce cerebellar vermis hypoplasia with radial "dandelion-sign" foliation and white-matter (longitudinal fasciculus) abnormalities. *This leads to →*
6. **Clinical manifestations:** congenital hypotonia, ataxia, global developmental delay with prominent expressive-speech impairment, intellectual disability (variable), facial dysmorphism, strabismus, and neurogenic bladder (via inferred detrusor-sphincter dyssynergia).

Branch point: severity tracks with **where** the missense variant falls — ZNF-domain variants (which destabilize the EBF3–DNA interaction) define a high-severity subtype, whereas certain DNA-binding-domain variants such as p.Arg209Trp retain partial function and are associated with milder outcomes, including rare normal-IQ cases.



Ontology annotations

- **Gene/protein:** *EBF3* (HGNC:19087), COE-family DNA-binding transcription factor.
- **GO biological process:** GO:0001764 (neuron migration), GO:0021954 (central nervous system neuron development), GO:0045944 (positive regulation of transcription by RNA Pol II), GO:0006357 (regulation of transcription by RNA Pol II).

- **GO cellular component:** GO:0005634 (nucleus), GO:0000785 (chromatin); mutants aberrantly in GO:0005737 (cytoplasm).
- **Cell types (CL):** CL:0000099 (interneuron), CL:0012001 (neuron of the cerebral cortex), Cajal–Retzius cell, cerebellar neurons.
- **Anatomy (UBERON):** UBERON:0004720 (cerebellar vermis), UBERON:0002037 (cerebellum), UBERON:0000955 (brain), UBERON:0001255 (urinary bladder).
- **Disease (MONDO):** MONDO:0015021.

Evidence Base

PMID	Title (abbrev.)	Evidence type	Role in this report
28017372	*A Syndromic NDD Caused by De Novo Variants in EBF3*	Human clinical + fly	Founding paper; de novo variants, ZNF localization (F001, F002)
28017373	*Mutations in EBF3 Disturb Transcriptional Profiles...*	In vitro + human	Transactivation/ChIP-seq loss; mosaic inheritance (F002, F005)
29162653	*De novo variants in EBF3*	Human clinical	Core phenotype delineation (hypotonia, ataxia, dysmorphism)
34050706	*Clinical spectrum of EBF3 variants/deletions*	Human clinical	Haploinsufficiency; deletion phenocopy; early neurogenic bladder (F001, F003)
37718233	*Urologic manifestations of HADDS*	Human clinical	52-patient series; DSD mechanism; ultra-rarity (F003, F012)
37090941	*Neurologic/neuroradiologic features*	Human clinical	Dandelion-sign cerebellar malformation; SARA ataxia (F004)
36317217	*Clinical features and EBF3 variant in a child*	Human clinical	Recurrent p.Arg209Trp, de novo (F005)
35340043	*High-risk subtype for ZNF-domain missense*	Human + fly/ luciferase	ZNF severity genotype–phenotype (F006)
34256850	*Coding and noncoding variants in EBF3*	Human WGS	Enhancer hs737 DNVs in autism (F007)
36937983	*EBF3-related NDD in 12 Chinese patients*	Human clinical	Phenotype frequencies; variable expressivity (F010)
34177436	*Neuroimaging findings in EBF3-NDD*	Human clinical	Normal-IQ cases; DTI white-matter asymmetry (F010)
40073166	*Nonsense variant in C-terminal transactivation domain*	Human clinical	C-terminal truncation; behavioral phenotype expansion
22421355	*EBF2/3 regulate Cajal–Retzius migration*	Mouse	Migration role (F011)
19627984		Mouse	

PMID	Title (abbrev.)	Evidence type	Role in this report
	Arx as regional key selector gene		ARX represses EBF3; tangential migration (F011)
18799476	*Arx transcriptional targets in basal forebrain*	Mouse	EBF3 a direct ARX target (F011)
29113800	*Prdm8 target Ebf3 regulates multipolar-to-bipolar transition*	Mouse/in vitro	Prdm8–EBF3–NeuroD1 axis (F011)
42711312	*Non-cell-autonomous roles for UNC-3/EBF*	*C. elegans*	Ortholog circuit-level mechanism; CEBP-1 driver
42182317	*Negative autoregulation of dosage-sensitive TF*	*C. elegans*	EBF/UNC-3 dosage sensitivity concept (supports F008)

Consistency across the evidence base: The haploinsufficiency model (F001, F008) is supported convergently by human genetics (deletion phenocopy), population constraint (pLI ≈ 1), and in vitro loss of function (F002). The migration mechanism (F011) is drawn entirely from model-organism work and is therefore an *inferred* explanation for the human imaging phenotype rather than a directly demonstrated human mechanism. No contradicting evidence was identified; the main tension is the breadth of expressivity (F010), which the ZNF-severity model (F006) partially reconciles.

Section-by-Section Characterization

1. Disease Information. HADDS (MONDO:0015021; OMIM:617330; Orphanet:658843) is an autosomal-dominant neurodevelopmental disorder caused by *EBF3*. Synonyms: HADDS; EBF3 neurodevelopmental disorder; developmental delay–ataxia–hypotonia–facial dysmorphism syndrome. No dedicated ICD-10/11 or MeSH code; it maps to generic neurodevelopmental/cerebellar categories. Information is derived from aggregated disease-level resources and small patient case series, not large EHR datasets.

2. Etiology. Primary cause is genetic — heterozygous LoF variants in *EBF3* (germline, usually *de novo*; rarely transmitted via parental mosaicism). No environmental, infectious, or lifestyle causes are known. The only recognized recurrence-risk modifier is parental germline/somatic mosaicism (F005). No protective alleles or gene–environment interactions are established.

3. Phenotypes. Near-universal: global developmental delay (HP:0001263) with expressive-speech impairment. Common: hypotonia (HP:0001252), ataxia (HP:0001251), intellectual disability (HP:0001249, variable), facial dysmorphism, strabismus (HP:0000486), neurogenic bladder (HP:0000011), cerebellar vermis hypoplasia (HP:0001320). Onset is congenital/neonatal; course is largely static; severity is variable (F006, F010). Quality-of-life impact is driven by mobility/communication impairment and urologic morbidity.

4. Genetic/Molecular. Causal gene *EBF3* (HGNC:19087, gene MIM 607407, chr10q26.3). ~230 P/LP ClinVar records; recurrent DNA-binding residues Arg163 and Arg209; variant classes include missense, nonsense, frameshift, splice, and whole-gene deletion (F009). Functional consequence: loss of function/haploinsufficiency (F001, F002, F008). No established modifier genes beyond the ZNF-domain positional effect; no disease-specific epigenetic mechanism reported.

5. Environmental. Not applicable — HADDS is a monogenic disorder with no known environmental, lifestyle, or infectious contributors.

6. Mechanism/Pathophysiology. See the causal chain above (F001–F002, F008, F011).

7. Anatomical Structures. Primary: central nervous system — cerebellum/vermis (UBERON:0004720), cerebral cortex, white-matter long tracts. Secondary: urinary bladder (UBERON:0001255) and upper urinary tract (via VUR/neurogenic bladder); eyes (strabismus). Subcellular: nucleus/chromatin (normal EBF3 localization) with pathologic cytoplasmic mislocalization of mutants.

8. Temporal Development. Congenital onset; insidious recognition in infancy. Static/non-progressive neurodevelopmental course (lifelong). Neurogenic bladder presents early (median 6.5 months). Critical period is early neurodevelopment; the main intervention window is early supportive/rehabilitative therapy and renal-protective urologic surveillance.

9. Inheritance and Population. Autosomal dominant; mostly *de novo*; incomplete reporting of penetrance but highly variable expressivity. No formal prevalence/incidence (ultra-rare). Female predominance in ascertained urologic cohorts (33F:19M) likely reflects ascertainment and the urologic phenotype rather than a true sex bias. Parental mosaicism documented; no founder effects or consanguinity role established.

10. Diagnostics. Molecular diagnosis via exome/genome sequencing, NDD gene panels including *EBF3*, single-gene testing, or chromosomal microarray (detects 10q26.3 deletions). Brain MRI (cerebellar vermis hypoplasia, dandelion-sign foliation) and SARA ataxia scoring support the clinical picture; urodynamic/renal imaging for bladder dysfunction. Differential diagnoses include other congenital ataxias/cerebellar hypoplasias, ARX-related disorders, and 10q26 deletion/ring-chromosome-10 syndromes.

11. Outcome/Prognosis. Lifelong disability with static course; no reported reduction in survival from the neurodevelopmental phenotype itself, though neurogenic bladder poses a risk of renal complications if unmanaged. Prognostic factor: ZNF-domain missense variants predict greater severity. No validated prognostic biomarkers.

12. Treatment. No disease-modifying therapy. Supportive/rehabilitative care: physical, occupational, and speech therapy; clean intermittent catheterization or cystostomy and VUR/UTI surveillance for neurogenic bladder; ophthalmologic management of strabismus; educational support. Suggested NCIT terms: physical therapy, occupational therapy, speech therapy, intermittent catheterization.

13. Prevention. No primary prevention. Secondary/tertiary measures are genetic counseling (recognizing mosaicism-related recurrence risk), prenatal/preimplantation testing for known familial variants, and proactive renal-protective urologic surveillance to prevent complications.

14. Other Species / Natural Disease. Orthologs: mouse *Ebf3*, *C. elegans unc-3*, *Drosophila knot/collier*-family. No naturally occurring companion-animal disease is cataloged; relevance is as experimental models. Mechanisms are evolutionarily conserved (migration, dosage-sensitive terminal-selector function).

15. Model Organisms. *Drosophila* (UAS-GAL4 rescue assays demonstrating variant deleteriousness and ZNF severity), mouse (*Ebf3*/*Ebf2* migration studies, Prdm8–EBF3–NeuroD1 axis), and *C. elegans* (*unc-3* dosage-sensitivity and non-cell-autonomous circuit roles). Models recapitulate the transcription-factor loss-of-function and neuronal-migration phenotypes but do not fully reproduce the human cerebellar/urologic clinical syndrome.

Limitations and Knowledge Gaps

1. **Mechanism is largely inferred for humans.** The neuronal-migration causal chain (F011) rests on mouse and invertebrate data; the direct link between EBF3 haploinsufficiency and

the human dandelion-sign cerebellar malformation has not been demonstrated at the tissue level in patients.

2. **No epidemiology.** No prevalence or incidence estimate exists (Orphanet:658843); the total reported literature is a few hundred patients at most, limiting frequency estimates for individual phenotypes.
3. **Ascertainment bias.** Female predominance and the high urologic burden derive from cohorts partly ascertained through urology clinics; true population sex ratio and penetrance of each feature are uncertain.
4. **Genotype–phenotype resolution is incomplete.** The ZNF-severity correlation (F006) is robust but does not explain normal-IQ outliers (F010) or the full spectrum; C-terminal transactivation-domain variants are under-characterized.
5. **No therapeutics.** There are no interventional trials, no biomarkers of progression, and no validated outcome measures beyond SARA for ataxia.
6. **Epigenetic, metabolomic, and proteomic data are absent** for this disorder specifically.

Proposed Follow-up Experiments / Actions

1. **Human neuropathology/imaging correlation.** Systematic volumetric MRI and, where available, neuropathologic study to directly test whether the dandelion-sign malformation reflects the migration defect predicted by mouse models.
2. **Variant-function atlas.** Deep mutational scanning of *EBF3* across DNA-binding, ZNF, and transactivation domains with transactivation/localization readouts to build a quantitative severity map and reclassify variants of uncertain significance.
3. **Prospective natural-history registry.** Expand the Simons Searchlight (NCT01238250) EBF3 cohort with standardized developmental, ataxia (SARA), and urodynamic assessments to establish phenotype frequencies, penetrance, and progression.
4. **Urologic mechanism study.** Test the detrusor-sphincter-dyssynergia hypothesis directly with urodynamics in a defined EBF3 cohort and correlate with variant type.
5. **Dosage-restoration proof of concept.** Given the haploinsufficiency mechanism and extreme constraint, evaluate dosage-upregulation strategies (e.g., ASO-mediated or CRISPR-activation of the wild-type allele, enhancer hs737 targeting) in neuronal/organoid models.

6. **iPSC and brain-organoid models** from patients carrying ZNF-domain vs DNA-binding vs truncating variants to resolve genotype-specific cellular phenotypes and screen candidate modifiers.

Report compiled from 12 confirmed findings and 27 reviewed papers over 5 investigation iterations. Evidence types are labeled per finding (human clinical, model organism, in vitro, computational). Causal-chain steps that rest on model-organism data are flagged as inferred.
