

Müllerian Aplasia and Hyperandrogenism (MONDO:0008019 / OMIM #158330): A Comprehensive Disease Characterization Report

Summary

Müllerian Aplasia and Hyperandrogenism (OMIM #158330; MONDO:0008019) is a rare **Mendelian, autosomal-dominant** disorder caused by **heterozygous loss-of-function (missense) variants in *WNT4*** (1p36.12). In 46,XX women it produces a distinctive dyad: **congenital aplasia of the uterus and upper vagina** (Müllerian agenesis) combined with **clinical and biochemical androgen excess**. It represents the "atypical," hyperandrogenic form of Mayer-Rokitansky-Küster-Hauser (MRKH) syndrome and is, to date, the only MRKH subtype with an established single-gene (Mendelian) cause. The index description was a landmark 2004 report of an 18-year-old 46,XX woman with primary amenorrhea, absent Müllerian derivatives, unilateral renal agenesis, and androgen excess, carrying a heterozygous *WNT4* loss-of-function allele ([PMID: 15317892](#)).

Mechanistically, *WNT4* serves a dual developmental role in the female reproductive tract. During embryogenesis it is required for **formation of the Müllerian (paramesonephric) duct**; in the differentiating ovary it **represses steroidogenic enzyme expression**, thereby suppressing ectopic androgen synthesis. Loss of *WNT4* function therefore simultaneously **abolishes Müllerian-duct development** and **de-represses ovarian androgen biosynthesis** (via re-expression of CYP17A1/17 α -hydroxylase and HSD3B2/3 β -hydroxysteroid dehydrogenase), explaining the coincidence of Müllerian aplasia and hyperandrogenism in a single gene defect. This mechanism is strongly supported by the *Wnt4*-null mouse, in which females lack the Müllerian duct, retain the Wolffian duct, and ectopically activate testosterone biosynthesis — a phenotype the human disorder was described as "remarkably similar to" ([PMID: 9989404](#); [PMID: 15317892](#)).

The disorder is **non-lethal**, with the principal burden being **absolute uterine-factor infertility** and substantial **psychosocial impact** (elevated anxiety/depression, impaired sexual function). Management is **symptomatic and multidisciplinary**: non-surgical vaginal dilation (first-line) or surgical neovagina creation, anti-androgen therapy for androgen excess, assisted reproduction (IVF with gestational surrogacy, and increasingly uterine transplantation), and psychological support. It must be carefully distinguished from the dominant disorder's **recessive, embryonic/perinatally lethal counterpart** — **SERKAL syndrome**, caused by **biallelic (homozygous) *WNT4* mutation**, which features 46,XX female-to-male sex reversal with renal, adrenal, and lung dysgenesis ([PMID: 18179883](#)).

1. Disease Information

Overview. Müllerian Aplasia and Hyperandrogenism is a congenital disorder of the female reproductive tract in genotypic (46,XX) females, defined by the combination of **Müllerian aplasia** (absence/hypoplasia of the uterus and upper vagina — "uterovaginal aplasia") and **hyperandrogenism** (clinical signs such as hirsutism/acne and/or biochemical elevation of androgens such as testosterone). It is nested within the broader **MRKH / Müllerian agenesis** spectrum but is set apart by androgen excess and by its monogenic *WNT4* etiology. Ovaries are present and typically of normal size on imaging, but are the source of excess androgens; secondary sexual characteristics are otherwise female. Patients characteristically present at adolescence with **primary amenorrhea** and are found to have a 46,XX karyotype.

Key identifiers.

Resource	Identifier
OMIM	#158330 ("Müllerian aplasia and hyperandrogenism"; <i>WNT4</i> -related)
MONDO	MONDO:0008019
Gene (HGNC)	<i>WNT4</i> , HGNC:12783; locus 1p36.12; gene OMIM *603490
Orphanet	Within the MRKH / Müllerian aplasia spectrum (atypical/hyperandrogenic form)
ICD-10	Q51.x (congenital malformations of uterus/cervix); vaginal agenesis Q52.x
ICD-11	LA14 (structural developmental anomalies of female genitalia)
MeSH	46,XX Disorders of Sex Development; Müllerian aplasia / Mullerian ducts

Synonyms / alternative names. Müllerian aplasia and hyperandrogenism; *WNT4* deficiency; atypical MRKH syndrome with hyperandrogenism; MRKH-like syndrome with Müllerian-duct regression and virilization; *WNT4* Müllerian aplasia. (Note: the classic, androgen-normal form is "typical MRKH" and is genetically distinct — see §10.)

Source of information. The characterization derives predominantly from **aggregated disease-level resources** (OMIM, review articles) and a small number of **individual-patient reports and small cohorts** (the index case and subsequent *WNT4*-mutation-positive adolescents), supplemented by **model-organism** (mouse) and **in-vitro functional** data. There is no large EHR-derived cohort specific to the *WNT4* hyperandrogenic form; EHR-based data exist for the broader Müllerian aplasia category ([PMID: 39391975](#)).

2. Etiology

Primary cause (genetic). The disorder is caused by **heterozygous loss-of-function missense variants in *WNT4***. The index patient carried p.Glu226Gly (p.E226G) ([PMID: 15317892](#)); subsequently reported variants include p.Leu12Pro (p.L12P, exon 1; [PMID: 18182450](#)) and p.Ala233Thr (p.A233T; [PMID: 21377155](#)). These are **germline, dominant-acting** alleles producing a partial loss of *WNT4* signalling.

Genetic risk factors. The causal variants themselves are the risk determinant; no additional susceptibility loci or validated modifier genes have been established for the hyperandrogenic *WNT4* form. Within the broader MRKH spectrum, *WNT4* is the only gene with mutations definitively linked to an atypical (hyperandrogenic) phenotype: "The only known MRKH-associated mutations are located within the *WNT4* gene and lead to an atypical form of MRKH syndrome associated with clinical and biochemical hyperandrogenism" ([PMID: 24641817](#)).

Environmental risk factors. None established as causal for the Mendelian *WNT4* form. For MRKH broadly, no reproducible environmental cause has been identified.

Protective factors. No genetic or environmental protective factors have been defined. Mechanistically, retention of one functional *WNT4* allele with sufficient residual signalling may mitigate severity (variable expressivity; see §9), but this is inferred rather than demonstrated.

Gene–environment interactions. None documented. The phenotype is driven by the monogenic lesion; environmental modulation of androgen expression has not been characterized in this disorder.

3. Phenotypes

Phenotypes fall into two cardinal domains — **structural (Müllerian aplasia)** and **endocrine (hyperandrogenism)** — plus extragenital features (chiefly renal).

Phenotype	Type	HPO suggestion	Onset	Frequency / notes
Primary amenorrhea	Symptom/sign	HP:0000786	Adolescence (presenting)	Near-universal presenting feature
Aplasia/hypoplasia of uterus	Physical malformation	HP:0000013 (Hypoplasia of the uterus)	Congenital	Defining; ranges hypoplastic→absent
Vaginal aplasia (absent upper vagina)	Physical malformation	HP:0000148 (Vaginal atresia)	Congenital	Defining
Hyperandrogenism (clinical)	Clinical sign	HP:0000858 (Hyperandrogenism)	Peripubertal/adult	Defining of this subtype
Hirsutism	Physical manifestation	HP:0001007	Peripubertal/adult	Variable
Acne (can be severe)	Physical manifestation	HP:0001061	Peripubertal	Present in index/variant carriers; acne overall less frequent in MRKH (24.6%)
Elevated serum testosterone	Lab abnormality	HP:0030348 (Increased circulating androgen level)	Peripubertal/adult	e.g., 1.8 vs 1.2 nmol/L in a carrier
Unilateral renal agenesis	Physical malformation	HP:0000122 / HP:0000104	Congenital	In index case; renal anomalies ~15.5% in MRKH
Follicle depletion / ovarian dysfunction	Lab/structural	HP:0008222 (Decreased ovarian reserve)	Variable	Reported in L12P carrier

Characteristics. - **Age of onset:** The structural defects are **congenital**; clinical recognition is typically at **adolescence** when primary amenorrhea prompts evaluation. Hyperandrogenic signs emerge peri-/post-pubertally. - **Severity: Variable.** Uterine involvement ranges from hypoplasia to complete absence; androgen excess ranges from biochemical only to clinically overt (severe acne, hirsutism). - **Progression:** The structural anomaly is **stable (non-**

progressive); androgen-related manifestations are **chronic** and may be modulated by therapy.

- **Frequency among affected:** Müllerian aplasia and androgen excess are the defining, essentially obligate features of this subtype; extragenital (renal) involvement is present in a subset.

Quality-of-life impact. The dominant QoL burdens are **infertility-related distress** and **sexual dysfunction** (dyspareunia, impaired arousal/lubrication/orgasm), plus psychological morbidity. A systematic review of MRKH reported that "MRKH could be associated with a higher prevalence of anxiety and depression symptoms and social insecurity compared with women of a similar age without the condition," and found impaired mental-health-related QoL though physical-health-related QoL was not affected ([PMID: 35455528](#)). An EHR cohort found anxiety/depressive disorders ~2× higher than male referents in Müllerian aplasia ([PMID: 39391975](#)).

4. Genetic / Molecular Information

Causal gene. *WNT4* (Wnt family member 4), HGNC:12783, locus **1p36.12**, gene OMIM *603490. *WNT4* is a secreted glycoprotein ligand of the Wnt signalling family.

Pathogenic variants (germline, heterozygous).

Variant (protein)	Exon	Source	Functional consequence
p.E226G	—	Index case, PMID: 15317892	Loss-of-function
p.L12P	exon 1	PMID: 18182450 ; 1/28 MRKH girls, absent in 100 controls	↑ expression of HSD3B2 and CYP17A1 (de-repression)
p.A233T	—	PMID: 21377155	Partial loss of repression; abnormal CYP17A1 re-expression

- **Variant classification (ACMG/AMP):** The recurrent/characterized missense alleles behave as **pathogenic/likely pathogenic** on the strength of functional assays, segregation, and absence in controls; unselected MRKH screening yields many variants of uncertain significance, and the diagnostic yield of *WNT4* in unselected MRKH is low (~1/28) ([PMID: 18182450](#)).
- **Variant type/class:** **Missense** (functional loss-of-function), not gross structural.

- **Allele frequency:** Pathogenic alleles are **private/rare**, absent from control panels (e.g., absent in 100 controls for L12P) and population databases.
- **Somatic vs germline: Germline.**
- **Functional consequence: Loss of function** — reduced WNT4/ β -catenin signalling. WNT4 loss downregulates WNT4-dependent inhibition of β -catenin degradation (PMID: 18179883), and de-represses ovarian steroidogenic enzymes: "This mutant partially lacks the capability to repress ovarian steroidogenic enzymes, with abnormal expression of 17 α -hydroxylase" (PMID: 21377155); and the L12P mutation "induces significantly increased expression of the enzymes involved in androgen biosynthesis (3 β -hydroxysteroid dehydrogenase and 17 α -hydroxylase)" (PMID: 18182450).

Modifier genes. None validated for this disorder. Pathway partners RSPO1 and FOXL2 are mechanistically related (see §6) but are not established modifiers of the *WNT4* phenotype.

Epigenetic information. No disease-specific epigenetic signature has been defined for Müllerian aplasia and hyperandrogenism. (Of pathway interest, DNA methylation via DNMT1 regulates nephron progenitor renewal in kidney development — PMID: 30518531 — but this is not a documented mechanism in this disorder.)

Chromosomal abnormalities. The hyperandrogenic *WNT4* form is a **point-mutation (single-gene)** disorder; it is not caused by aneuploidy or large rearrangements. (By contrast, portions of the broader MRKH spectrum involve copy-number variants, e.g., 16p11.2 and 17q12 — not specific to the *WNT4* subtype.)

5. Environmental Information

- **Environmental factors:** None established as causal.
 - **Lifestyle factors:** None established as causal; weight/obesity may modulate androgenic phenotype expression generally but is not a defined cause here.
 - **Infectious agents:** Not applicable — this is a genetic developmental disorder with no infectious etiology.
-

6. Mechanism / Pathophysiology

Ordered causal chain

1. A **heterozygous loss-of-function missense variant in WNT4** (e.g., p.E226G, p.L12P, p.A233T) **reduces** secreted WNT4 ligand activity.
2. Reduced WNT4 **decreases** WNT4-dependent inhibition of β -catenin degradation, which **lowers** canonical Wnt/ β -catenin signalling in the developing reproductive tract (demonstrated in vitro — [PMID: 18179883](#)).
3. **Branch A (structural):** Loss of WNT4 signalling in the paramesonephric mesenchyme **prevents formation of the Müllerian (paramesonephric) duct**, which **results in** uterine and upper-vaginal aplasia ($\rightarrow$ primary amenorrhea, absolute uterine-factor infertility). Demonstrated in mouse: "the Mullerian duct is absent while the Wolffian duct continues to develop" ([PMID: 9989404](#)).
4. **Branch B (endocrine):** Loss of WNT4's repressive action in the differentiating ovary **de-represses ovarian steroidogenic enzymes — abnormal re-expression of CYP17A1 (17 α -hydroxylase)** and increased **HSD3B2 (3 β -hydroxysteroid dehydrogenase)** — which **leads to** ectopic/elevated ovarian androgen (testosterone) biosynthesis ($\rightarrow$ hyperandrogenism, hirsutism, acne). Demonstrated functionally in OVCAR3 cells for p.A233T ([PMID: 21377155](#)) and for p.L12P ([PMID: 18182450](#)); in mouse, "Wnt-4 in the developing ovary appears to suppress the development of Leydig cells; consequently, Wnt-4-mutant females ectopically activate testosterone biosynthesis" ([PMID: 9989404](#)).
5. **Branch C (renal, subset):** Because WNT4 is also required in nephrogenesis (Wnt4-null mutants fail to form nephron structures — [PMID: 41358813](#)), the lesion can **result in** renal anomalies such as unilateral renal agenesis in a subset of patients ([PMID: 15317892](#)).
6. The convergence of Branches A + B in a 46,XX woman **produces** the clinical syndrome: uterovaginal aplasia with androgen excess.

Detailed mechanism

Molecular pathways. WNT4 acts within the **canonical Wnt/ β -catenin pro-ovarian pathway** together with **RSPO1** (R-spondin-1) and **FOXL2**, which collectively promote ovarian fate and antagonize the male (SRY/SOX9) testis-determining program. "The male pathway is opposed by the expression of two signals, WNT4 and R-SPONDIN-1 (RSPO1), that promote the ovarian fate and block testis development. Both of these ligands can activate the canonical Wnt signaling pathway" ([PMID: 18617533](#)). "Female sex-determining genes RSPO1, Wnt4/ β -catenin and Foxl2,

are involved in the formation of female genitalia and the maintenance of female ovarian development" (PMID: 27481580). Ovarian somatic differentiation is regulated by FOXL2 and the β -catenin pathway activated by Wnt4 and Rspo1 (PMID: 22251856). Suggested pathway/GO terms: **Wnt signaling pathway** (GO:0016055), **canonical Wnt signaling pathway** (GO:0060070), **female sex differentiation** (GO:0046660), **Müllerian duct development** (GO:0061205), **negative regulation of steroid biosynthetic process** (GO:0010894).

Cellular processes. Developmental patterning and lineage specification — **Müllerian duct formation, ovarian somatic (pre-granulosa) differentiation, and suppression of ectopic steroidogenic (Leydig-like) cell development.** Loss leads to failure of duct morphogenesis (Branch A) and inappropriate steroidogenic gene expression (Branch B).

Protein dysfunction. Missense WNT4 variants cause **loss of function** (reduced ligand signalling), not aggregation or classic dominant-negative gain; the heterozygous dominant phenotype reflects **haploinsufficiency/partial loss** sufficient to perturb dosage-sensitive developmental and repressive functions.

Metabolic/biochemical abnormalities. The core biochemical defect is **de-repression of steroidogenic enzymes: CYP17A1 (17 α -hydroxylase/17,20-lyase) and HSD3B2 (3 β -HSD),** driving **androgen (testosterone) overproduction** (CHEBI: testosterone CHEBI:17347; androgen CHEBI:50113).

Immune system / tissue damage. Not implicated — this is a developmental/endocrine disorder, not inflammatory, fibrotic, or ischemic.

Molecular profiling. In-vitro transcriptional assays (OVCAR3, steroidogenic cell models) demonstrate the enzyme de-repression; no large-scale transcriptomic/proteomic/metabolomic atlas exists specific to this disorder.

Cell types (CL suggestions): ovarian granulosa cell (CL:0000501), theca/steroid-hormone-secreting cell, Müllerian duct epithelial/mesenchymal cells, nephron progenitor cell (for renal branch).

WNT4 LOF (het missense)

| ↓ canonical Wnt/ β -catenin signalling

| → Müllerian duct NOT formed → uterine + upper-vaginal aplasia → primary amenorrhea

| → loss of ovarian repression → CYP17A1↑ / HSD3B2↑ → ↑androgens → hirsutism, acne, etc.

| → nephron defect (subset) → renal agenesis/anomaly

7. Anatomical Structures Affected

Organ level. - **Primary:** Uterus (UBERON:0000995) and upper vagina (UBERON:0000996) — aplastic/hypoplastic. Ovaries (UBERON:0000992) are present but functionally abnormal (androgen over-secreting). - **Secondary / associated:** Kidney (UBERON:0002113) — unilateral renal agenesis in a subset. Skin/pilosebaceous unit secondarily affected by androgen excess (hirsutism, acne). - **Body systems:** **Reproductive/genitourinary** (primary) and **endocrine** (androgen axis); **renal/urinary** in a subset.

Tissue and cell level. Müllerian-duct-derived **epithelial and mesenchymal** tissue (fails to develop); **ovarian somatic cells** (granulosa/theca-steroidogenic lineage) exhibit aberrant steroidogenesis. Suggested CL terms: granulosa cell (CL:0000501), steroidogenic/theca cell.

Subcellular level. **Mitochondria** and **smooth endoplasmic reticulum** (sites of steroidogenesis — CYP17A1 resides in the ER); **nucleus** (transcriptional de-repression of steroidogenic genes); **plasma membrane/extracellular** (WNT4 ligand secretion and receptor signalling). GO cellular-component suggestions: extracellular space (GO:0005615), endoplasmic reticulum membrane (GO:0005789).

Localization / lateralization. Uterovaginal aplasia is **midline/bilateral** (a single fused midline structure fails to form). Renal agenesis, when present, is typically **unilateral** ([PMID: 15317892](#)).

8. Temporal Development

- **Onset:** Structural anomalies are **congenital**; clinical presentation is at **adolescence** (primary amenorrhea, often ages 14–18). Onset pattern is **insidious/chronic** — the anomaly is present from birth but detected when menarche fails.
 - **Progression:** The anatomical defect is **stable and non-progressive**. Hyperandrogenism is **chronic**, variable, and treatment-responsive. There are no formal disease "stages."
 - **Course / duration:** **Chronic, lifelong** congenital condition.
 - **Patterns:** No spontaneous remission of the structural defect; androgenic features can be controlled (treatment-induced improvement). **Critical periods:** the vulnerable developmental window is **embryonic reproductive-tract morphogenesis** (Müllerian duct and ovarian differentiation); the therapeutic windows are adolescence/adulthood for neovagina creation and fertility interventions.
-

9. Inheritance and Population

Epidemiology. Precise prevalence of the *WNT4* hyperandrogenic subtype is **not established** (very rare; described in case reports and small cohorts). For context, **MRKH overall affects ~1:4,500 women** (PMID: 28434104) and is "the second most common cause of primary amenorrhea, accounting for 10%-15% of cases" (PMID: 41636313). Hyperandrogenemia is biochemically common across MRKH cohorts (60.1% in one series) but overt acne (24.6%) and PCOS are relatively infrequent, and *WNT4* mutations account for only the rare atypical, genetically-defined subset (PMID: 24641817).

For genetic etiology. - **Inheritance pattern: Autosomal dominant** (heterozygous *WNT4* LOF). Contrast with the **autosomal recessive** biallelic-*WNT4* disorder SERKAL (PMID: 18179883). - **Penetrance / expressivity: Variable expressivity**; most reported variants arise as rare/private alleles. Penetrance is presumed high for the structural defect in carriers described, but formal estimates are lacking given the small number of families. - **Genetic anticipation:** Not applicable (not a repeat-expansion disorder). - **Germline mosaicism / founder effects / consanguinity:** Not documented for the dominant form. (Consanguinity is relevant to recessive SERKAL.) - **Carrier frequency:** Pathogenic alleles are **private/rare**; no meaningful population carrier frequency for the dominant disorder.

Population demographics. - **Affected populations:** Reported across multiple populations; no established ethnic predilection for the *WNT4* subtype. - **Geographic distribution:** No endemic pattern (Mendelian, globally distributed). - **Sex ratio:** Affects **46,XX (phenotypic females)** — effectively female-only presentation given the reproductive-tract phenotype. - **Age distribution:** Diagnosed predominantly in **adolescence/young adulthood**.

10. Diagnostics

Clinical tests. - **Laboratory:** Karyotype confirming **46,XX**; **serum androgens** (testosterone — elevated, e.g., 1.8 vs control 1.2 nmol/L in a carrier; also free testosterone, androstenedione, DHEAS), gonadotropins (FSH/LH), estradiol, AMH (ovarian reserve); 17-hydroxyprogesterone to screen for CAH. LOINC-codable panels for testosterone and gonadotropins. - **Biomarkers:** Elevated circulating androgens are the key biochemical discriminator; *WNT4* mutation is the molecular biomarker. - **Imaging: Pelvic ultrasound and MRI** are the mainstay to document

uterovaginal aplasia and confirm presence of ovaries (ultrasound and MRI are the main diagnostic techniques — [PMID: 28434104](#)). Renal imaging to detect associated renal agenesis. - **Biopsy/pathology:** Not routinely required for diagnosis.

Genetic testing. Recommended approach: confirm 46,XX karyotype, then **targeted *WNT4* sequencing** (single-gene) or an MRKH/DSD **gene panel**; **WES/WGS** are useful for atypical/syndromic cases or when panels are negative. Chromosomal microarray can detect CNVs relevant to the broader MRKH spectrum (not the *WNT4* point-mutation form). Yield of *WNT4* in unselected MRKH is low (~1/28), so testing is highest-yield in **hyperandrogenic** MRKH ([PMID: 18182450](#)).

Clinical criteria & differential diagnosis. Diagnosis rests on 46,XX uterovaginal aplasia **plus androgen excess**. Key differentials:

Condition	Karyotype	Uterus	Gonads	Androgens	<i>WNT4</i>	Discriminator
WNT4 Müllerian aplasia + hyperandrogenism	46,XX	Absent/hypoplastic	Ovaries	High	Mutated	Androgen excess + <i>WNT4</i>
Classic (typical) MRKH	46,XX	Absent	Ovaries	Normal	Negative	Normal androgens
CAIS (complete androgen insensitivity)	46,XY	Absent	Testes	— (resistance)	—	46,XY, AR mutation, no ovaries
MURCS association	46,XX	Absent	Ovaries	±	Not causal	Skeletal/renal; <i>WNT4</i> excluded (PMID: 24356390)
PCOS / androgen-secreting tumor / CAH	46,XX	Present	Ovaries/adrenal	High	—	Uterus present; distinct endocrine profile

A MURCS patient with obesity and hyperandrogenism showed no exonic *WNT4* mutation: "Our finding excludes *WNT4* gene as a candidate for MURCS association" ([PMID: 24356390](#)).

Screening. No population newborn screening. **Cascade genetic testing** and genetic counseling are appropriate for families with an identified *WNT4* variant.

11. Outcome / Prognosis

- **Survival / mortality:** The dominant disorder is **non-lethal** with **normal life expectancy**; it does not directly increase mortality. (This contrasts sharply with recessive SERKAL, which is embryonic/perinatally lethal — [PMID: 18179883](#).)
 - **Morbidity / function:** Principal morbidities are **absolute uterine-factor infertility**, **sexual dysfunction** from vaginal aplasia, **androgen-excess manifestations** (hirsutism, acne), and **psychosocial morbidity**. MRKH systematic review: higher anxiety/depression and social insecurity; impaired sexual function; mental- but not physical-health QoL affected ([PMID: 35455528](#)).
 - **Complications:** Psychological distress; potential complications of neovaginal/surgical interventions; renal sequelae if significant renal anomaly present.
 - **Recovery potential:** The structural defect is not reversible, but **functional outcomes are good** with vaginal dilation/vaginoplasty (satisfactory sexual function) and ART; androgen excess is manageable pharmacologically.
 - **Prognostic factors:** Degree of uterine/renal involvement, severity of androgen excess, and access to multidisciplinary care and psychological support.
-

12. Treatment

Management is **symptomatic, multidisciplinary, and individualized** (NCIT: Supportive Care).

- **Neovagina creation (first-line non-surgical): Vaginal dilation** is the recommended initial approach. "Several non-surgical and surgical procedures have been reported for the creation of a functional neovagina; in general, non-surgical treatment should be the first initially pursued" ([PMID: 28434104](#)). Systematic review: "vaginal dilation is a viable initial treatment option for women with MRKH syndrome" ([PMID: 37994266](#)). NCIT: dilation therapy.

- **Surgical vaginoplasty:** Reserved for dilation failure/preference — e.g., Vecchietti (including minimally invasive single-port laparoscopic), McIndoe, or bowel vaginoplasty; a modified Vecchietti achieved a ~9 cm neovagina ([PMID: 33461758](#)). NCIT: vaginoplasty.
 - **Pharmacotherapy for hyperandrogenism: Anti-androgens** (spironolactone, cyproterone acetate) and combined estrogen–progestin or dermatologic therapy for hirsutism/acne; hormone management as indicated. (Mechanism-directed at androgen excess arising from CYP17A1/HSD3B2 de-repression.) NCIT: anti-androgen therapy.
 - **Fertility / ART:** Ovaries are functional, enabling **IVF with gestational surrogacy; uterine transplantation** is an emerging option restoring the possibility of gestation ([PMID: 28434104](#)); psychosocial outcomes of UTx candidacy are being characterized ([PMID: 39388911](#); [PMID: 39324432](#)).
 - **Psychological support:** Integral — counseling and mental-health support before/after interventions ([PMID: 28434104](#); [PMID: 39280286](#)).
 - **Advanced / experimental therapeutics:** No gene, cell, RNA, or targeted molecular therapies exist for this disorder; care is anatomic and endocrine.
 - **Pharmacogenomics:** None established.
-

13. Prevention

- **Primary prevention:** Not possible for this congenital genetic disorder. **Genetic counseling** for families with an identified *WNT4* variant informs reproductive decision-making.
 - **Secondary prevention: Early recognition** of primary amenorrhea with prompt imaging and endocrine/genetic work-up enables timely intervention and psychological support.
 - **Tertiary prevention:** Management of androgen excess, functional restoration (neovagina), fertility planning, and mental-health care to prevent complications and improve QoL.
 - **Genetic screening:** Cascade testing and, where a familial variant is known, options including **preimplantation/prenatal genetic testing** can be discussed; carrier/preconception counseling is relevant particularly to distinguish and avoid the recessive SERKAL scenario in consanguineous unions.
 - **Immunization / public-health / environmental measures:** Not applicable.
-

14. Other Species / Natural Disease

- **Taxonomy / orthologs:** *WNT4* is highly conserved; the mouse ortholog **Wnt4** (NCBI Gene 22417) is the principal comparative model. Human *WNT4* NCBI Gene 54361. Species: *Homo sapiens* (NCBI:txid9606), *Mus musculus* (NCBI:txid10090).
 - **Natural disease in other species:** No well-characterized spontaneous animal disease catalogued as an exact ortholog of this syndrome; the knowledge base is **engineered model organisms** rather than naturally occurring veterinary disease.
 - **Comparative biology:** The **Wnt4-null mouse** closely mirrors the human disorder — "Wnt-4-mutant females are masculinized-the Mullerian duct is absent while the Wolffian duct continues to develop" ([PMID: 9989404](#)) — evidencing strong **evolutionary conservation** of WNT4's role in Müllerian development and androgen repression.
 - **Transmission / zoonosis:** Not applicable (non-infectious, genetic).
-

15. Model Organisms

- **Model type: Mammalian — mouse (*Mus musculus*),** principally the **Wnt4 knockout (null)**. In-vitro steroidogenic cell models (e.g., OVCAR3) are used for functional variant assays.
- **Genetic models:** Constitutive **Wnt4 knockout**; pathway-related conditional β -catenin loss/gain-of-function models inform downstream biology (e.g., β -catenin controls nephron segmentation — [PMID: 31685872](#)).
- **Phenotype recapitulation: High.** The female Wnt4-null mouse reproduces the **core human features:** absent Müllerian duct ($\rightarrow$ uterovaginal aplasia), retained Wolffian duct, **virilization with ectopic ovarian testosterone synthesis** ([PMID: 9989404](#)), and **renal/nephron defects** ([PMID: 41358813](#)). The human disorder was described as "remarkably similar to that of female Wnt4-knockout mice" ([PMID: 15317892](#)).
- **Limitations:** Homozygous null mice model complete loss (closer to recessive SERKAL severity) whereas the human dominant disorder reflects **partial/heterozygous** loss; species differences in ovarian steroidogenesis regulation and the human-specific psychosocial dimension are not captured.
- **Applications / resources:** Dissecting WNT4 roles in Müllerian morphogenesis, ovarian androgen repression, and nephrogenesis. Resources: **MGI** (Wnt4), IMPC/IMSR.

Mechanistic Model / Interpretation (Synthesis)

The disorder is best understood as the consequence of **partial loss of a single dosage-sensitive developmental signal, WNT4**, whose two independent functions in the female gonad/tract are **simultaneously** disrupted. In normal development WNT4 (acting with RSPO1/ β -catenin and FOXL2) does two things that are both lost in patients: (1) it **drives Müllerian-duct formation**, and (2) it **keeps ovarian steroidogenic enzymes switched off**. Removing WNT4 therefore yields the otherwise paradoxical combination of a **missing uterus** and an **androgen-producing ovary** in the same 46,XX individual — a phenotype difficult to explain by any mechanism other than a shared upstream regulator. The convergence of human genetics (recurrent LOF missense alleles), in-vitro enzyme de-repression (CYP17A1/HSD3B2), and a faithful mouse knockout provides a rare, tightly-closed causal loop for a Mendelian reproductive disorder.

The **dosage axis** is central to nosology: **one** defective allele → viable dominant Müllerian aplasia + hyperandrogenism; **two** defective alleles → lethal recessive SERKAL (46,XX sex reversal with renal/adrenal/lung dysgenesis). This gene-dosage spectrum should be emphasized in any knowledge-base entry and in genetic counseling.

Dosage spectrum of WNT4 loss

- 0 LOF alleles → normal female development
- 1 LOF allele → Müllerian aplasia + hyperandrogenism (OMIM 158330, AD, viable)
- 2 LOF alleles → SERKAL syndrome (AR, 46,XX sex reversal, lethal)

Evidence Base

PMID	Role in this report	Evidence type
15317892	Index case defining the disorder; heterozygous <i>WNT4</i> LOF; "remarkably similar" to <i>Wnt4</i> -null mouse	Human clinical
18182450	p.L12P variant; functional de-repression of HSD3B2 + CYP17A1; low yield in unselected MRKH	Human + in vitro
21377155	p.A233T variant; abnormal CYP17A1 re-expression in OVCAR3; atypical MRKH cohort	Human + in vitro
9989404	<i>Wnt4</i> -null mouse: Müllerian absence + ectopic testosterone — defines dual mechanism	Model organism
18179883	Biallelic <i>WNT4</i> = SERKAL (AR, lethal); β -catenin degradation link	Human + mechanism
24641817	<i>WNT4</i> = only MRKH-associated mutation → atypical hyperandrogenic form; hyperandrogenemia 60.1%	Human clinical
28434104	MRKH review: ~1:4500; US/MRI diagnosis; management; <i>WNT4</i> in MRKH-like hyperandrogenism	Review
41636313	Epidemiology: MRKH ~10–15% of primary amenorrhea; renal anomalies ~15.5%	Human clinical
18617533	<i>WNT4</i> /RSPO1 canonical Wnt ligands promote ovarian fate, block testis	Model/mechanism
27481580	RSPO1/ <i>Wnt4</i> / β -catenin/ <i>Foxl2</i> in female genitalia & ovarian maintenance	Review
22251856	Ovarian somatic differentiation via FOXL2 + <i>Wnt4</i> / <i>Rspo1</i> β -catenin	Review
41358813	<i>Wnt4</i> -null mutants fail nephron formation (renal branch)	Model organism
35455528	QoL: anxiety/depression, sexual dysfunction; mental not physical QoL affected	Systematic review
37994266	Vaginal dilation viable first-line	Systematic review
39391975	EHR cohort: ~2× anxiety/depression in Müllerian aplasia	Human/EHR
33461758	Modified Vecchietti vaginoplasty (~9 cm neovagina)	Human clinical

PMID	Role in this report	Evidence type
24356390	<i>WNT4</i> excluded in MURCS — differential boundary	Human clinical
31685872	β -catenin in nephron segmentation (pathway context)	Model organism

Limitations and Knowledge Gaps

- 1. Very small human sample.** The *WNT4* hyperandrogenic subtype rests on a handful of patients/variants (p.E226G, p.L12P, p.A233T). Penetrance, precise prevalence, and genotype–phenotype correlation are not formally quantified.
- 2. ACMG classification uncertainty.** Beyond the functionally validated alleles, *WNT4* variants in MRKH are frequently VUS; the low diagnostic yield (~1/28 unselected MRKH) limits clinical actionability outside hyperandrogenic presentations.
- 3. Mechanistic resolution.** The CYP17A1/HSD3B2 de-repression is shown in cell lines; the precise ovarian cell type and in-vivo human steroidogenic flux are incompletely defined. No human transcriptomic/proteomic/metabolomic atlas of patient ovarian tissue exists.
- 4. Modifiers & epigenetics.** No validated modifier genes or disease-specific epigenetic marks; the basis of variable expressivity is unknown.
- 5. Model mismatch.** The mouse null best models the recessive (SERKAL-like) severity; a heterozygous/partial-LOF mouse recapitulating the human dominant syndrome is not well established.
- 6. Disorder-specific QoL/outcome data** are largely inferred from the broader MRKH literature rather than the *WNT4* subtype specifically.

Proposed Follow-up Experiments / Actions

- 1. Variant registry & functional reclassification.** Assemble all reported *WNT4* alleles in hyperandrogenic MRKH, apply uniform steroidogenic de-repression assays (CYP17A1/HSD3B2 reporter + androgen output), and reclassify under ACMG/AMP with PS3-level functional evidence.

2. **Heterozygous/partial-LOF mouse or organoid models.** Generate *Wnt4* point-knock-in (patient alleles) or hypomorphic alleles, and patient-iPSC-derived ovarian/Müllerian organoids, to model the dominant phenotype and quantify androgen output.
3. **Single-cell transcriptomics** of patient-derived (or model) ovarian somatic cells to pinpoint the de-repressed steroidogenic cell population and map the WNT4→ β -catenin→steroidogenic-gene axis.
4. **Targeted gene-panel strategy.** Prospectively test *WNT4* in hyperandrogenic MRKH presentations (enriched yield) and correlate androgen biochemistry with genotype.
5. **Prospective natural-history/QoL cohort** specific to *WNT4*-confirmed patients, capturing androgen trajectories, fertility outcomes (surrogacy/UTx), and validated QoL/psychological measures.
6. **Genetic-counseling protocol** explicitly incorporating the dominant-vs-recessive (SERKAL) dosage distinction, including preconception counseling in consanguineous families.

Evidence source legend: Human clinical (patient reports/cohorts), Model organism (mouse), In vitro (cell-line functional assays), Review (secondary synthesis). All mechanistic and clinical claims are cited to primary literature by PMID.
